

A₃-FORMALITY FOR PRO-2 DEMUSHKIN GROUPS

AMBRUS PÁL AND GEREON QUICK

ABSTRACT. We study a weak form of formality for differential graded algebras, called A_3 -formality, and show that the differential graded $\mathbb{F}_2$ -algebras of continuous cochains of all pro-2 Demushkin groups are A_3 -formal. We prove this result by an explicit computation of the Benson–Krause–Schwede canonical class using the classification of pro-2 Demushkin groups by Demushkin, Serre, and Labute. Compared to the case of odd primes, the new idea is to interpret the data of the canonical class as defining systems of higher Massey products.

1. INTRODUCTION

Let F be a field and let G_F denote its absolute Galois group. Let $\mathcal{C}^\bullet(G_F, \mathbb{F}_2)$ denote the differential graded algebra of inhomogeneous continuous cochains of G_F with coefficients in the constant discrete G_F -module $\mathbb{F}_2$. In [11], Hopkins and Wickelgren showed that, for F a local or global field, all triple Massey products of elements in $H^1(G_F, \mathbb{F}_2)$ vanish whenever they are defined. Since triple Massey products are the first obstruction to formality, Hopkins–Wickelgren therefore asked in [11, Question 1.4] whether $\mathcal{C}^\bullet(G_F, \mathbb{F}_2)$ is A_∞ -formal, i.e., the higher multiplication maps m_n on its minimal model are trivial for all $n \geq 3$. However, Positselski showed in [37, Section 9.11] and [38, §6] that $\mathcal{C}^\bullet(G_F, \mathbb{F}_2)$ is not A_∞ -formal in general for local fields. Then Harpaz–Wittenberg in [9, Example A.15] and more recently Merkurjev–Scavia in [23, Theorem 1.6] and [25, Theorem 1.3] provided examples which show that the second obstruction to formality is not trivial in general, i.e., not all fourfold Massey products are defined when the neighbouring cup-products vanish. For local fields, or more generally, for Demushkin groups, we may then ask the weaker question whether the differential graded algebra $\mathcal{C}^\bullet(G, \mathbb{F}_2)$ is A_3 -formal, i.e., whether the first higher multiplication map m_3 on its minimal model is trivial. A_3 -formality is much stronger than the vanishing of triple Massey products.

1.1. Demushkin groups and the main result. The purpose of this paper is to study A_3 -formality for $\mathcal{C}^\bullet(G, \mathbb{F}_2)$ for pro-2 Demushkin groups.

Definition 1.1. Let p be a prime number and let G be a pro- p -group. Then G is called a *Demushkin group* if $\dim_{\mathbb{F}_p} H^1(G, \mathbb{F}_p) < \infty$, $\dim_{\mathbb{F}_p} H^2(G, \mathbb{F}_p) = 1$, and the cup product $H^1(G, \mathbb{F}_p) \times H^1(G, \mathbb{F}_p) \rightarrow H^2(G, \mathbb{F}_p)$ is a non-degenerate bilinear form.

Demushkin groups are Poincaré groups of dimension two, and arise, for example, as pro- p completions of fundamental groups of compact surfaces. They also play a key role in number theory since the maximal pro- p quotients of absolute Galois

2020 *Mathematics Subject Classification.* 20J06, 12G05, 16E40, 20E18, 55S30.

Both authors were partially supported by RCN Project No. 313472 *Equations in Motivic Homotopy*, and the project *Pure Mathematics in Norway* funded by the Trond Mohn Foundation.

groups of local fields that contain a primitive p -th root of unity are Demushkin groups or trivial (see [3, 4], [41], [17], and Section 3.3). Demushkin groups are fundamental building blocks of the class of elementary type pro- p groups in the sense of Efrat. The only finite Demushkin group is $\mathbb{Z}/2$ and $\mathcal{C}^\bullet(\mathbb{Z}/2, \mathbb{F}_2)$ is known to be formal. We therefore consider from now on infinite Demushkin groups. Our main result in this paper is the following theorem, see Theorems 4.1, 5.1, and 6.1.

Theorem 1.2. *Let G be a pro-2 Demushkin group. Then $\mathcal{C}^\bullet(G, \mathbb{F}_2)$ is A_3 -formal.*

Before we outline the proof of Theorem 1.2 in Section 1.3 and comment on the new ideas used compared to the case of odd primes in [34], we first summarise the now complete characterisation of which Demushkin groups are A_3 -formal and will then describe the relation of our work to the Massey vanishing conjecture.

Let p be a prime number, and let G be a pro- p Demushkin group with $d := \dim_{\mathbb{F}_p} H^1(G, \mathbb{F}_p)$. Let G^{ab} denote the abelianisation of G . Then, by for example [32, Section 3.9 on page 232], either $G^{\text{ab}} \cong \mathbb{Z}_p^d$ or $G^{\text{ab}} \cong \mathbb{Z}_p/p^f \mathbb{Z}_p \times \mathbb{Z}_p^{d-1}$ with $f \geq 1$. In the latter case, we set $q := p^f$ and in the former case $q := 0$. The numbers d and q are invariants of G . Theorem 1.2 and [34, Theorem 1.2], then combine to the following result.

Theorem 1.3. *Let p be a prime number, and let G be a pro- p Demushkin group with invariants $d \geq 2$ and q . For $q \neq 3$, $\mathcal{C}^\bullet(G, \mathbb{F}_p)$ is A_3 -formal. For $q = p = 3$, $\mathcal{C}^\bullet(G, \mathbb{F}_3)$ is not A_3 -formal.*

1.2. Relation to the Massey vanishing conjecture of Mináč–Tân. Mináč and Tân conjectured in [29, Conjecture 1.6] that, for every field F and prime p , G_F satisfies n -Massey vanishing with respect to p , i.e., all n -fold Massey products of elements in $H^1(G_F, \mathbb{F}_p)$ vanish whenever they are defined. By the work of Matzri [20], Efrat–Matzri [8] and Mináč–Tân [28], all fields satisfy triple Massey vanishing with respect to all primes. For a more recent, independent proof, see [18, Section 6.2]. In [10], Harpaz–Wittenberg showed that number fields satisfy n -Massey vanishing with respect to all primes. More recently, Merkurjev–Scavia proved in [24] that all fields satisfy fourfold Massey vanishing with respect to $p = 2$. Other cases of the conjecture have been proven in [33], [35] and [39]. The vanishing of Massey products has concrete consequences for the structure of the Zassenhaus filtration of an absolute Galois group and thereby led to new examples of profinite groups which are not absolute Galois groups of a field (see for example [28, 29]).

In [27, Definition 4.5], Mináč and Tân also formulate the following related property. Let G be a profinite group and p be a prime number. Then G is said to have the *cup-defining n -fold Massey product property* (with respect to $\mathbb{F}_p$) if for every $\chi_1, \dots, \chi_n \in H^1(G, \mathbb{F}_p)$ with $0 = \chi_1 \cup \chi_2 = \chi_2 \cup \chi_3 = \dots = \chi_{n-1} \cup \chi_n$ the n -fold Massey product $\langle \chi_1, \dots, \chi_n \rangle$ is defined. For $n \geq 4$, this is a non-trivial condition, and, in [27, Remark 4.4], Mináč–Tân show that not all pro- p groups have the cup-defining n -fold Massey product property. In [27, Question 4.2], Mináč–Tân ask whether every Galois group of a maximal p -extension of a field F containing a primitive p -th root of unity has the cup-defining n -fold Massey product property with respect to $\mathbb{F}_p$ (see also [26, Section 8]). Moreover, in [27, Proposition 4.1], Mináč–Tân prove that pro- p Demushkin groups have the cup-defining n -fold Massey product property with respect to $\mathbb{F}_p$. Together with their work in [29], this implies that pro- p Demushkin groups have the following stronger property.

We say that a profinite group G satisfies *strong n -Massey vanishing* with respect to p if for every $\chi_1, \dots, \chi_n \in H^1(G, \mathbb{F}_p)$ with $0 = \chi_1 \cup \chi_2 = \dots = \chi_{n-1} \cup \chi_n$ the n -fold Massey product $\langle \chi_1, \dots, \chi_n \rangle$ is *defined and vanishes*. For $n \geq 4$, this is a strictly stronger condition than n -Massey vanishing. By the work of Mináč–Tân in [27, Proposition 4.1] and [28, Theorem 4.3], pro- p Demushkin groups satisfy strong n -Massey vanishing with respect to p and all $n \geq 3$. An independent proof that pro- p Demushkin groups satisfy strong n -Massey vanishing for all $n \geq 3$ was given by Pál–Szabó in [35, Theorem 3.5]. Moreover, by [19, Theorem 1], the absolute Galois groups of number fields which do not contain a primitive p -th root of unity satisfy strong n -Massey vanishing with respect to p for all $n \geq 3$. Strong vanishing of triple Massey products is a necessary condition for the A_3 -formality of $\mathcal{C}^\bullet(G, \mathbb{F}_p)$. In fact, A_3 -formality implies triple Massey vanishing and the cup-defining fourfold Massey product property. We note, however, that A_3 -formality is a significant strengthening of the vanishing of the Massey product obstructions since A_3 -formality requires that a specific element in the triple Massey product of elements in $H^\bullet(G, \mathbb{F}_p)$ vanishes and that defining systems of triple Massey products can be chosen compatibly.

1.3. Outline of the proof. We now give a brief outline of the proof of Theorem 1.2. While n -Massey vanishing for Demushkin groups is a direct consequence of the non-degeneracy of the cup product, showing A_3 -formality is much more involved. By the work of Kadeishvili, the differential graded algebra $\mathcal{C}^\bullet(G, \mathbb{F}_2)$ is A_3 -formal if and only if a certain *canonical class* γ_G in the Hochschild cohomology group $\mathrm{HH}^{3,-1}(H^\bullet(G, \mathbb{F}_2))$ is zero. The canonical class of a differential graded algebra has also been studied by Benson–Krause–Schwede in [1] in the context of the realizability of modules over Tate cohomology. The strategy for proving Theorem 1.2 is then to show that γ_G vanishes in $\mathrm{HH}^{3,-1}(H^\bullet(G, \mathbb{F}_2))$ for all pro-2 Demushkin groups. The task to compute the class γ_G in $\mathrm{HH}^{3,-1}(H^\bullet(G, \mathbb{F}_2))$ is simplified by the work of Mináč–Pasini–Quadrelli–Tân in [30, Theorem 5.2] who showed that the graded $\mathbb{F}_2$ -algebra $H^\bullet(G, \mathbb{F}_2)$ for a pro-2 Demushkin group is Koszul. A second key tool is Dwyer’s Theorem on unipotent representations and Massey products in group cohomology. A third key ingredient is the classification of pro-2 Demushkin groups by Demushkin, Serre, and Labute. A Demushkin group has a finite number of generators subject to a single relation. There are four different types of relations, each leading to a specific type of pro-2 Demushkin group. For each type, we compute the canonical class in a separate section. In each case, we first determine a basis for the relevant spaces R and K_3^3 in the Koszul complex of $H^\bullet(G, \mathbb{F}_2)$. Then we compute a sufficiently large part of the differential in the complex which computes the Hochschild cohomology of $H^\bullet(G, \mathbb{F}_2)$. Next, we construct the map $\kappa_3: K_3^3(H^\bullet(G, \mathbb{F}_2)) \rightarrow H^2(G, \mathbb{F}_2)$ which represents the canonical class. Finally, we compute the canonical class of G for each case.

The key new idea for $p = 2$ compared to the argument for odd primes in [34] is that we interpret κ_3 as higher Massey products. In [34], we computed triple Massey products in $H^1(G, \mathbb{F}_p)$ via Dwyer’s Theorem. Since there are enough relations in $K_3^3(H^\bullet(G, \mathbb{F}_p))$ for an odd prime p , we could then determine all values of κ_3 . For $p = 2$, however, there are not enough relations. Hence we use a different strategy and interpret the data in $K_3^3(H^\bullet(G, \mathbb{F}_2))$ as defining systems in certain n -fold Massey products for $n = 3, \dots, 8$, and then we compute the specific values of the corresponding element in $H^2(G, \mathbb{F}_2)$. After adding signs, the new method also

works for p odd, and hence we can use the methods in Section 4 to provide a new computation of the canonical class of a pro- p Demushkin for odd primes.

1.4. Contents. All vector spaces and algebras in this paper are over $\mathbb{F}_2$. In Section 2.1, we recall the definition of graded Hochschild cohomology groups of a graded algebra. In Section 2.2, we recall the definition of Koszul algebras. In Section 2.3, we define A_3 -formality for differential graded algebras and recall Massey products. Then we construct the canonical class, first for a general differential graded algebra in Section 2.4, and then for a differential graded algebra whose cohomology algebra is Koszul in Section 2.5. In Section 3.1, we recall continuous group cohomology, and in Section 3.2, we recall Dwyer's Theorem. In Section 3.3, we recall the classification of pro-2 Demushkin groups of Demushkin, Serre, and Labute, and we determine the algebra structure on $H^\bullet(G, \mathbb{F}_2)$ for pro-2 Demushkin groups. In Sections 4, 5, and 6, we compute the canonical class of pro-2 Demushkin groups for the three cases we need to consider according to the classification.

1.5. Acknowledgements. We thank Jan Mináč for helpful comments.

2. HOCHSCHILD COHOMOLOGY, A_3 -FORMALITY, AND MASSEY PRODUCTS

In this section, we recall the background for the computation of the canonical class of a differential graded algebra.

2.1. Graded Hochschild cohomology. Let A be a graded unital $\mathbb{F}_2$ -algebra. We recall that the bar resolution $B(A)$ of A is the non-negative chain complex of free graded A -bimodules given by $B_n(A) := A^{\otimes n+2}$ for $n \geq 0$. The differential $d_n: B_n(A) \rightarrow B_{n-1}(A)$ is given by

$$(1) \quad a_0 \otimes \cdots \otimes a_{n+1} \mapsto \sum_{i=0}^n a_0 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_{n+1}.$$

We write $A^e := A \otimes A^{\text{op}}$, where A^{op} denotes the opposite algebra. Note that $A^{\otimes n+2} \cong A^e \otimes A^{\otimes n}$ as a graded A -bimodule, and hence $B(A)$ indeed consists of free modules.

Proposition 2.1. *The bar resolution $B(A)$ is a free resolution of A as a graded A -bimodule.*

Proof. It suffices to show that the extended complex $\tilde{B}(A)$ is acyclic, where $\tilde{B}(A)$ is extended from $B(A)$ by adjoining $\tilde{B}_{-1}(A) := A$ in degree -1 via the multiplication map $\mu: A \otimes A \rightarrow A$. The map $h: \tilde{B}(A) \rightarrow \tilde{B}(A)$ of degree 1 given by

$$a_0 \otimes \cdots \otimes a_{n+1} \mapsto 1 \otimes a_0 \otimes \cdots \otimes a_{n+1}$$

is a contracting homotopy i.e., $dh + hd = 1$, since

$$dh(a_0 \otimes \cdots \otimes a_{n+1}) = a_0 \otimes \cdots \otimes a_{n+1} - hd(a_0 \otimes \cdots \otimes a_n). \quad \square$$

Definition 2.2. Let M and N be graded A -bimodules. We define $\underline{\text{Hom}}_A(M, N)$ as the graded $\mathbb{F}_2$ -vector space with degree s component given by A -linear graded maps $f: M \rightarrow N[s]$, where $N[s]$ is the graded A -module given by $N[s]^n = N^{s+n}$.

Definition 2.3. Let M be a graded A -bimodule. We define the Hochschild cohomology $\mathrm{HH}^{n,\bullet}(A, M)$ as the n th cohomology of the cochain complex

$$\underline{\mathrm{Hom}}_{A^e}(B(A), M)$$

of graded $\mathbb{F}_2$ -vector spaces with differential $\delta^n(f) = f \circ d_n$ where d_n is the differential of $B(A)$ in (1). When $M = A$ we will write $\mathrm{HH}(A) := \mathrm{HH}(A, A)$.

We note that the groups $\mathrm{HH}^{n,\bullet}(A, M)$ are equipped with a cohomological grading, and an internal grading induced by the grading of A and M . We can describe $\mathrm{HH}^{n,s}(A, M)$ more concretely as follows. Using the natural contracting isomorphism

$$\underline{\mathrm{Hom}}_{A^e}(A^e \otimes A^{\otimes n}, M) \cong \underline{\mathrm{Hom}}_{\mathbb{F}_2}(A^{\otimes n}, M)$$

we see that $\mathrm{HH}^{n,\bullet}(A, M)$ is isomorphic to the n th cohomology of the complex

$$(2) \quad \cdots \rightarrow \underline{\mathrm{Hom}}_{\mathbb{F}_2}(A^{\otimes n-1}, M) \xrightarrow{\partial} \underline{\mathrm{Hom}}_{\mathbb{F}_2}(A^{\otimes n}, M) \xrightarrow{\partial} \underline{\mathrm{Hom}}_{\mathbb{F}_2}(A^{\otimes n+1}, M) \rightarrow \cdots,$$

where the differentials are given by

$$\begin{aligned} \partial(f)(a_1 \otimes \cdots \otimes a_{n+1}) &= a_1 f(a_2 \otimes \cdots \otimes a_{n+1}) + \sum_{i=1}^n f(a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_{n+1}) \\ &\quad + f(a_1 \otimes \cdots \otimes a_n) a_{n+1}. \end{aligned}$$

Remark 2.4. By Proposition 2.1, we see that $\mathrm{HH}(A, M)$ computes the graded Ext modules $\underline{\mathrm{Ext}}_{A^e}(A, M)$. In particular, we can compute $\mathrm{HH}(A, M)$ using any free resolution of A as a graded A -bimodule.

2.2. Koszul algebras. Let V denote a finite-dimensional $\mathbb{F}_2$ -vector space, and let $T(V)$ denote its graded tensor algebra over $\mathbb{F}_2$. For $R \subseteq V \otimes V$, let (R) denote the two-sided ideal in $T(V)$ generated by R . We recall from [36] that a graded algebra of the form $T(V)/(R)$ is called a *quadratic algebra*. For any quadratic algebra A , we can define the following chain complex of free graded A -bimodules.

Definition 2.5. Let $A = T(V)/(R)$ be a quadratic algebra. Let $K_0^0 = \mathbb{F}_2$, $K_1^1 = V$, and, for $n \geq 2$, let

$$K_n^n := \bigcap_{i=1}^{n-1} V^{\otimes i-1} \otimes R \otimes V^{\otimes n-i-1} \subseteq V^{\otimes n}.$$

The Koszul complex $K(A^e, A)$ of A is defined as the non-negative chain complex of graded A -bimodules with

$$K_n(A^e, A) := A \otimes K_n^n \otimes A,$$

and differential d_n induced by the one in the bar resolution $B(A)$, i.e.,

$$d_n : a \otimes v_1 \otimes \cdots \otimes v_n \otimes b \mapsto av_1 \otimes v_2 \otimes \cdots \otimes v_n \otimes b + a \otimes v_1 \otimes \cdots \otimes v_n b.$$

Note that for the differential d_n in the Koszul complex, the middle terms of the bar construction differential in (1) vanish. This is because each product $v_i v_{i+1}$ in a middle term has its factors v_i, v_{i+1} in the space of relations R , so the product $v_i v_{i+1}$ vanishes in A , which is where the product in the expression of the differential is taking place.

Definition 2.6. A quadratic algebra A is called *Koszul* if its Koszul complex $K(A^e, A)$ is a resolution of A as a graded A -bimodule, i.e., if $H_n(K(A^e, A)) = 0$ for $n > 0$ and $H_0(K(A^e, A)) = A$.

Let A be a Koszul algebra. Then the natural inclusion $K(A^e, A) \hookrightarrow B(A)$ is a quasi-isomorphism. Hence we can compute $\mathrm{HH}(A)$ as the cohomology of the complex $\underline{\mathrm{Hom}}_{A^e}(K(A^e, A), A)$. We first observe that we have

$$K(A^e, A)_n = A \otimes K_n^n \otimes A \cong A^e \otimes K_n^n$$

as graded A -bimodules. Using the contracting isomorphism

$$\underline{\mathrm{Hom}}_{A^e}(A^e \otimes K_n^n, A) \cong \underline{\mathrm{Hom}}_{\mathbb{F}_2}(K_n^n, A)$$

of graded vector spaces we see that $\mathrm{HH}(A)$ can be computed as the cohomology of the following complex of graded vector spaces:

$$\cdots \rightarrow \underline{\mathrm{Hom}}_{\mathbb{F}_2}(K_{n-1}^{n-1}, A) \xrightarrow{\partial^{n-1}} \underline{\mathrm{Hom}}_{\mathbb{F}_2}(K_n^n, A) \xrightarrow{\partial^n} \underline{\mathrm{Hom}}_{\mathbb{F}_2}(K_{n+1}^{n+1}, A) \rightarrow \cdots$$

where the differential ∂^n is given by

$$\partial^n(f)(v_1 \otimes \cdots \otimes v_{n+1}) = v_1 f(v_2 \otimes \cdots \otimes v_{n+1}) + f(v_1 \otimes \cdots \otimes v_n) v_{n+1}.$$

In particular, the group $\mathrm{HH}^{3,-1}(A)$ is thus isomorphic to the cohomology of the complex

$$(3) \quad \mathrm{Hom}_{\mathbb{F}_2}(K_2^2, A^1) \xrightarrow{\partial^2} \mathrm{Hom}_{\mathbb{F}_2}(K_3^3, A^2) \xrightarrow{\partial^3} \mathrm{Hom}_{\mathbb{F}_2}(K_4^4, A^3).$$

2.3. A_3 -formality and Massey products. We now recall the definition of an A_∞ -algebra. For an introduction to the theory of A_∞ -algebras we refer to [16].

Definition 2.7. Let $\mathcal{A} = \bigoplus_{i \geq 0} \mathcal{A}^i$ be a non-negatively graded $\mathbb{F}_2$ -vector space with $\mathcal{A}^0 = \mathbb{F}_2$. Then $\mathcal{A}$ is called an A_∞ -algebra over $\mathbb{F}_2$ if, for all $i \geq 1$, there are graded $\mathbb{F}_2$ -linear maps $m_i: \mathcal{A}^{\otimes i} \rightarrow \mathcal{A}[2-i]$ such that, for all $n \geq 1$,

$$\sum_{r+s+t=n} m_{n-s+1}(\mathbf{1}^{\otimes r} \otimes m_s \otimes \mathbf{1}^{\otimes t}) = 0$$

as maps $\mathcal{A}^{\otimes n} \rightarrow \mathcal{A}$, where $\mathbf{1}$ denotes the identity map of $\mathcal{A}$.

Remark 2.8. For $n = 1, 2, 3$, the maps m_n satisfy the following identities: first, $m_1 m_1 = 0$, i.e., $(\mathcal{A}, m_1)$ is a cochain complex; second, $m_1 m_2 = m_2(m_1 \otimes \mathbf{1} + \mathbf{1} \otimes m_1)$ i.e., m_1 is a graded derivation with respect to the multiplication m_2 ; and

$$m_2(\mathbf{1} \otimes m_2 + m_2 \otimes \mathbf{1}) = m_1 m_3 + m_3(m_1 \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes m_1 \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes m_1),$$

i.e., m_2 is associative up to homotopy.

Example 2.9. Every graded $\mathbb{F}_2$ -algebra is an A_∞ -algebra with trivial m_1 and m_n for $n \geq 3$. Every differential graded algebra $(\mathcal{C}^\bullet, \delta, \cup)$ over $\mathbb{F}_2$ is an A_3 -algebra with $m_1 = \delta$, $m_2 = \cup$, and $m_n = 0$ for all $n \geq 3$.

Let $(\mathcal{C}^\bullet, \delta, \cup)$ be a differential graded algebra (DGA) over $\mathbb{F}_2$ with cohomology algebra $H^\bullet$. By the work of Kadeishvili [13, 14] (see also [15], [16], and [22]), one can equip $H^\bullet$ with the structure of an A_∞ -algebra $(H^\bullet, \{m_n\}_{n \geq 1})$ such that $m_1 = 0$ together with a quasi-isomorphism of A_∞ -algebras $(H^\bullet, \{m_n\}) \xrightarrow{\cong} (\mathcal{C}^\bullet, \delta, \cup)$. The A_∞ -algebra $(H^\bullet, \{m_n\})$ is called a *minimal model* of $(\mathcal{C}^\bullet, \delta, \cup)$. Since any two minimal models of $\mathcal{C}^\bullet$ are isomorphic as A_∞ -algebras, we speak of *the* minimal model from now on. A DGA is called A_∞ -formal, or *formal* as an A_∞ -algebra, if its minimal model can be chosen such that $m_n = 0$ for all $n \geq 3$. We now consider the following weaker notion. We refer to [34] for a more detailed discussion.

Definition 2.10. Let $\mathcal{C}^\bullet$ be a DGA. We say that $\mathcal{C}^\bullet$ is *A₃-formal* if its minimal model can be chosen such that $m_3 = 0$.

For a simple but non-trivial example of an *A₃-formal* DGA we refer to [5]. The notion of an *A_∞-algebra* is closely related to Massey products. We recall the definition of Massey products for degree one classes next.

Definition 2.11. Let $(\mathcal{C}^\bullet, \delta, \cup)$ be a DGA over $\mathbb{F}_2$ with cohomology algebra $H^\bullet$. For an integer $n \geq 2$, let $a_1, a_2, \dots, a_n$ be cohomology classes in H^1 . A *defining system* for the *n-fold Massey product* of $a_1, a_2, \dots, a_n$ is a set $\{a_{ij}\}$ of elements of $\mathcal{C}^1$ for $1 \leq i < j \leq n+1$ and $(i, j) \neq (1, n+1)$ such that

$$\delta(a_{i,j}) = \sum_{k=i+1}^{j-1} a_{i,k} \cup a_{k,j}$$

and, for each i , the class a_i is represented by $a_{i,i+1}$. We say that the *n-fold Massey product* of $a_1, a_2, \dots, a_n$ is *defined* if there exists a defining system. The *n-fold Massey product* $\langle a_1, a_2, \dots, a_n \rangle_{\{a_{i,j}\}}$ of $a_1, a_2, \dots, a_n$ with respect to the defining system $\{a_{i,j}\}$ is the cohomology class of

$$\sum_{k=2}^n a_{1,k} \cup a_{k,n+1}$$

in H^2 . Let $\langle a_1, a_2, \dots, a_n \rangle$ denote the subset of H^2 consisting of the *n-fold Massey products* of $a_1, a_2, \dots, a_n$ with respect to all defining systems. The set $\langle a_1, a_2, \dots, a_n \rangle$ is called the *n-fold Massey product* of $a_1, a_2, \dots, a_n$. We say that the *n-fold Massey product* of $a_1, a_2, \dots, a_n$ *vanishes* if the set $\langle a_1, a_2, \dots, a_n \rangle \subseteq H^2$ contains zero.

Example 2.12. For $n = 2$, the Massey product $\langle a_1, a_2 \rangle$ is just the cup product $a_1 \cup a_2$. Let $a_1, a_2, a_3 \in H^1$ such that $a_1 \cup a_2 = 0$ and $a_2 \cup a_3 = 0$. Then the *triple Massey product* $\langle a_1, a_2, a_3 \rangle$ is defined.

We recall the following special case from [2, Theorem C]. Let $a, b, c \in H^\bullet$ be cohomology classes such that $a \cup b = 0$ and $b \cup c = 0$. Then $m_3(a \otimes b \otimes c) \in \langle a, b, c \rangle$. This implies the following well-known fact.

Proposition 2.13. Let $\mathcal{C}^\bullet$ be a DGA with cohomology algebra $H^\bullet$. Let $a, b, c \in H^\bullet$ be cohomology classes such that $a \cup b = 0$ and $b \cup c = 0$. Assume that $\mathcal{C}^\bullet$ is *A₃-formal*. Then the triple Massey product $\langle a, b, c \rangle$ contains zero. $\square$

2.4. The canonical class. Let $(\mathcal{C}^\bullet, \delta, \cup)$ be a DGA with cohomology algebra $H^\bullet$. Let $(H^\bullet, \{m_n\})$ be the minimal model of $\mathcal{C}^\bullet$ as an *A_∞-algebra*. We note that we can construct the map $m_3: (H^\bullet)^{\otimes 3} \rightarrow H^\bullet[-1]$ as follows (see for example [1, Section 5], [22], [34]). We choose an $\mathbb{F}_2$ -linear graded map $f_1: H^\bullet \rightarrow \text{Ker } \delta$ which induces the identity when taking cohomology. Since f_1 is multiplicative on cohomology, we can find a graded $\mathbb{F}_2$ -linear map $f_2: H^\bullet \otimes H^\bullet \rightarrow \mathcal{C}^\bullet$ of degree -1 satisfying

$$\delta(f_2(a \otimes b)) = f_1(a \cup b) + f_1(a) \cup f_1(b).$$

Now we define a graded $\mathbb{F}_2$ -linear map $\Phi_3: (H^\bullet)^{\otimes 3} \rightarrow \mathcal{C}^\bullet[-1]$ by

$$\Phi_3(a \otimes b \otimes c) := f_1(a)f_2(b \otimes c) + f_2(a \otimes b)f_1(c) + f_2((ab) \otimes c + a \otimes (bc))$$

for all homogeneous elements $a, b, c \in H^\bullet$ where we write xy for the product $x \cup y$ to shorten the notation. We check that Φ_3 has image in the cocycles of $\mathcal{C}^\bullet$, and hence Φ_3 induces a graded map $[\Phi_3]: (H^\bullet)^{\otimes 3} \rightarrow H^\bullet[-1]$. We set $m_3 := [\Phi_3]$. By

[1, Proposition 5.4], m_3 is a cocycle in the complex (2). By [1, Corollary 5.7], the corresponding Hochschild cohomology class $[m_3] \in \mathrm{HH}^{3,-1}(H^\bullet)$ is independent of the choice of f_1 and f_2 , and it is called the *canonical class* of $\mathcal{C}^\bullet$ following Benson–Krause–Schwede who studied this class as an obstruction to the realizability of modules over Tate cohomology in [1]. We note that the canonical class is also sometimes called the *universal triple Massey product*, see for example [31]. The following result is a modified version of Kadeishvili’s theorem [14] (see also [34, Theorem 3.9], [5, Theorem 2.7], and [40, Theorem 4.7]).

Theorem 2.14. *Let $\mathcal{C}^\bullet$ be a DGA with canonical class $[m_3] \in \mathrm{HH}^{3,-1}(H^\bullet)$. Then $\mathcal{C}^\bullet$ is A_3 -formal if and only if $[m_3] = 0$.*

2.5. Canonical class for Koszul cohomology algebras. Let $(\mathcal{C}^\bullet, \delta, \cup)$ be a differential graded algebra over $\mathbb{F}_2$ with cohomology algebra $H^\bullet$. We now assume that $H^\bullet$ is a Koszul algebra. In this case, the canonical class of $\mathcal{C}^\bullet$ can be constructed in a simpler way as follows. Let $R \subset H^1 \otimes H^1$ denote the relations such that $H^\bullet = T(H^1)/(R)$. Let $\mathcal{Z}^1 = \ker \delta \subset \mathcal{C}^1$ denote the cocycles in degree one. Let $f_1: K_1^1(H^\bullet) = H^1 \rightarrow \mathcal{Z}^1$ be an $\mathbb{F}_2$ -linear map which induces the identity when taking cohomology. Let $f_2: K_2^2(H^\bullet) = R \rightarrow \mathcal{C}^1$ be an $\mathbb{F}_2$ -linear map such that $\delta f_2(a, b) = f_1(a) \cup f_1(b)$. We define the $\mathbb{F}_2$ -linear map

$$(4) \quad \Psi_3(a, b, c) := f_1(a) \cup f_2(b, c) + f_2(a, b) \cup f_1(c): K_3^3(H^\bullet) \rightarrow \mathcal{Z}^2$$

where, by slight abuse of notation, we allow (a, b) , (b, c) , and (a, b, c) to denote a sum of tensors in R and $K_3^3(H^\bullet)$ respectively. By a direct computation, we check that Ψ_3 has image in the cocycles of $\mathcal{C}^2$. Hence Ψ_3 induces an $\mathbb{F}_2$ -linear map

$$\kappa_3(a, b, c) = f_1(a) \cup f_2(b, c) + f_2(a, b) \cup f_1(c): K_3^3(H^\bullet) \rightarrow H^2.$$

In fact, the map κ_3 represents the canonical class of $\mathcal{C}^\bullet$:

Proposition 2.15. *The class of κ_3 in the cohomology of the complex (3) is the image of the canonical class of $\mathcal{C}^\bullet$ under the automorphism of $\mathrm{HH}^{3,-1}(H^\bullet)$ induced by the inclusion $K_3^3(H^\bullet) \hookrightarrow (H^1)^{\otimes 3}$. In particular, the differential graded algebra $\mathcal{C}^\bullet$ is A_3 -formal if and only if $[\kappa_3] = 0$.*

Proof. This follows from [34, Proposition 4.12] and Theorem 2.14. $\square$

3. DWYER’S CRITERION AND DEMUSHKIN GROUPS

We now specialise to differential graded algebras which arise from continuous group cohomology of profinite groups.

3.1. Continuous group cohomology and the canonical class. Let G be a profinite group, and let G^n denote the n -fold direct product of G with itself. Let $\mathcal{C}^n(G, \mathbb{F}_2)$ denote the $\mathbb{F}_2$ -vector space of continuous functions $G^n \rightarrow \mathbb{F}_2$ with respect to the discrete topology on $\mathbb{F}_2$ and the profinite topology on G . Following [42, §2.2] the differential $\delta: \mathcal{C}^n(G, \mathbb{F}_2) \rightarrow \mathcal{C}^{n+1}(G, \mathbb{F}_2)$, which is defined by

$$\begin{aligned} (\delta\varphi)(g_1, \dots, g_{n+1}) &= \varphi(g_2, \dots, g_{n+1}) + \sum_{i=1}^n \varphi(g_1, \dots, g_i g_{i+1}, \dots, g_{n+1}) \\ &\quad + \varphi(g_1, \dots, g_n), \end{aligned}$$

turns $\mathcal{C}^\bullet(G, \mathbb{F}_2)$ into a cochain complex whose cohomology $H^\bullet(G, \mathbb{F}_2)$ is the continuous cohomology of G with coefficients in the trivial G -module $\mathbb{F}_2$. In particular,

$H^1(G, \mathbb{F}_2)$ is the group of continuous group homomorphisms $G \rightarrow \mathbb{F}_2$. For every $\varphi \in \mathcal{C}^i(G, \mathbb{F}_2)$ and $\psi \in \mathcal{C}^j(G, \mathbb{F}_2)$, we define $\varphi \cup \psi \in \mathcal{C}^{i+j}(G, \mathbb{F}_2)$ by the formula:

$$(\varphi \cup \psi)(g_1, \dots, g_{i+j}) = \varphi(g_1, \dots, g_i) \cdot \psi(g_{i+1}, \dots, g_{i+j}).$$

This induces the cup product on cohomology which turns $H^\bullet(G, \mathbb{F}_2)$ into a graded $\mathbb{F}_2$ -algebra.

Definition 3.1. Let G be a pro-2 group. We say that G is *A₃-formal* if the differential graded $\mathbb{F}_2$ -algebra $\mathcal{C}^\bullet(G, \mathbb{F}_2)$ given by continuous cochains is *A₃-formal*. We write γ_G for the canonical class $\gamma_{\mathcal{C}^\bullet(G, \mathbb{F}_2)}$ in $\mathrm{HH}^{3,-1}(H^\bullet(G, \mathbb{F}_2))$ and call it the *canonical class of G* .

3.2. Dwyer's theorem on Massey products. We now recall from [6, Theorem 2.4] that the definition and vanishing of Massey products can be characterised as follows. Let $U_n(\mathbb{F}_2)$ denote the group of all upper triangular unipotent $(n \times n)$ -matrices with coefficients in $\mathbb{F}_2$. Let $Z_n(\mathbb{F}_2)$ denote the center of $U_n(\mathbb{F}_2)$, i.e., the subgroup of all matrices in $U_n(\mathbb{F}_2)$ with all off-diagonal entries being 0 except at position $(1, n)$. We write $\bar{U}_n(\mathbb{F}_2) = U_n(\mathbb{F}_2)/Z_n(\mathbb{F}_2)$.

Notation 3.2. We let $e_{ij}: U_n(\mathbb{F}_2) \rightarrow \mathbb{F}_2$ denote the projection to the (i, j) -coordinate.

The following result was proven by Dwyer in [6, Theorem 2.4].

Theorem 3.3 (Dwyer). *Let G be a profinite group. Let $a_1, \dots, a_n \in H^1(G, \mathbb{F}_2)$. There is a one-to-one correspondence $M \leftrightarrow \bar{\rho}_M$ between defining systems M for $\langle a_1, \dots, a_n \rangle$ and continuous group homomorphisms $\bar{\rho}_M: G \rightarrow \bar{U}_{n+1}(\mathbb{F}_2)$ such that $e_{i,i+1} \circ (\bar{\rho}_M) = a_i$ for $i = 1, \dots, n$, where we identify a_i with the corresponding continuous group homomorphism $G \rightarrow \mathbb{F}_2$. The correspondence is given by sending a defining system $M = \{a_{i,j}\}$ to the continuous group homomorphism $\bar{\rho}_M: G \rightarrow \bar{U}_{n+1}(\mathbb{F}_2)$ such that $e_{i,j} \circ \bar{\rho} = a_{i,j}$ for $1 \leq i < j \leq n+1$. Moreover, the element $\langle a_1, \dots, a_n \rangle_M \in H^2(G, \mathbb{F}_2)$ vanishes if and only if there is a continuous group homomorphism $\bar{\rho}_M: G \rightarrow \bar{U}_{n+1}(\mathbb{F}_2)$ making the following diagram commutative*

$$\begin{array}{ccccccc} & & & & G & & \\ & & & \swarrow \rho_M & \downarrow \bar{\rho}_M & & \\ 0 & \longrightarrow & Z_{n+1}(\mathbb{F}_2) & \longrightarrow & U_{n+1}(\mathbb{F}_2) & \longrightarrow & \bar{U}_{n+1}(\mathbb{F}_2) \longrightarrow 0. \end{array}$$

Example 3.4. Let $\chi_1, \chi_2 \in H^1(G, \mathbb{F}_2)$. Then we have $\chi_1 \cup \chi_2 = 0$ if and only if there is a continuous group homomorphism $\rho: G \rightarrow U_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho = \chi_1$ and $e_{2,3} \circ \rho = \chi_2$. In particular, the continuous map $\eta := e_{1,3} \circ \rho: G \rightarrow \mathbb{F}_2$ is a cochain in $\mathcal{C}^1(G, \mathbb{F}_2)$ such that $\delta\eta = \chi_1 \cup \chi_2$.

3.3. The classification of pro-2 Demushkin groups following Demushkin, Serre, and Labute. By the work of Demushkin [3, 4], Serre [41], and Labute [17], Demushkin groups are completely classified and pro-2 Demushkin groups can be characterised as follows. Let G be a pro-2 Demushkin group. For elements $x, y \in G$, let $[x, y] = x^{-1}y^{-1}xy$ denote the commutator. In the following, we use the convention $2^\infty := 0$. Let $d := \dim_{\mathbb{F}_2} H^1(G, \mathbb{F}_2)$, which is a finite number by definition of a Demushkin group. Then G is generated by elements $x_1, \dots, x_d$

subject to a single relation $r = r(x_1, \dots, x_d)$. The number d and the relation r satisfy one of the following characterisations:

(I) Demushkin[4]: $d \geq 2$ is even, $f \in \{2, 3, \dots\} \cup \{\infty\}$, and r is of the form

$$(5) \quad x_1^{2^f} [x_1, x_2] [x_3, x_4] \cdots [x_{d-1}, x_d] = 1;$$

(II) Serre [41]: $d \geq 3$ is odd, $f \in \{2, 3, \dots\} \cup \{\infty\}$, and r is of the form

$$(6) \quad x_1^2 x_2^{2^f} [x_2, x_3] \cdots [x_{d-1}, x_d] = 1;$$

(III) Labute [17]: $d \geq 2$ is even, $f \in \{2, 3, \dots\} \cup \{\infty\}$, and r is of the form

$$(7) \quad x_1^{2+2^f} [x_1, x_2] \cdots [x_{d-1}, x_d] = 1;$$

(IV) Labute [17]: $d \geq 4$ is even, $f \in \{2, 3, \dots\}$, and r is of the form

$$(8) \quad x_1^2 [x_1, x_2] x_3^{2^f} [x_3, x_4] \cdots [x_{d-1}, x_d] = 1.$$

Example 3.5. Let ℓ be an odd prime number. Let $G_{\mathbb{Q}_\ell}(2)$ be the maximal pro-2 quotient of the absolute Galois group of $\mathbb{Q}_\ell$, and let $q = 2^f$ be the maximum power of 2 such that $\mathbb{Q}_\ell$ contains a q -th root of unity. By [32, Proposition 7.5.9], $G_{\mathbb{Q}_\ell}(2)$ is pro-2 group with generators x_1, x_2 subject to the relation $1 = x_1^q [x_1, x_2]$. This group is of type I if $q \geq 4$, and it is of type III if $q = 2$.

Example 3.6. Following [41, Section 4], let $\mathbb{Q}_2$ denote the field of 2-adic numbers, and let K be a finite extension of $\mathbb{Q}_2$ of degree n . Let G be the maximal pro-2 quotient of the absolute Galois group of K . Let $q = 2^f$ denote the highest power of 2 such that K contains a primitive q -th root of unity. Assume $q \neq 1$. Then, by [41, Théorème 4.2], G is a pro-2 Demushkin group with $d = n + 2$. By [4] and [41, Corollaire 4.3], if $q \neq 2$, then G is defined by relation (5). By [41, Corollaire 4.4], if $q = 2$ and n is odd, then G is defined by relation (6). In particular, for $K = \mathbb{Q}_2$, we obtain the pro-2 Demushkin group G with generators x_1, x_2, x_3 subject to the single relation $1 = x_1^2 x_2^4 [x_2, x_3]$.

Example 3.7. We recall the following example provided by Labute in [17, §5]. Let $\mathbb{Q}_2$ denote the field of 2-adic numbers, and let $K = \mathbb{Q}_2(\sqrt{-2})$. Let G be the maximal pro-2 quotient of the absolute Galois group of K . By [17, Example below Theorem 9 on page 132], G is defined by relation (7) with $d = 4$, i.e., G is generated by x_1, x_2, x_3, x_4 subject to the relation $x_1^{2+4} [x_1, x_2] [x_3, x_4] = 1$. For other examples of this type, we refer to [32, Chapter 7.5, page 417].

Example 3.8. Let G be the pro-2 group with generators x_1, x_2, x_3 subject to the single relation $1 = x_1^2 [x_2, x_3]$. According to [12, Remark 5.5], it is not known whether this group arises as the maximal pro-2 quotient of an absolute Galois group.

In the following proofs and constructions we will frequently use the following observation, often without explicitly mentioning it.

Notation 3.9. For matrices M and N in $U_n(\mathbb{F}_2)$, we write $[M, N] := M^{-1}N^{-1}MN$.

Lemma 3.10. *Let G be a pro-2 group with generators $x_1, \dots, x_d$ subject to the single relation $r(x_1, \dots, x_d) = 1$. Let $M_1, \dots, M_d \in U_n(\mathbb{F}_2)$ and let I_n denote the $(n \times n)$ -identity matrix. If*

$$(9) \quad r(M_1, \dots, M_d) = I_n$$

in $\overline{U}_n(\mathbb{F}_2)$, then the assignment $\bar{\rho}: x_i \mapsto M_i$ for $x_1, \dots, x_d$ defines a continuous group homomorphism $\bar{\rho}: G \rightarrow \overline{U}_n(\mathbb{F}_2)$. Moreover, $\bar{\rho}$ can be lifted to a continuous group homomorphism $\rho: G \rightarrow U_n(\mathbb{F}_2)$ if and only if (9) holds in $U_n(\mathbb{F}_2)$.

Proof. This follows directly from the defining relation for G and the fact that $\overline{U}_n(\mathbb{F}_2)$ and $U_n(\mathbb{F}_2)$ are finite 2-groups. $\square$

Lemma 3.11. *Let G be a pro-2 Demushkin group with generators $x_1, \dots, x_d$. We let $\chi_i: G \rightarrow \mathbb{F}_2$ denote the group homomorphism determined by $\chi_i(x_j) = \delta_{ij}$, where δ_{ij} denotes the Kronecker symbol. For each i , we consider χ_i as an element in $H^1(G, \mathbb{F}_2)$. In $H^2(G, \mathbb{F}_2)$, we have the following relations:*

- If G is defined by relation (5), then $\chi_1 \cup \chi_2 \neq 0$, $\chi_1 \cup \chi_2 = \chi_{2i-1} \cup \chi_{2i} = \chi_{2i} \cup \chi_{2i-1}$ for $1 \leq i \leq d/2$, and $\chi_i \cup \chi_j = 0$ for all other cases. We have $H^2 = \mathbb{F}_2 \langle \chi_1 \cup \chi_2 \rangle$.
- If G is defined by relation (6), then $\chi_1 \cup \chi_1 \neq 0$, $\chi_1 \cup \chi_1 = \chi_{2i} \cup \chi_{2i+1} = \chi_{2i+1} \cup \chi_{2i}$ for $1 \leq i \leq (d-1)/2$, and $\chi_j \cup \chi_k = 0$ for all other cases. We have $H^2 = \mathbb{F}_2 \langle \chi_1 \cup \chi_1 \rangle$.
- If G is defined either by relation (7) or by relation (8), then $\chi_1 \cup \chi_1 \neq 0$, $\chi_1 \cup \chi_1 = \chi_{2i-1} \cup \chi_{2i} = \chi_{2i} \cup \chi_{2i-1}$ for $1 \leq i \leq d/2$, and $\chi_j \cup \chi_k = 0$ for all other cases. We have $H^2 = \mathbb{F}_2 \langle \chi_1 \cup \chi_1 \rangle$.

Proof. First, we assume that G is defined by relation (5). We write $q := 2^f$. Let $A_{10} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ and $A_{01} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ be matrices in $U_3(\mathbb{F}_2)$. We have $A_{10}^{2^n} = A_{01}^{2^n} = I_3$ in $U_3(\mathbb{F}_2)$ for all $n \geq 1$. Since $[A_{10}, A_{01}] = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, we have $A_{10}^q[A_{10}, A_{01}] = I_3$ in $\overline{U}_3(\mathbb{F}_2)$ but $A_{10}^q[A_{10}, A_{01}] \neq I_3$ in $U_3(\mathbb{F}_2)$. By Lemma 3.10, the assignment $x_1 \mapsto A_{10}$, $x_2 \mapsto A_{01}$, and $x_j \mapsto I_3$ for $j \neq 1, 2$ defines a continuous homomorphism $G \rightarrow \overline{U}_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho = \chi_1$, and $e_{2,3} \circ \rho = \chi_2$ which does not lift to a continuous homomorphism $G \rightarrow U_3(\mathbb{F}_2)$. By Dwyer's Theorem 3.3 and Example 3.4, this implies that $\chi_1 \cup \chi_2 \neq 0$ in $H^2(G, \mathbb{F}_2)$. Now assume that $i \neq j$ and $(i, j) \neq (2k-1, 2k), (2k, 2k-1)$. Since $[M, I_3] = [I_3, M] = I_3$ for every $M \in U_3(\mathbb{F}_2)$, the assignment $x_i \mapsto A_{10}$, $x_j \mapsto A_{01}$, and $x_s \mapsto I_3$ for $s \neq i, j$ defines, by Lemma 3.10, a continuous homomorphism $\rho_{i,j}: G \rightarrow U_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho_{i,j} = \chi_i$, and $e_{2,3} \circ \rho_{i,j} = \chi_j$. By Dwyer's Theorem 3.3 and Example 3.4, this implies $\chi_i \cup \chi_j = 0$ in $H^2(G, \mathbb{F}_2)$. Now let A_{11} denote the matrix $A_{11} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ in $U_3(\mathbb{F}_2)$. We have $A_{1,1}^n = \begin{pmatrix} 1 & n & \binom{n}{2} \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix}$ with $\binom{n}{2} = 0$ for $n = 0, 1$, and hence $A_{11}^q = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ in $U_3(\mathbb{F}_2)$ since either $q = 0$ or $q = 2^f$ with $f \geq 2$. Hence, by Lemma 3.10, the assignment $x_i \mapsto A_{11}$, and $x_j \mapsto I_3$ for $j \neq i$ defines a continuous homomorphism $\rho_{i,i}: G \rightarrow U_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho_{i,i} = \chi_i = e_{2,3} \circ \rho_{i,i}$. By Dwyer's Theorem 3.3 and Example 3.4, this implies $\chi_i \cup \chi_i = 0$ in $H^2(G, \mathbb{F}_2)$. Since $A_{10}^q[A_{10}, A_{01}][A_{01}, A_{10}] = I_3$ in $U_3(\mathbb{F}_2)$, the assignment $x_1 \mapsto A_{10}$, $x_2 \mapsto A_{01}$, $x_{2i-1} \mapsto A_{01}$, $x_{2i} \mapsto A_{10}$, and $x_j \mapsto I_3$ else defines a continuous homomorphism $\rho: G \rightarrow U_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho = \chi_1 + \chi_{2i}$, and $e_{2,3} \circ \rho = \chi_2 + \chi_{2i-1}$. By Dwyer's Theorem 3.3 and Example 3.4, this implies $(\chi_1 + \chi_{2i}) \cup (\chi_2 + \chi_{2i-1}) = 0$ in $H^2(G, \mathbb{F}_2)$. Since we already know $\chi_1 \cup \chi_{2i-1} = 0$ and $\chi_{2i} \cup \chi_2 = 0$, this implies $\chi_1 \cup \chi_2 = \chi_{2i} \cup \chi_{2i-1}$ in $H^2(G, \mathbb{F}_2)$. By switching the assignments for x_1, x_2, x_{2i-1} , and x_{2i} accordingly we get the remaining relation. This finishes the proof of the first case.

Now we assume that G is defined by relation (6). Since $A_{11}^2 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, we have $A_{11}^2 = I_3$ in $\overline{U}_3(\mathbb{F}_2)$ but $A_{11}^2 \neq I_3$ in $U_3(\mathbb{F}_2)$. By Lemma 3.10, the assignment $x_1 \mapsto A_{11}$ and $x_j \mapsto I_3$ for $j \neq 1$ defines a continuous homomorphism $G \rightarrow \overline{U}_3(\mathbb{F}_2)$ which does not lift to a continuous homomorphism $G \rightarrow U_3(\mathbb{F}_2)$. By Dwyer's Theorem 3.3 and Example 3.4, this implies that $\chi_1 \cup \chi_1 \neq 0$ in $H^2(G, \mathbb{F}_2)$. We have $A_{10}^{2^n} = A_{01}^{2^n} = I_3$ for all $n \geq 1$. Assume $(j, k) \neq (1, 1), (2i, 2i+1), (2i+1, 2i)$. By Lemma 3.10, the assignment $x_j \mapsto A_{10}$, $x_k \mapsto A_{01}$, and $x_r \mapsto I_3$ for $r \neq j, k$ defines a continuous homomorphism $\rho: G \rightarrow U_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho = \chi_j$, and $e_{2,3} \circ \rho = \chi_k$. By Dwyer's Theorem 3.3 and Example 3.4, this implies $\chi_j \cup \chi_k = 0$ in $H^2(G, \mathbb{F}_2)$. Since $[A_{10}, A_{01}] = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, we have $A_{11}^2[A_{10}, A_{01}] = I_3$ in $U_3(\mathbb{F}_2)$. By Lemma 3.10, the assignment $x_1 \mapsto A_{11}$, $x_{2i} \mapsto A_{10}$, $x_{2i+1} \mapsto A_{01}$, and $x_j \mapsto I_3$ for $j \neq 1, 2i, 2i+1$ defines a continuous homomorphism $\rho: G \rightarrow U_3(\mathbb{F}_2)$ such that $e_{1,2} \circ \rho = \chi_1 + \chi_{2i}$, and $e_{2,3} \circ \rho = \chi_1 + \chi_{2i+1}$. By Dwyer's Theorem 3.3 and Example 3.4, this implies $(\chi_1 + \chi_{2i}) \cup (\chi_1 + \chi_{2i+1}) = 0$ in $H^2(G, \mathbb{F}_2)$. Since we already know that $\chi_1 \cup \chi_{2i+1} = \chi_{2i} \cup \chi_1 = 0$ in $H^2(G, \mathbb{F}_2)$, this implies $\chi_1 \cup \chi_1 + \chi_{2i} \cup \chi_{2i+1} = 0$ in $H^2(G, \mathbb{F}_2)$. By switching the assignments for x_{2i} and x_{2i+1} , we get the assertion $\chi_1 \cup \chi_1 + \chi_{2i+1} \cup \chi_{2i} = 0$ in $H^2(G, \mathbb{F}_2)$, too. This finishes the second case.

The assertion for relations (7) and (8), follows as for relation (6) by replacing x_{2i} and x_{2i+1} with x_{2i-1} and x_{2i} respectively. See also Section 6.3. Here we note that every matrix $M \in U_3(\mathbb{F}_2)$ we used satisfies $M^4 = I_3$ in $U_3(\mathbb{F}_2)$. Hence an additional factor M^{2^f} in a matrix relation of the form (9), corresponding to the additional factor $x_1^{2^f}$ or $x_3^{2^f}$ in the defining relation for G , does not impose an obstruction for defining a continuous group homomorphism $G \rightarrow U_3(\mathbb{F}_2)$. $\square$

4. PRO-2 DEMUSHKIN GROUPS OF TYPE I

The goal of this section is to prove the following theorem.

Theorem 4.1. *Let $d \geq 2$ be an even number, and let $q = 2^f$ with $f \geq 2$, or $q = 0$. Let G be the pro-2 group with minimal set of generators $\{x_1, \dots, x_d\}$ satisfying the single relation*

$$1 = x_1^q[x_1, x_2][x_3, x_4] \cdots [x_{d-1}, x_d].$$

Then the canonical class of G is trivial, and G is A_3 -formal.

For the rest of this section, we let G be the pro-2 group as in Theorem 4.1. Let $\mathcal{C}^\bullet = \mathcal{C}^\bullet(G, \mathbb{F}_2)$ denote the complex of continuous inhomogeneous cochains, let $\mathcal{Z}^\bullet$ denote the cocycles, and let $H^\bullet = H^\bullet(G, \mathbb{F}_2)$ denote the corresponding cohomology algebra. We consider $\mathcal{C}^n$ and $\mathcal{Z}^n$ as $\mathbb{F}_2$ -vector spaces with addition and multiplication defined pointwise. We let $\chi_i: G \rightarrow \mathbb{F}_2$ denote the group homomorphism determined by $\chi_i(x_j) = \delta_{ij}$, where δ_{ij} denotes the Kronecker symbol. The collection $\{\chi_1, \dots, \chi_d\}$ forms a basis of H^1 . We write $J := \{1, \dots, d\}$, $J/2 := \{1, \dots, d/2\}$, and we set $\tilde{J}/2 := (J/2) \setminus \{1\} = \{2, \dots, d/2\}$. In Section 4.1, we determine the complex which computes the Hochschild cohomology of $H^\bullet$. In Section 4.2, we determine the effect of the corresponding differential. In Section 4.3, we give a concrete construction of a map which represents the canonical class of G . In Section 4.5, we finish the proof of Theorem 4.1.

4.1. The Koszul–Hochschild complex. By [30, Theorem 5.2], the graded $\mathbb{F}_2$ -cohomology algebra of G is a Koszul algebra. In this section, we determine a basis for the $\mathbb{F}_2$ -vector spaces R such that $H^\bullet = T(H^1)/R$ and of $K_3^3(H^\bullet)$.

Proposition 4.2. *The $\mathbb{F}_2$ -vector subspace $R \subset H^1 \otimes H^1$ such that $H^\bullet = T(H^1)/(R)$ admits the set $\mathbf{B}_2$ as a basis where*

$$\begin{aligned} \mathbf{B}_2 := & \{\chi_i \otimes \chi_j \text{ for all } (i, j) \notin \{(2k-1, 2k), (2k, 2k-1) : k \in J/2\}; \\ & \chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}, \chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \text{ for } i \in J/2\}. \end{aligned}$$

Proof. This follows from the description of the cup product on $H^\bullet(G, \mathbb{F}_2)$ in Lemma 3.11. Note that we have $\#\mathbf{B}_2 = d(d-1) + 2(d/2) - 1 = d^2 - 1$ as required for a basis by definition of a Demushkin group. $\square$

Proposition 4.3. *The set $\mathbf{B}_3 := \mathbf{S} \cup \mathbf{D} \cup \tilde{\mathbf{D}} \cup \mathbf{T}$ is a basis for the $\mathbb{F}_2$ -vector space $K_3^3(H^\bullet)$ where the sets $\mathbf{S}$ (single terms), $\mathbf{D}$, $\tilde{\mathbf{D}}$ (double sums), and $\mathbf{T}$ (triple sums) are*

$$\mathbf{S} = \left\{ \chi_i \otimes \chi_j \otimes \chi_k \text{ with } \begin{cases} i, k \neq j+1 & \text{if } j \text{ is odd} \\ i, k \neq j-1 & \text{if } j \text{ is even;} \end{cases} \right\},$$

$$\begin{aligned} \mathbf{D} = & \{\chi_k \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}), \\ & (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_k \text{ for } i \in J/2, k \neq 2i-1, 2i, \\ & \chi_k \otimes (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}), \\ & (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_k \text{ for } i \in \tilde{J}/2, k \neq 1, 2, 2i-1, 2i\} \end{aligned}$$

$$\begin{aligned} \tilde{\mathbf{D}} = & \{\chi_1 \otimes (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}), \chi_{2i} \otimes (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}), \\ & \chi_2 \otimes (\chi_2 \otimes \chi_1 + \chi_{2i-1} \otimes \chi_{2i}), \chi_{2i-1} \otimes (\chi_2 \otimes \chi_1 + \chi_{2i-1} \otimes \chi_{2i}), \\ & (\chi_2 \otimes \chi_1 + \chi_{2i-1} \otimes \chi_{2i}) \otimes \chi_1, (\chi_2 \otimes \chi_1 + \chi_{2i-1} \otimes \chi_{2i}) \otimes \chi_{2i}, \\ & (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_2, (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_{2i-1} \text{ for } i \in \tilde{J}/2\}. \end{aligned}$$

$$\begin{aligned} \mathbf{T} = & \{\chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i-1} + \chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i-1} \otimes \chi_{2i-1} \otimes \chi_{2i}, \\ & \chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i} \otimes \chi_{2i-1} \text{ for } i \in J/2\}. \end{aligned}$$

Proof. Proposition 4.2 shows that R has the same form as for pro- p Demushkin groups for an odd prime p . Since $K_3^3(H^\bullet)$ only depends on the relations in H^2 , the same arguments as in [34, Lemma 7.2] then prove the assertion that $\mathbf{B}_3$ is a basis of $K_3^3(H^\bullet)$. (Note that we explain the argument in the other cases below.) $\square$

4.2. The differential in the Koszul–Hochschild complex. Now, we analyse the differential $\partial: \text{Hom}_{\mathbb{F}_2}(K_2^2(H^\bullet), H^1) \rightarrow \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ in the complex (3) which computes the Hochschild cohomology of $H^\bullet$. For any $\sigma \in \text{Hom}_{\mathbb{F}_2}(K_2^2(H^\bullet), H^1)$, $\partial\sigma$ vanishes on elements in $\mathbf{T}$ and on elements in the set $\Delta_{\mathbf{S}} := \{\chi_i \otimes \chi_i \otimes \chi_i : i \in J\} \subseteq \mathbf{S}$. In particular, for $d = 2$, the differential is trivial. For $d \geq 4$, however, ∂ is non-trivial as we now show. First, we note that it is easy to see that for every element (χ_a, χ_b, χ_c) in $\mathbf{S} \setminus \Delta_{\mathbf{S}} \cup \mathbf{D}$, there is a $\sigma \in \mathbf{B}_2^2$ such that $\partial\sigma(\chi_a, \chi_b, \chi_c) \neq 0$. The same $\partial\sigma$ may be non-trivial on several elements of $\mathbf{B}_3^3$ simultaneously. Since we will show in Section 4.5 that the map which represents the canonical class of G vanishes on the sets $\mathbf{S}$ and $\mathbf{D}$, we only compute the effect on the set $\tilde{\mathbf{D}}$.

Notation 4.4. We will often write $\chi_{i,j}$ for $\chi_i \otimes \chi_j \in H^1 \otimes H^1$ and $\chi_{i,j,k}$ for $\chi_i \otimes \chi_j \otimes \chi_k \in H^1 \otimes H^1 \otimes H^1$ when it makes formulas easier to fit into the text.

Notation 4.5. Let $\chi_{i,j} \in \mathbf{B}_2$, i.e., $(i,j) \neq (2k-1, 2k), (2k, 2k-1)$. For $s \in J$, we write $\sigma_{i,j}^s$ for the unique $\mathbb{F}_2$ -linear map $R \rightarrow H^1$ which sends $\chi_{i,j}$ to χ_s and every other element of $\mathbf{B}_2$ to 0. For $\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1} \in \mathbf{B}_2$, $i \in J/2$, and $s \in J$, we write $\sigma_{1,2'+2i,2i-1}^s$ for the unique $\mathbb{F}_2$ -linear map $R \rightarrow H^1$ which sends $\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}$ to χ_s and every other element of $\mathbf{B}_2$ to 0. Similarly, for $\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \in \mathbf{B}_2$, we write $\sigma_{2i-1,2i'+2i,2i-1}^s$ for the unique $\mathbb{F}_2$ -linear map $R \rightarrow H^1$ which sends $\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}$ to χ_s and every other element of $\mathbf{B}_2$ to 0.

Notation 4.6. For an element $(\chi_a, \chi_b, \chi_c) \in \mathbf{B}_3$, we write $(\chi_a, \chi_b, \chi_c)^*$ for the unique $\mathbb{F}_2$ -linear map in $\text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ such that $(\chi_a, \chi_b, \chi_c)^*(\chi_a, \chi_b, \chi_c) \neq 0$ and $(\chi_a, \chi_b, \chi_c)^*$ sends all other elements in $\mathbf{B}_3$ to 0. By slight abuse of terminology, using the isomorphism $\text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2) \cong \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), \mathbb{F}_2)$ induced by the isomorphism $H^2 \cong \mathbb{F}_2$, we refer to $(\chi_a, \chi_b, \chi_c)^*$ as the *dual of* (χ_a, χ_b, χ_c) . We write $\mathbf{B}_3^*$ for the set of $(\chi_a, \chi_b, \chi_c)^*$ for all elements in $\mathbf{B}_3$.

Lemma 4.7. Assume $d \geq 4$ and let $i \in \tilde{J}/2$. Then

$$\begin{aligned} (\chi_{2,1,2i} + \chi_{2i-1,2i,2i})^* &= \partial \sigma_{2i-1,2i'+2i,2i-1}^{2i-1}, \\ (\chi_{2i-1,2,1} + \chi_{2i-1,2i-1,2i})^* &= \partial \sigma_{2i-1,2i'+2i,2i-1}^{2i}, \\ (\chi_{2i,1,2} + \chi_{2i,2i,2i-1})^* &= \partial \sigma_{2i-1,2i'+2i,2i-1}^{2i-1} + \partial \sigma_{1,2'+2i,2i-1}^{2i-1}, \\ \text{and } (\chi_{1,2,2i-1} + \chi_{2i,2i-1,2i-1})^* &= \partial \sigma_{2i-1,2i'+2i,2i-1}^{2i} + \partial \sigma_{1,2'+2i,2i-1}^{2i}. \end{aligned}$$

Moreover, we have

$$\begin{aligned} \sum_{j \in \tilde{J}/2} (\chi_{2,2,1} + \chi_{2,2j-1,2j})^* &= \partial \sigma_{1,2'+2,1}^1, \quad \sum_{j \in \tilde{J}/2} (\chi_{2,1,1} + \chi_{2j-1,2j,1})^* = \partial \sigma_{1,2'+2,1}^2, \\ \sum_{j \in \tilde{J}/2} (\chi_{1,2,2} + \chi_{2j,2j-1,2})^* &= \partial \sigma_{1,2'+2,1}^1 + \sum_{j \in \tilde{J}/2} \partial \sigma_{1,2'+2j,2j-1}^1, \end{aligned}$$

and

$$\sum_{j \in \tilde{J}/2} (\chi_{1,1,2} + \chi_{1,2j,2j-1})^* = \partial \sigma_{1,2'+2,1}^2 + \sum_{j \in \tilde{J}/2} \partial \sigma_{1,2'+2j,2j-1}^2.$$

Proof. First, we consider $\partial \sigma_{2i-1,2i'+2i,2i-1}^{2i}$ and compute

$$\begin{aligned} &\partial \sigma_{2i-1,2i'+2i,2i-1}^{2i} (\chi_{2i-1,2,1} + \chi_{2i-1,2i-1,2i}) \\ &= \chi_{2i-1} \cup \sigma_{2i-1,2i'+2i,2i-1}^{2i} (\chi_{2,1} + \chi_{2i-1,2i}) \\ &\quad + \sigma_{2i-1,2i'+2i,2i-1}^{2i} (\chi_{2i-1,2}) \cup \chi_1 + \sigma_{2i-1,2i'+2i,2i-1}^{2i} (\chi_{2i-1,2i-1}) \cup \chi_{2i}. \end{aligned}$$

By definition of $\sigma_{2i-1,2i+2i,2i-1}^{2i}$, we know that $\sigma_{2i-1,2i+2i,2i-1}^{2i}(\chi_{2i-1,2}) = 0$ and $\sigma_{2i-1,2i+2i,2i-1}^{2i}(\chi_{2i-1,2i-1}) = 0$. Moreover, we have

$$(10) \quad \chi_{2,1} + \chi_{2i-1,2i} = (\chi_{1,2} + \chi_{2,1}) + (\chi_{2i-1,2i} + \chi_{2i,2i-1}) + (\chi_{1,2} + \chi_{2i,2i-1}).$$

By definition of $\sigma_{2i-1,2i+2i,2i-1}^{2i}$, we have $\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{1,2} + \chi_{2,1}) = 0$ and $\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{1,2} + \chi_{2i,2i-1}) = 0$, and hence

$$\begin{aligned} & \partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2i-1,2,1} + \chi_{2i-1,2i-1,2i}) \\ &= \chi_{2i-1} \cup \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2i-1,2i} + \chi_{2i,2i-1}) = \chi_{2i-1} \cup \chi_{2i} \neq 0. \end{aligned}$$

It remains to show that $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}$ vanishes on all other elements of $\mathbf{B}_3$. For $\chi_{a,b,c} \in \mathbf{S}$, we have $\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{a,b}) = 0 = \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{b,c})$ and hence $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{a,b,c}) = 0$. Now let $b \in J/2$ and $a \in J$ with $a \neq 2b-1, 2b$. Then

$$\begin{aligned} & \partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{a,2b-1,2b} + \chi_{a,2b,2b-1}) \\ &= \chi_a \cup \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2b-1,2b} + \chi_{2b,2b-1}) \\ & \quad + \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{a,2b-1}) \cup \chi_{2b} + \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{a,2b}) \cup \chi_{2b-1}. \end{aligned}$$

By definition of $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}$, the second and third summands are zero, and we have $\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2b-1,2b} + \chi_{2b,2b-1}) \neq 0$ if and only if $b = i$ by Lemma 3.11. However, the cup-product $\chi_a \cup \chi_{2i}$ is non-zero if and only if $a = 2i-1$. But we have $a \neq 2b-1$ by assumption. This shows $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{a,2b-1,2b} + \chi_{a,2b,2b-1}) = 0$. Similarly, we get $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2b-1,2b,a} + \chi_{2b,2b-1,a}) = 0$. We can check that $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}$ vanishes on the remaining elements of $\mathbf{D}$. For another example of an element in $\tilde{\mathbf{D}}$, we compute, for $b \in \tilde{J}/2$,

$$\begin{aligned} & \partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2,1,1} + \chi_{2b-1,2b,1}) \\ &= \chi_2 \cup \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{1,1}) + \chi_{2b-1} \cup \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2b,1}) \\ & \quad + \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2,1} + \chi_{2b-1,2b}) \cup \chi_1 \\ &= \sigma_{2i-1,2i'+2i,2i-1}^{2i}(\chi_{2b-1,2b} + \chi_{2b,2b-1}) \cup \chi_1 \end{aligned}$$

where we use again (10). The last cup product is non-zero if and only if $b = i$ and $i = 1$ by Lemma 3.11. However, we have $b \geq 2$ by assumption. Similar computations show that $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}$ vanishes on the remaining elements in $\tilde{\mathbf{D}}$ as well. Since $\partial\sigma$ vanishes on $\mathbf{T}$ for all σ , this proves the assertion for $\partial\sigma_{2i-1,2i'+2i,2i-1}^{2i}$.

Second, we consider $\partial\sigma_{1,2'+2,1}^1$. Let $j \in \tilde{J}/2$. By definition of $\partial\sigma_{1,2'+2,1}^1$, we have $\sigma_{1,2'+2,1}^1(\chi_{2,2}) = 0$ and $\sigma_{1,2'+2,1}^1(\chi_{2,2j-1}) = 0$. Hence we get

$$\begin{aligned} & \partial\sigma_{1,2'+2,1}^1(\chi_{2,2,1} + \chi_{2,2j-1,2j}) \\ &= \chi_2 \cup \sigma_{1,2'+2,1}^1(\chi_{2,1} + \chi_{2j-1,2j}) + \sigma_{1,2'+2,1}^1(\chi_{2,2}) \cup \chi_1 + \sigma_{1,2'+2,1}^1(\chi_{2,2j-1}) \cup \chi_{2j} \\ &= \chi_2 \cup \sigma_{1,2'+2,1}^1(\chi_{2,1} + \chi_{2j-1,2j}). \end{aligned}$$

Now we use again (10). We have $\sigma_{1,2'+2,1}^1(\chi_{2j-1,2j} + \chi_{2j,2j-1}) = 0$ and $\sigma_{1,2'+2,1}^1(\chi_{1,2} + \chi_{2j,2j-1}) = 0$ since $j \geq 2$ by assumption. Thus, for every $j \in \tilde{J}/2$,

$$\partial\sigma_{1,2'+2,1}^1(\chi_{2,2,1} + \chi_{2,2j-1,2j}) = \chi_2 \cup \sigma_{1,2'+2,1}^1(\chi_{1,2} + \chi_{2,1}) = \chi_2 \cup \chi_1 \neq 0.$$

It remains to show that $\partial\sigma_{1,2'+2,1}^1$ vanishes on all other elements of $\mathbf{B}_3^3$. For $\chi_{a,b,c} \in \mathbf{S}$, we have $\sigma_{1,2'+2,1}^1(\chi_{a,b}) = 0 = \sigma_{1,2'+2,1}^1(\chi_{b,c})$ and hence $\partial\sigma_{1,2'+2,1}^1(\chi_{a,b,c}) = 0$.

Now let $b \in J/2$ and $a \in J$ with $a \neq 2b - 1, 2b$. Then

$$\begin{aligned} & \partial\sigma_{1,2'+2,1}^1(\chi_{a,2b-1,2b} + \chi_{a,2b,2b-1}) \\ &= \chi_a \cup \sigma_{1,2'+2,1}^1(\chi_{2b-1,2b} + \chi_{2b,2b-1}) \\ &+ \sigma_{1,2'+2,1}^1(\chi_{a,2b-1}) \cup \chi_{2b} + \sigma_{1,2'+2,1}^1(\chi_{a,2b}) \cup \chi_{2b-1}. \end{aligned}$$

By definition of $\partial\sigma_{1,2'+2,1}^1$, the second and third summands are zero, and we have $\sigma_{1,2'+2,1}^1(\chi_{2b-1,2b} + \chi_{2b,2b-1}) \neq 0$ if and only if $b = 1$. However, the cup-product $\chi_a \cup \chi_1$ is non-zero if and only if $a = 2$ by Lemma 3.11. But we have $a \neq 2b = 2$ by assumption. This shows $\partial\sigma_{1,2'+2,1}^1(\chi_{a,2b-1,2b} + \chi_{a,2b,2b-1}) = 0$. Similarly, we get $\partial\sigma_{1,2'+2,1}^1(\chi_{2b-1,2b,a} + \chi_{2b,2b-1,a}) = 0$. Now we assume $b \in \tilde{J}/2$. Then $\partial\sigma_{1,2'+2,1}^1$ sends both $\chi_{a,1,2} + \chi_{a,2b,2b-1}$ and $\chi_{1,2,a} + \chi_{2b,2b-1,a}$ to zero, since $\sigma_{1,2'+2,1}^1(\chi_{1,2} + \chi_{2b,2b-1}) = 0$ for $b \geq 2$. It remains to show that $\partial\sigma_{1,2'+2,1}^1$ vanishes on the other elements in $\tilde{\mathbf{D}}$. Then, by relation (10), we get $\sigma_{1,2'+2,1}^1(\chi_{2,1} + \chi_{2b-1,2b}) = \chi_1$. However, since $\chi_1 \cup \chi_1 = \chi_1 \cup \chi_{2b} = \chi_{2b-1} \cup \chi_1 = 0$, we conclude that $\partial\sigma_{1,2'+2,1}^1$ vanishes on the elements $(\chi_{2,1,1} + \chi_{2b-1,2b,1})$, $(\chi_{2,1,2b} + \chi_{2b-1,2b,2b})$, and $(\chi_{2b-1,2,1} + \chi_{2b-1,2b-1,2b})$ in $\tilde{\mathbf{D}}$. Since $\sigma_{1,2'+2,1}^1(\chi_{1,2} + \chi_{2b,2b-1}) = 0$ for $b \geq 2$, we get that $\partial\sigma_{1,2'+2,1}^1$ also sends the elements $\chi_{1,1,2} + \chi_{1,2b,2b-1}$, $\chi_{1,2,2} + \chi_{2b,2b-1,2}$, $\chi_{2b,1,2} + \chi_{2b,2b,2b-1}$, and $\chi_{1,2,2b-1} + \chi_{2b,2b-1,2b-1}$ in $\tilde{\mathbf{D}}$ to zero. This proves that $\partial\sigma_{1,2'+2,1}^1$ is non-zero exactly on all elements of the form $(\chi_{2,2,1} + \chi_{2,2j-1,2j})^*$ for all $j \in \tilde{J}/2$. This proves the claim for $\partial\sigma_{1,2'+2,1}^1$. The remaining assertions of the lemma follow from similar computations. $\square$

Proposition 4.8. *Let $\tilde{\mathbf{D}}_\Sigma^* := \tilde{\mathbf{D}}_1^* \oplus \tilde{\mathbf{D}}_2^*$ denote the subspace of $\text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ given by the sum of*

$$\begin{aligned} \tilde{\mathbf{D}}_1^* := \text{Span}_{\mathbb{F}_2} \{ & (\chi_{2,1,2i} + \chi_{2i-1,2i,2i})^*, (\chi_{2i-1,2,1} + \chi_{2i-1,2i-1,2i})^*, \\ & (\chi_{2i,1,2} + \chi_{2i,2i,2i-1})^*, (\chi_{1,2,2i-1} + \chi_{2i,2i-1,2i-1})^* : i \in \tilde{J}/2 \} \end{aligned}$$

and

$$\begin{aligned} \tilde{\mathbf{D}}_2^* := \text{Span}_{\mathbb{F}_2} \{ & \sum_{j \in \tilde{J}/2} (\chi_{2,2,1} + \chi_{2,2j-1,2j})^*, \sum_{j \in \tilde{J}/2} (\chi_{2,1,1} + \chi_{2j-1,2j,1})^*, \\ & \sum_{j \in \tilde{J}/2} (\chi_{1,2,2} + \chi_{2j,2j-1,2})^*, \sum_{j \in \tilde{J}/2} (\chi_{1,1,2} + \chi_{1,2j,2j-1})^* \}. \end{aligned}$$

Then $\tilde{\mathbf{D}}_\Sigma^* \subset \text{Im}(\partial)$. In particular, if $\kappa \in \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ sends $\mathbf{S}$, $\mathbf{D}$, and $\mathbf{T}$ to zero, and has the same values on $(\chi_{2,2,1} + \chi_{2,2i-1,2i})$, $(\chi_{2,1,1} + \chi_{2i-1,2i,1})$, $(\chi_{1,2,2} + \chi_{2i,2i-1,2})$, and $(\chi_{1,1,2} + \chi_{1,2i,2i-1})$, respectively, for all $i \in \tilde{J}/2$, then κ is a coboundary and $[\kappa] = 0$ in $\text{HH}^{3,-1}(H^\bullet)$.

Proof. This follows directly from Lemma 4.7. $\square$

Corollary 4.9. *Let $\kappa \in \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$, and assume that κ sends $\mathbf{S}$, $\mathbf{D}$, and $\mathbf{T}$ to zero, and has the same values on $(\chi_{2,2,1} + \chi_{2,2i-1,2i})$, $(\chi_{2,1,1} + \chi_{2i-1,2i,1})$, $(\chi_{1,2,2} + \chi_{2i,2i-1,2})$, and $(\chi_{1,1,2} + \chi_{1,2i,2i-1})$, respectively. Then κ is a coboundary and $[\kappa] = 0$ in $\text{HH}^{3,-1}(H^\bullet)$.*

Proof. This follows directly from Proposition 4.8. $\square$

4.3. Construction of the canonical class. We will now construct the canonical class of G . Let $\text{Hom}(G, \mathbb{F}_2)$ denote the $\mathbb{F}_2$ -vector space of continuous group homomorphisms. We choose f_1 to be the identity $\text{Hom}(G, \mathbb{F}_2) = H^1 \rightarrow \mathcal{Z}^1 = \text{Hom}(G, \mathbb{F}_2)$ and omit it from the notation. We construct an $\mathbb{F}_2$ -linear map $f_2: R \rightarrow \mathcal{C}^1$ by defining it on each element of $\mathbf{B}_2$ and then extend it $\mathbb{F}_2$ -linearly.

Let $A_{10} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, $A_{01} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$, and $A_{11} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ denote the matrices used in the proof of Lemma 3.11. Let $i \neq j$ and $(i, j) \neq (2k-1, 2k), (2k, 2k-1)$. As in the proof of Lemma 3.11, we define the continuous homomorphism $\rho_{i,j}: G \rightarrow U_3(\mathbb{F}_2)$ by the assignment $x_i \mapsto A_{10}$, $x_j \mapsto A_{01}$, and $x_s \mapsto I_3$ for $s \neq i, j$. We define the continuous map $f_2(\chi_i \otimes \chi_j): G \rightarrow \mathbb{F}_2$ by setting $f_2(\chi_i \otimes \chi_j) := e_{1,3} \circ \rho_{i,j}$. We define the continuous homomorphism $\rho_{i,i}: G \rightarrow U_3(\mathbb{F}_2)$ by the assignment $x_i \mapsto A_{11}$, and $x_j \mapsto I_3$ for $j \neq i$. We define the continuous map $f_2(\chi_i \otimes \chi_i): G \rightarrow \mathbb{F}_2$ by setting $f_2(\chi_i \otimes \chi_i) := e_{1,3} \circ \rho_{i,i}$. Now let $i \in J/2$. We would like to define continuous map $\eta_{2i-1, 2i'+2i, 2i-1}: G \rightarrow \mathbb{F}_2$ such that $\delta\eta_{2i-1, 2i'+2i, 2i-1} = \chi_{2i-1} \cup \chi_{2i} + \chi_{2i} \cup \chi_{2i-1}$. To do so, we use Dwyer's Theorem 3.3 and the following notation.

Notation 4.10. By slight abuse of notation, we will often write a continuous homomorphism $\rho: G \rightarrow U_n(\mathbb{F}_2)$ as an $(n \times n)$ -matrix with (i, j) -entry the map $e_{i,j} \circ \rho$.

For example, for $n = 4$, we write $\rho = \begin{pmatrix} 1 & a_{1,2} & a_{1,3} & a_{1,4} \\ & 1 & a_{2,3} & a_{2,4} \\ & & 1 & a_{3,4} \\ & & & 1 \end{pmatrix}$ where each $a_{i,j}: G \rightarrow \mathbb{F}_2$ is a continuous map, and $a_{1,2}$, $a_{2,3}$, and $a_{3,4}$ are group homomorphisms.

Notation 4.11. Let $n \geq 3$. For $1 \leq s < t \leq n$, we let $E_{s,t}$ denote the matrix in $U_n(\mathbb{F}_2)$ with only non-zero entry above the diagonal in row s and column t . We have $(E_{s,t})^q = I_n$ in $U_n(\mathbb{F}_2)$ for each pair (s, t) . We note that it will be clear from the context which n we consider and we therefore omit it from notation for the matrices $E_{s,t}$.

To construct the cochain $\eta_{2i-1, 2i'+2i, 2i-1}$ we use Dwyer's Theorem 3.3. We would like to define a continuous homomorphism $\rho_{2i-1, 2i'+2i, 2i-1}: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the continuous homomorphism $\bar{\rho}_{2i-1, 2i'+2i, 2i-1}: G \rightarrow \bar{U}_4(\mathbb{F}_2)$ given by $\bar{\rho}_{2i-1, 2i'+2i, 2i-1} = \begin{pmatrix} 1 & \chi_{2i-1} & \chi_{2i} & * \\ & 1 & 0 & \chi_{2i} \\ & & 1 & \chi_{2i-1} \\ & & & 1 \end{pmatrix}$. To construct the desired lift, consider the matrices

$$A := \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \text{ and } B := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $A^q = I_4$ and $[A, B] = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{2i-1, 2i'+2i, 2i-1}$ by the assignment $x_{2i-1} \mapsto A$, $x_{2i} \mapsto B$, and $x_s \mapsto I_4$ for $s \neq 2i-1, 2i$. We define the desired continuous map $G \rightarrow \mathbb{F}_2$ by setting

$$f_2(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) := e_{1,4} \circ \rho_{2i-1, 2i'+2i, 2i-1}.$$

Similarly, for $i \in \tilde{J}/2$, we construct a continuous homomorphism $\rho_{1, 2'+2i, 2i-1}: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the continuous homomorphism $\bar{\rho}_{1, 2'+2i, 2i-1}: G \rightarrow \bar{U}_4(\mathbb{F}_2)$ given

by $\bar{\rho}_{1, 2'+2i, 2i-1} = \begin{pmatrix} 1 & \chi_1 & \chi_{2i} & * \\ & 1 & 0 & \chi_2 \\ & & 1 & \chi_{2i-1} \\ & & & 1 \end{pmatrix}$. We have $[E_{1,2}, E_{2,4}] = [E_{3,4}, E_{1,3}] = E_{1,4}$ and

hence $[E_{1,2}, E_{2,4}][E_{3,4}, E_{1,3}] = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{1, 2'+2i, 2i-1}: G \rightarrow U_4(\mathbb{F}_2)$ by the assignment $x_1 \mapsto E_{1,2}$, $x_2 \mapsto E_{2,4}$, $x_{2i-1} \mapsto E_{3,4}$,

$x_{2i} \mapsto E_{1,3}$, and $x_s \mapsto I_4$ for $s \neq 1, 2, 2i-1, 2i$. We then define the desired continuous map $G \rightarrow \mathbb{F}_2$ by

$$f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) := e_{1,4} \circ \rho_{1,2'+2i,2i-1}.$$

We now define the map $f_2: R \rightarrow \mathcal{C}^1$ on all of R by extending it $\mathbb{F}_2$ -linearly from $\mathbf{B}_2$ to R . We then deduce from Proposition 2.15:

Proposition 4.12. *We define the map $\Psi_3: K_3^3(H^\bullet) \rightarrow \mathcal{Z}^2$ by*

$$(11) \quad \Psi_3(\chi_a, \chi_b, \chi_c) = \chi_a \cup f_2(\chi_b, \chi_c) + f_2(\chi_a, \chi_b) \cup \chi_c.$$

Taking the cohomology class of Ψ_3 defines an $\mathbb{F}_2$ -linear map $\kappa_3: K_3^3(H^\bullet) \rightarrow H^2$. The map κ_3 is a cocycle in the complex $(\underline{\mathrm{Hom}}_{\mathbb{F}_2}(K_\bullet^3(H^\bullet), H^\bullet[-1]), \partial)$, and the class of κ_3 in $\mathrm{HH}^{3,-1}(H^\bullet)$ is the canonical class of G . $\square$

4.4. Two compatibility results. For the proof of Theorem 4.1 in Section 4.5 we need the compatibility statements of Propositions 4.15 and 4.18 we prove below. To formulate and prove these results, we need the following lemmas in linear algebra.

Lemma 4.13. *Let $T_4^l(\mathbb{F}_2)$ be the subset of $U_4(\mathbb{F}_2)$ consisting of matrices of the form*

$$\begin{pmatrix} 1 & x & v & z \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ and let } W_4^l(\mathbb{F}_2) \text{ denote the subset of matrices of the form } \begin{pmatrix} 1 & 0 & v & 0 \\ 0 & 1 & w & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Then $T_4^l(\mathbb{F}_2)$ is a subgroup of $U_4(\mathbb{F}_2)$, and $W_4^l(\mathbb{F}_2)$ is a normal subgroup of $T_4^l(\mathbb{F}_2)$.

*Moreover, the map $\tau^l: U_3(\mathbb{F}_2) \rightarrow T_4^l(\mathbb{F}_2)/W_4^l(\mathbb{F}_2)$ defined by $\tau^l: \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & x & * & z \\ 0 & 1 & * & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$*

is an isomorphism of groups.

Proof. The following computation shows that $T_4^l(\mathbb{F}_2)$ is closed under the group operation:

$$(12) \quad \begin{pmatrix} 1 & x_1 & v_1 & z_1 \\ 0 & 1 & w_1 & y_1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & x_2 & v_2 & z_2 \\ 0 & 1 & w_2 & y_2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & x_1+x_2 & v_1+v_2+x_1w_2 & z_1+z_2+x_1y_2 \\ 0 & 1 & w_1+w_2 & y_1+y_2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Since $U_4(\mathbb{F}_2)$ is a finite group, this implies that $T_4^l(\mathbb{F}_2)$ is a subgroup. We note

that the inverse of $\begin{pmatrix} 1 & x & v & z \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ in $T_4^l(\mathbb{F}_2)$ is given by $\begin{pmatrix} 1 & x & v+xw & z+xy \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$. We also see

from (12) that $W_4^l(\mathbb{F}_2)$ is a subgroup of $T_4^l(\mathbb{F}_2)$. To show that $W_4^l(\mathbb{F}_2)$ is a normal subgroup of $T_4^l(\mathbb{F}_2)$, we observe

$$\begin{aligned} \begin{pmatrix} 1 & x & v & z \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & v' & 0 \\ 0 & 1 & w' & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & x & v+xw & z+xy \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} &= \begin{pmatrix} 1 & x & v+v'+xw' & z \\ 0 & 1 & w+w' & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & x & v+xw & z+xy \\ 0 & 1 & w & y \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & v'+xw' & 0 \\ 0 & 1 & w' & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in W_4^l(\mathbb{F}_2). \end{aligned}$$

Since $\#U_3(\mathbb{F}_2) = \#(T_4^l(\mathbb{F}_2)/W_4^l(\mathbb{F}_2))$, to show that τ^l is an isomorphism, it suffices to show that it is a homomorphism which follows from (12) and

$$(13) \quad \begin{pmatrix} 1 & x_1 & z_1 \\ 0 & 1 & y_1 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & x_2 & z_2 \\ 0 & 1 & y_2 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & x_1+x_2 & z_1+z_2+x_1y_2 \\ 0 & 1 & y_1+y_2 \\ 0 & 0 & 1 \end{pmatrix}$$

in $U_3(\mathbb{F}_2)$. $\square$

Lemma 4.14. *Let $T_4^r(\mathbb{F}_2)$ be the subset of $U_4(\mathbb{F}_2)$ consisting of matrices of the form*

$$\begin{pmatrix} 1 & 0 & x & z \\ 0 & 1 & w & v \\ 0 & 0 & 1 & y \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ and let } W_4^r(\mathbb{F}_2) \text{ denote the subset of matrices of the form } \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & w & v \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Then $T_4^r(\mathbb{F}_2)$ is a subgroup of $U_4(\mathbb{F}_2)$, and $W_4^r(\mathbb{F}_2)$ is a normal subgroup of $T_4^r(\mathbb{F}_2)$.
 Moreover, the map $\tau^r: U_3(\mathbb{F}_2) \rightarrow T_4^r(\mathbb{F}_2)/W_4^r(\mathbb{F}_2)$ defined by $\tau^r: \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & x & z \\ 0 & 1 & * & * \\ 0 & 0 & 1 & y \\ 0 & 0 & 0 & 1 \end{pmatrix}$
 is an isomorphism of groups.

Proof. This follows from an analogous but symmetric computation as in the proof of Lemma 4.13. $\square$

Proposition 4.15. Assume $\chi_i \otimes \chi_j \in R$. Let $f_2(\chi_i \otimes \chi_j): G \rightarrow \mathbb{F}_2$ be defined as in Section 4.3. Let $\rho_4^l: G \rightarrow T_4^l(\mathbb{F}_2)$ be a continuous group homomorphism satisfying $\rho_4^l(x_i) = E_{1,2}$ and $\rho_4^l(x_j) = E_{2,4}$, and $\rho_4^l(x_k) = I_4$ for $k \neq i, j$. Let $\rho_4^r: G \rightarrow T_4^r(\mathbb{F}_2)$ be a continuous group homomorphism satisfying $\rho_4^r(x_i) = E_{1,3}$ and $\rho_4^r(x_j) = E_{3,4}$, and $\rho_4^r(x_k) = I_4$ for $k \neq i, j$. Then

$$e_{1,4} \circ \rho_4^l = f_2(\chi_i \otimes \chi_j) = e_{1,4} \circ \rho_4^r$$

as continuous maps $G \rightarrow \mathbb{F}_2$.

Proof. The assertion for ρ_4^l follows from Lemma 4.13, and the assertion for ρ_4^r follows from Lemma 4.14. $\square$

We also need to consider the case of matrices in $U_5(\mathbb{F}_2)$.

Lemma 4.16. Let $T_5^l(\mathbb{F}_2)$ be the subset of $U_5(\mathbb{F}_2)$ consisting of matrices of the form $\begin{pmatrix} 1 & t & x & u & z \\ 0 & 1 & 0 & v & 0 \\ 0 & 0 & 1 & w & y \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$ and let $W_5^l(\mathbb{F}_2)$ denote the subset of matrices of the form $\begin{pmatrix} 1 & t & 0 & u & 0 \\ 0 & 1 & 0 & v & 0 \\ 0 & 0 & 1 & w & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$.
 Then $T_5^l(\mathbb{F}_2)$ is a subgroup of $U_5(\mathbb{F}_2)$, and $W_5^l(\mathbb{F}_2)$ is a normal subgroup of $T_5^l(\mathbb{F}_2)$.

Moreover, the map $\tau_5^l: U_3(\mathbb{F}_2) \rightarrow T_5^l(\mathbb{F}_2)/W_5^l(\mathbb{F}_2)$ defined by $\tau_5^l: \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & * & x & * & z \\ 0 & 1 & 0 & * & 0 \\ 0 & 0 & 1 & * & y \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
 is an isomorphism of groups.

Proof. This follows from a direct computation as in the proof of Lemma 4.13. $\square$

Lemma 4.17. Let $T_5^r(\mathbb{F}_2)$ be the subset of $U_5(\mathbb{F}_2)$ consisting of matrices of the form $\begin{pmatrix} 1 & 0 & x & 0 & z \\ 0 & 1 & w & v & u \\ 0 & 0 & 1 & 0 & y \\ 0 & 0 & 0 & 1 & t \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$ and let $W_5^r(\mathbb{F}_2)$ denote the subset of matrices of the form $\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & w & v & u \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & t \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$.
 Then $T_5^r(\mathbb{F}_2)$ is a subgroup of $U_5(\mathbb{F}_2)$, and $W_5^r(\mathbb{F}_2)$ is a normal subgroup of $T_5^r(\mathbb{F}_2)$.

Moreover, the map $\tau_5^r: U_3(\mathbb{F}_2) \rightarrow T_5^r(\mathbb{F}_2)/W_5^r(\mathbb{F}_2)$ defined by $\tau_5^r: \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & x & 0 & z \\ 0 & 1 & * & * & * \\ 0 & 0 & 1 & 0 & y \\ 0 & 0 & 0 & 1 & * \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
 is an isomorphism of groups.

Proof. This follows from a direct computation as in the proof of Lemma 4.13. $\square$

Proposition 4.18. Assume $\chi_i \otimes \chi_j \in R$. Let $f_2(\chi_i \otimes \chi_j): G \rightarrow \mathbb{F}_2$ be defined as in Section 4.3. Let $\rho_5^l: G \rightarrow T_5^l(\mathbb{F}_2)$ be a continuous group homomorphism satisfying $\rho_5^l(x_i) = E_{1,3}$ and $\rho_5^l(x_j) = E_{3,5}$, and $\rho_5^l(x_k) = I_5$ for $k \neq i, j$. Let $\rho_5^r: G \rightarrow T_5^r(\mathbb{F}_2)$ be a continuous group homomorphism satisfying $\rho_5^r(x_i) = E_{1,3}$ and $\rho_5^r(x_j) = E_{3,5}$, and $\rho_5^r(x_k) = I_5$ for $k \neq i, j$. Then

$$e_{1,5} \circ \rho_5^l = f_2(\chi_i \otimes \chi_j) = e_{1,5} \circ \rho_5^r$$

as continuous maps $G \rightarrow \mathbb{F}_2$.

Proof. The assertion for ρ_5^l follows from Lemma 4.16, and the assertion for ρ_5^r follows from Lemma 4.17. $\square$

4.5. Computation of the canonical class. We will now compute the values of κ_3 . To do so, we will make frequent use of the following notation.

Notation 4.19. For $i = 1, 2, 3$ and $\varepsilon_i \in \mathbb{F}_2$, we define the matrix

$$B_{\varepsilon_1, \varepsilon_2, \varepsilon_3} = \begin{pmatrix} 1 & \varepsilon_1 & 0 & 0 \\ 0 & 1 & \varepsilon_2 & 0 \\ 0 & 0 & 1 & \varepsilon_3 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in U_4(\mathbb{F}_2).$$

Note that a direct computation shows $B_{\varepsilon_1, \varepsilon_2, \varepsilon_3}^q = I_4$ in $U_4(\mathbb{F}_2)$.

Convention 4.20. We embed $U_n(\mathbb{F}_2)$ as a subgroup in $U_{n+1}(\mathbb{F}_2)$ either by sending a matrix $A \in U_n(\mathbb{F}_2)$ to $\begin{pmatrix} 1 & 0 \\ 0 & A \end{pmatrix} \in U_{n+1}(\mathbb{F}_2)$, or to $\begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} \in U_{n+1}(\mathbb{F}_2)$, where the 0s denote rows and columns with entry 0 respectively.

Proposition 4.21. *Let $\kappa_3: K_3^3(H^\bullet) \rightarrow H^2$ be the map defined in Proposition 4.12. The map κ_3 vanishes on all elements in the sets **S**, **D** and **T**.*

Proof. Let Ψ_3 denote the cocycle defined in (11). We prove the assertion by constructing cochains $\vartheta \in \mathcal{C}^1$ such that $\delta\vartheta = \Psi_3$ for the elements in each of the sets using Dwyer's Theorem 3.3. We consider each set separately. We begin with elements in **S**. By definition of $K_3^3(H^\bullet)$, if $\chi_a \otimes \chi_b \otimes \chi_c \in H^1 \otimes H^1 \otimes H^1$ is an element in **S**, then the triple Massey product $\langle \chi_a, \chi_b, \chi_c \rangle$ is defined. Moreover, $\{\chi_a, \chi_b, \chi_c, f_2(\chi_a \otimes \chi_b), f_2(\chi_b \otimes \chi_c)\}$ is a defining system, and $\kappa_3(\chi_a, \chi_b, \chi_c)$ is the class of this defining system. Thus, by Dwyer's Theorem 3.3, $\kappa_3(\chi_a, \chi_b, \chi_c) = 0$ if and only if the continuous group homomorphism $\bar{\rho}: G \rightarrow \bar{U}_4(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_a & f_2(\chi_a \otimes \chi_b) & * \\ & 1 & \chi_b & f_2(\chi_b \otimes \chi_c) \\ & & 1 & \chi_c \\ & & & 1 \end{pmatrix}$$

lifts to a continuous group homomorphism $\rho: G \rightarrow U_4(\mathbb{F}_2)$. If a lift ρ exists, then the continuous map $\vartheta := e_{1,4} \circ \rho: G \rightarrow \mathbb{F}_2$ is a cochain such that $\delta\vartheta = \Psi_3(\chi_a \otimes \chi_b \otimes \chi_c)$. Thus, to prove $\kappa_3(\chi_a, \chi_b, \chi_c) = 0$ for an element in **S**, it suffices to construct the corresponding continuous homomorphism ρ .

For elements of the form $\chi_i \otimes \chi_i \otimes \chi_i$, we define the continuous group homomorphism $\rho_{i,i,i}^{\mathbf{S}}: G \rightarrow U_4(\mathbb{F}_2)$ by setting $\rho_{i,i,i}^{\mathbf{S}}(x_i) := B_{1,1,1}$ and $\rho_{i,i,i}^{\mathbf{S}}(x_j) := I_4$ for $j \neq i$. For elements of the form $\chi_i \otimes \chi_i \otimes \chi_j$ with $i \neq j$ and $i \neq j-1$ if j is even, we define the continuous group homomorphism $\rho_{i,i,j}^{\mathbf{S}}: G \rightarrow U_4(\mathbb{F}_2)$ by setting

$$\rho_{i,i,j}^{\mathbf{S}}(x_i) := B_{1,1,0}, \rho_{i,i,j}^{\mathbf{S}}(x_j) := B_{0,0,1}, \text{ and } \rho_{i,i,j}^{\mathbf{S}}(x_k) := I_4 \text{ for } k \neq i, j.$$

For elements of the form $\chi_i \otimes \chi_j \otimes \chi_j$ with $i \neq j$ and $i \neq j-1$ if j is even, we define the continuous group homomorphism $\rho_{i,j,j}^{\mathbf{S}}: G \rightarrow U_4(\mathbb{F}_2)$ by setting

$$\rho_{i,j,j}^{\mathbf{S}}(x_i) := B_{1,0,0}, \rho_{i,j,j}^{\mathbf{S}}(x_j) := B_{0,1,1}, \text{ and } \rho_{i,j,j}^{\mathbf{S}}(x_k) := I_4 \text{ for } k \neq i, j.$$

For elements of the form $\chi_i \otimes \chi_j \otimes \chi_k$ with $i, k \neq j$, $i \neq j-1$ if j is even and $k \neq j+1$ if j is odd, we define a continuous group homomorphism $\rho_{i,j,k}^{\mathbf{S}}: G \rightarrow U_4(\mathbb{F}_2)$ by setting

$$\rho_{i,j,k}^{\mathbf{S}}(x_i) := B_{1,0,0}, \rho_{i,j,k}^{\mathbf{S}}(x_j) := B_{0,1,0}, \rho_{i,j,k}^{\mathbf{S}}(x_k) := B_{0,0,1},$$

and $\rho_{i,j,k}^{\mathbf{S}}(x_l) := I_4$ for $l \neq i, j, k$. This finishes the proof for the set **S**.

Next we consider elements in **D**. First, let $i \in J/2$ and $k \neq 2i-1, 2i$, and we consider an element of the form $\chi_k \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1})$. While $\chi_k \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1})$ does not correspond to a triple Massey product, the fourfold Massey product $\langle \chi_k, \chi_{2i-1}, 0, \chi_{2i-1} \rangle$ is defined, and by Dwyer's Theorem

3.3, the continuous homomorphism $\bar{\rho}_{k,2i-1,2i}^{\mathbf{D}}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho}_{k,2i-1,2i}^{\mathbf{D}} = \begin{pmatrix} 1 & \chi_k & f_2(\chi_k \otimes \chi_{2i-1}) & f_2(\chi_k \otimes \chi_{2i}) & * \\ & 1 & \chi_{2i-1} & \chi_{2i} & f_2(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) \\ & & 1 & 0 & \chi_{2i} \\ & & & 1 & \chi_{2i-1} \\ & & & & 1 \end{pmatrix}$$

corresponds to a defining system. Moreover, we have $\kappa_3(\chi_k \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1})) = 0$ if and only if $\bar{\rho}_{k,2i-1,2i}^{\mathbf{D}}$ lifts to a continuous homomorphism $\rho_{k,2i-1,2i}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$. To construct $\rho_{k,2i-1,2i}^{\mathbf{D}}$, we consider the matrices

$$A := \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ & 1 & 1 & 0 & 0 \\ & & 1 & 0 & 0 \\ & & & 1 & 1 \\ & & & & 1 \end{pmatrix}, \quad B := \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 0 \\ & & 1 & 0 & 1 \\ & & & 1 & 0 \\ & & & & 1 \end{pmatrix}$$

in $U_5(\mathbb{F}_2)$. We have $[A, B] = I_5$. Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho_{k,2i-1,2i}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_k \mapsto E_{1,2}, \quad x_{2i-1} \mapsto A, \quad x_{2i} \mapsto B,$$

and $\rho_{k,2i-1,2i}^{\mathbf{D}}(x_j) = I_5$ for $j \neq k, 2i-1, 2i$. We use Proposition 4.15 to identify $e_{1,4} \circ \rho_{k,2i-1,2i}^{\mathbf{D}}$ with $f_2(\chi_k \otimes \chi_{2i})$. We define the continuous map $\vartheta_{k,2i-1,2i}^{\mathbf{D}}: G \rightarrow \mathbb{F}_2$ by $e_{1,5} \circ \rho_{k,2i-1,2i}^{\mathbf{D}}$ and get

$$\delta \vartheta_{k,2i-1,2i}^{\mathbf{D}} = \Psi_3(\chi_k \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1})).$$

We prove the other cases by specifying the corresponding homomorphisms $\bar{\rho}$ and constructing the required lifts ρ . For elements of the form $(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_k$, we need to show that the continuous homomorphism $\bar{\rho}_{2i-1,2i,k}^{\mathbf{D}}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho}_{2i-1,2i,k}^{\mathbf{D}} = \begin{pmatrix} 1 & \chi_{2i-1} & \chi_{2i} & f_2(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) & * \\ & 1 & 0 & \chi_{2i} & f_2(\chi_{2i} \otimes \chi_k) \\ & & 1 & \chi_{2i-1} & f_2(\chi_{2i-1} \otimes \chi_k) \\ & & & 1 & \chi_k \\ & & & & 1 \end{pmatrix}$$

lifts to a continuous homomorphism $\rho_{2i-1,2i,k}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$. Consider the matrices

$$C := \begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ & 1 & 0 & 0 & 0 \\ & & 1 & 1 & 0 \\ & & & 1 & 0 \\ & & & & 1 \end{pmatrix}, \quad D := \begin{pmatrix} 1 & 0 & 1 & 0 & 0 \\ & 1 & 0 & 1 & 0 \\ & & 1 & 0 & 0 \\ & & & 1 & 0 \\ & & & & 1 \end{pmatrix}$$

in $U_5(\mathbb{F}_2)$. We have $[C, D] = I_5$. By Lemma 3.10, we can define a continuous group homomorphism $\rho_{2i-1,2i,k}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_k \mapsto E_{4,5}, \quad x_{2i-1} \mapsto C, \quad x_{2i} \mapsto D,$$

and $\rho_{2i-1,2i,k}^{\mathbf{D}}(x_j) = I_5$ for $j \neq k, 2i-1, 2i$. We use Proposition 4.15 to identify $e_{2,5} \circ \rho_{2i-1,2i,k}^{\mathbf{D}}$ with $f_2(\chi_{2i} \otimes \chi_k)$.

Second, let $i \in \tilde{J}/2$ and $k \neq 1, 2, 2i-1, 2i$. For elements of the form $\chi_k \otimes (\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1})$, we need to show that the continuous homomorphism $\bar{\rho}_{k,1,2+2i,2i-1}^{\mathbf{D}}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho}_{k,1,2+2i,2i-1}^{\mathbf{D}} = \begin{pmatrix} 1 & \chi_k & f_2(\chi_k \otimes \chi_1) & f_2(\chi_k \otimes \chi_{2i}) & * \\ & 1 & \chi_1 & \chi_{2i} & f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \\ & & 1 & 0 & \chi_2 \\ & & & 1 & \chi_{2i-1} \\ & & & & 1 \end{pmatrix}$$

lifts to a continuous homomorphism $\rho_{k,1,2+2i,2i-1}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$. We have the relation $[E_{2,3}, E_{3,5}][E_{4,5}, E_{2,4}] = I_5$. (Note, however, that $[E_{2,3}, E_{3,5}] = [E_{4,5}, E_{2,4}] =$

$E_{2,5} \neq I_5$.) By Lemma 3.10, we can define a continuous group homomorphism $\rho_{k,1,2+2i,2i-1}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_k \mapsto E_{1,2}, x_1 \mapsto E_{2,3}, x_2 \mapsto E_{3,5}, x_{2i-1} \mapsto E_{4,5}, x_{2i} \mapsto E_{2,4},$$

and $\rho_{k,1,2+2i,2i-1}^{\mathbf{D}}(x_j) = I_5$ for $j \neq k, 1, 2, 2i-1, 2i$. We use Proposition 4.15 to identify $e_{1,4} \circ \rho_{k,1,2+2i,2i-1}^{\mathbf{D}}$ with $f_2(\chi_k \otimes \chi_{2i})$. For elements of the form $(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_k$, we need to show that the continuous homomorphism $\bar{\rho}_{1,2+2i,2i-1,k}^{\mathbf{D}}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho}_{1,2+2i,2i-1,k}^{\mathbf{D}} = \begin{pmatrix} 1 & \chi_1 & \chi_{2i} & f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) & * \\ & 1 & 0 & \chi_2 & f_2(\chi_2 \otimes \chi_k) \\ & & 1 & \chi_{2i-1} & f_2(\chi_{2i-1} \otimes \chi_k) \\ & & & 1 & \chi_k \\ & & & & 1 \end{pmatrix}$$

lifts to a continuous homomorphism $\rho_{1,2+2i,2i-1,k}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$. We have the relation $[E_{1,2}, E_{2,4}][E_{3,4}, E_{1,3}] = I_5$. (Note, however, that $[E_{1,2}, E_{2,4}] = [E_{3,4}, E_{1,3}] = E_{1,4} \neq I_5$.) By Lemma 3.10, we can define a continuous group homomorphism $\rho_{1,2+2i,2i-1,k}^{\mathbf{D}}: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_k \mapsto E_{4,5}, x_1 \mapsto E_{1,2}, x_2 \mapsto E_{2,4}, x_{2i-1} \mapsto E_{3,4}, x_{2i} \mapsto E_{1,3},$$

and $\rho_{1,2+2i,2i-1,k}^{\mathbf{D}}(x_j) = I_5$ for $j \neq k, 1, 2, 2i-1, 2i$. We use Proposition 4.15 to identify $e_{2,5} \circ \rho_{1,2+2i,2i-1,k}^{\mathbf{D}}$ with $f_2(\chi_2 \otimes \chi_k)$. This proves the assertion for $\mathbf{D}$.

Finally, we compute κ_3 on elements in $\mathbf{T}$. For $i \in J/2$, we consider $\chi_{2i-1,2i-1,2i} + \chi_{2i-1,2i,2i-1} + \chi_{2i,2i-1,2i-1}$. By Dwyer's Theorem 3.3, the continuous group homomorphism $\bar{\rho}_{2i-1}^{\mathbf{T}}: G \rightarrow \bar{U}_6(\mathbb{F}_2)$ given by

$$\bar{\rho}_{2i-1}^{\mathbf{T}} = \begin{pmatrix} 1 & \chi_{2i-1} & \chi_{2i} & f_2(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) & f_2(\chi_{2i-1} \otimes \chi_{2i-1}) & * \\ & 1 & 0 & \chi_{2i} & \chi_{2i-1} & f_2(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) \\ & & 1 & \chi_{2i-1} & 0 & f_2(\chi_{2i-1} \otimes \chi_{2i-1}) \\ & & & 1 & 0 & \chi_{2i-1} \\ & & & & 1 & \chi_{2i} \\ & & & & & 1 \end{pmatrix}$$

corresponds to a defining system for the fivefold Massey product $\langle \chi_{2i-1}, 0, 0, 0, \chi_{2i-1} \rangle$. Consider the matrices

$$S := \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 0 & 1 & 0 \\ & & 1 & 1 & 0 & 0 \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}, T := \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 0 & 0 \\ & & 1 & 0 & 0 & 0 \\ & & & 1 & 0 & 0 \\ & & & & 1 & 1 \\ & & & & & 1 \end{pmatrix},$$

in $U_6(\mathbb{F}_2)$. We have $S^q = I_6$, $T^q = I_6$, and $[S, T] = I_6$ in $U_6(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define continuous group homomorphisms $\rho_{2i-1}^{\mathbf{T}}: G \rightarrow U_6(\mathbb{F}_2)$ by the assignment $x_{2i-1} \mapsto S$, $x_{2i} \mapsto T$, and $\rho_{2i-1}^{\mathbf{T}}(x_j) = I_6$ for $j \neq 2i-1, 2i$. Then $\rho_{2i-1}^{\mathbf{T}}$ lifts $\bar{\rho}_{2i-1}^{\mathbf{T}}$. This implies $\kappa_3(\chi_{2i-1,2i-1,2i} + \chi_{2i-1,2i,2i-1} + \chi_{2i,2i-1,2i-1}) = 0$. Here, we use Convention 4.20 and Proposition 4.15 to identify $e_{3,6} \circ \rho_{2i-1}^{\mathbf{T}}$ and Proposition 4.18 to identify $e_{1,5} \circ \rho_{2i-1}^{\mathbf{T}}$ with $f_2(\chi_{2i-1} \otimes \chi_{2i-1})$. By switching the roles of χ_{2i-1} and χ_{2i} , we define $\rho_{2i}^{\mathbf{T}}: G \rightarrow U_6(\mathbb{F}_2)$ by the assignment $x_{2i-1} \mapsto T$, $x_{2i} \mapsto S$, and $\rho_{2i}^{\mathbf{T}}(x_j) = I_6$ for $j \neq 2i-1, 2i$. The existence of $\rho_{2i}^{\mathbf{T}}$ implies $\kappa_3(\chi_{2i,2i,2i-1} + \chi_{2i,2i-1,2i} + \chi_{2i-1,2i,2i}) = 0$. This proves the assertion for the set $\mathbf{T}$ and finishes the proof. $\square$

For $d \geq 4$, however, κ_3 is a non-trivial map as the proof of the following proposition shows.

Proposition 4.22. *For all $i, j \in \tilde{J}/2$, we have*

$$\begin{aligned}\kappa_3(\chi_{1,1,2} + \chi_{1,2i,2i-1}) &= \kappa_3(\chi_{1,1,2} + \chi_{1,2j,2j-1}), \\ \kappa_3(\chi_{1,2,2} + \chi_{2i,2i-1,2}) &= \kappa_3(\chi_{1,2,2} + \chi_{2j,2j-1,2}), \\ \kappa_3(\chi_{2,1,1} + \chi_{2i-1,2i,1}) &= \kappa_3(\chi_{2,1,1} + \chi_{2j-1,2j,1}), \text{ and} \\ \kappa_3(\chi_{2,2,1} + \chi_{2,2i-1,2i}) &= \kappa_3(\chi_{2,2,1} + \chi_{2,2j-1,2j}).\end{aligned}$$

Proof. The claimed relations follow from the symmetry of the roles of the χ_i . However, we also give a more concrete proof by computing the respective values of κ_3 using Dwyer's Theorem 3.3 similar to the arguments in the proof of Proposition 4.21. For the first relation, we show that the continuous homomorphism $\bar{\rho}_1: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho}_1 = \begin{pmatrix} 1 & \chi_1 & f_2(\chi_1 \otimes \chi_1) & f_2(\chi_1 \otimes \chi_{2i}) & * \\ & 1 & \chi_1 & \chi_{2i} & f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \\ & & 1 & 0 & \chi_2 \\ & & & 1 & \chi_{2i-1} \\ & & & & 1 \end{pmatrix}$$

does not lift to a continuous homomorphism $G \rightarrow U_5(\mathbb{F}_2)$. Let $A := \begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ & 1 & 1 & 0 & 0 \\ & & 1 & 0 & 0 \\ & & & 1 & 0 \\ & & & & 1 \end{pmatrix}$ in $U_5(\mathbb{F}_2)$. Then we have $A^q = I_5$ and $[A, E_{3,5}][E_{4,5}, E_{2,4}] = I_5$ in $\bar{U}_5(\mathbb{F}_2)$, but $[A, E_{3,5}][E_{4,5}, E_{2,4}] = E_{1,5} \neq I_5$ in $U_5(\mathbb{F}_2)$. Thus, we can define a continuous group homomorphism $\bar{\rho}_1: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto A, \quad x_2 \mapsto E_{3,5}, \quad x_{2i-1} \mapsto E_{4,5}, \quad x_{2i} \mapsto E_{2,4},$$

and $\bar{\rho}_1(x_j) := I_5$ for $j \neq 1, 2, 2i-1, 2i$, but $\bar{\rho}_1$ does not lift to a continuous group homomorphism $G \rightarrow U_5(\mathbb{F}_2)$. Note that we use Proposition 4.15 to identify $e_{1,4} \circ \bar{\rho}_1$ with $f_2(\chi_1 \otimes \chi_{2i})$. This proves $\kappa_3(\chi_{1,1,2} + \chi_{1,2i,2i-1}) \neq 0$ for all $i \in \tilde{J}/2$. This implies the first relation since there is a unique non-zero element in $H^2 \cong \mathbb{F}_2$.

For the second relation, we show that the continuous group homomorphism $\bar{\rho}_2: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho}_2 = \begin{pmatrix} 1 & \chi_1 & \chi_{2i} & f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) & * \\ & 1 & 0 & \chi_2 & f_2(\chi_2 \otimes \chi_2) \\ & & 1 & \chi_{2i-1} & f_2(\chi_{2i-1} \otimes \chi_2) \\ & & & 1 & \chi_2 \\ & & & & 1 \end{pmatrix}$$

lifts to a continuous homomorphism $\rho_2: G \rightarrow U_5(\mathbb{F}_2)$. Let $B := \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 0 \\ & & 1 & 0 & 0 \\ & & & 1 & 1 \\ & & & & 1 \end{pmatrix}$ in $U_5(\mathbb{F}_2)$. Then we have $[E_{1,2}, B][E_{3,4}, E_{1,3}] = I_5$ in $U_5(\mathbb{F}_2)$. Note, however, that $[E_{1,2}, B] = [E_{3,4}, E_{1,3}] = E_{1,4} \neq I_5$. Thus, we can define a continuous group homomorphism $\rho_2: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto E_{1,2}, \quad x_2 \mapsto B, \quad x_{2i-1} \mapsto E_{3,4}, \quad x_{2i} \mapsto E_{1,3},$$

and $\rho_2(x_j) := I_5$ for $j \neq 1, 2, 2i-1, 2i$, which lifts $\bar{\rho}_2$. Here we use Proposition 4.15 to identify $e_{2,5} \circ \bar{\rho}_2$ and $e_{2,5} \circ \rho_2$ with $f_2(\chi_2 \otimes \chi_2)$. This proves $\kappa_3(\chi_{1,2,2} + \chi_{2i,2i-1,2}) = 0$ for all $i \in \tilde{J}/2$.

For the third relation, we note that in $H^1 \otimes H^1 \otimes H^1$ we have

$$\begin{aligned} & (\chi_{2,1} + \chi_{2i-1,2i}) \otimes \chi_1 \\ &= [(\chi_{1,2} + \chi_{2,1}) + (\chi_{2i-1,2i} + \chi_{2i,2i-1}) + (\chi_{1,2} + \chi_{2i,2i-1})] \otimes \chi_1. \end{aligned}$$

Since $\chi_{1,2,1} + \chi_{2,1,1}$ and $\chi_{1,2,1} + \chi_{2i,2i-1,1}$ are not in $K_3^3(H^\bullet)$, we add $\chi_{1,1,2}$ twice to the sum and get

$$\begin{aligned} (\chi_{2,1} + \chi_{2i-1,2i}) \otimes \chi_1 &= (\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1}) \\ &\quad + ((\chi_{2i-1,2i} + \chi_{2i,2i-1}) \otimes \chi_1) + (\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2i,2i-1,1}). \end{aligned}$$

Here we note that the latter is not a linear combination of elements in $\mathbf{B}_3$, and that κ_3 is defined independently of our choice of basis for $K_3^3(H^\bullet)$. We know $\kappa_3(\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1}) = 0$ by the computation for elements in $\mathbf{T}$, and $\kappa_3((\chi_{2i-1,2i} + \chi_{2i,2i-1}) \otimes \chi_1) = 0$ by the computation for elements in $\mathbf{D}$. Thus, $\kappa_3(\chi_{2,1,1} + \chi_{2i-1,2i,1}) = \kappa_3(\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2i,2i-1,1})$, and it suffices to compute the latter. To compute $\kappa_3(\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2i,2i-1,1})$, we show that the continuous homomorphism $\bar{\rho}_3: G \rightarrow \bar{U}_6(\mathbb{F}_2)$ given by

$$\bar{\rho}_3 = \begin{pmatrix} 1 & \chi_{2i} & \chi_1 & f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) & f_2(\chi_1 \otimes \chi_1) & * \\ & 1 & 0 & \chi_{2i-1} & 0 & f_2(\chi_{2i-1} \otimes \chi_1) \\ & & 1 & \chi_2 & \chi_1 & f_2(\chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_1) \\ & & & 1 & 0 & \chi_1 \\ & & & & 1 & \chi_2 \\ & & & & & 1 \end{pmatrix}$$

lifts to a continuous homomorphism $\rho_3: G \rightarrow U_6(\mathbb{F}_2)$. Let $M := \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ & 1 & 0 & 0 & 0 & 0 \\ & & 1 & 1 & 0 & 0 \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}$

in $U_6(\mathbb{F}_2)$. Then $M^q = I_6$ and $[M, N][E_{2,4}, E_{1,2}] = I_6$ in $U_6(\mathbb{F}_2)$. (Note that $[M, N] = [E_{2,4}, E_{1,2}] = E_{1,4}$ in $U_6(\mathbb{F}_2)$.) Thus, we can define a continuous group homomorphism $\rho_3: G \rightarrow U_6(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto M, \quad x_2 \mapsto N, \quad x_{2i-1} \mapsto E_{2,4}, \quad x_{2i} \mapsto E_{1,2},$$

and $\rho_3(x_j) := I_6$ for $j \neq 1, 2, 2i-1, 2i$, which lifts $\bar{\rho}_3$. Here we use Proposition 4.18 to identify $e_{1,4} \circ \bar{\rho}_3$ and $e_{1,4} \circ \rho_3$ with $f_2(\chi_1 \otimes \chi_1)$, and $e_{2,5} \circ \bar{\rho}_3$ and $e_{2,5} \circ \rho_3$ with $f_2(\chi_{2i-1} \otimes \chi_1)$. This proves

$$\kappa_3(\chi_{2,1,1} + \chi_{2i-1,2i,1}) = \kappa_3(\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2i,2i-1,1}) = 0 \text{ for all } i \in \tilde{J}/2,$$

and thus the third relation.

For the fourth relation, we note that in $H^1 \otimes H^1 \otimes H^1$ we have

$$\begin{aligned} &\chi_2 \otimes (\chi_{2,1} + \chi_{2i-1,2i}) \\ &= \chi_2 \otimes [(\chi_{1,2} + \chi_{2,1}) + (\chi_{2i-1,2i} + \chi_{2i,2i-1}) + (\chi_{1,2} + \chi_{2i,2i-1})]. \end{aligned}$$

Since $\chi_{2,1,2} + \chi_{2,2,1}$ and $\chi_{2,1,2} + \chi_{2,2i,2i-1}$ are not in $K_3^3(H^\bullet)$, we add $\chi_{1,2,2}$ twice to the sum and get

$$\begin{aligned} &\chi_2 \otimes (\chi_{2,1} + \chi_{2i-1,2i}) \\ &= (\chi_{1,2,2} + \chi_{2,1,2} + \chi_{2,2,1}) + (\chi_2 \otimes (\chi_{2i-1,2i} + \chi_{2i,2i-1})) + (\chi_{1,2,2} + \chi_{2,1,2} + \chi_{2,2i,2i-1}). \end{aligned}$$

We know $\kappa_3(\chi_{1,2,2} + \chi_{2,1,2} + \chi_{2,2,1}) = 0$ by the computation for elements in $\mathbf{T}$, and $\kappa_3(\chi_2 \otimes (\chi_{2i-1,2i} + \chi_{2i,2i-1})) = 0$ by the computation for elements in $\mathbf{D}$. Thus, $\kappa_3(\chi_{2,2,1} + \chi_{2,2i-1,2i}) = \kappa_3(\chi_{1,2,2} + \chi_{2,1,2} + \chi_{2,2i,2i-1})$, and it suffices to compute the latter. To compute $\kappa_3(\chi_{1,2,2} + \chi_{2,1,2} + \chi_{2,2i,2i-1})$, we show that the continuous

homomorphism $\bar{\rho}_4: G \rightarrow \bar{U}_6(\mathbb{F}_2)$ given by

$$\bar{\rho}_4 = \begin{pmatrix} 1 & \chi_1 & \chi_2 & f_2(\chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_1) & f_2(\chi_2 \otimes \chi_{2i}) & * \\ & 1 & 0 & \chi_2 & 0 & f_2(\chi_2 \otimes \chi_2) \\ & & 1 & \chi_1 & \chi_{2i} & f_2(\chi_1 \otimes \chi_2 + \chi_{2i} \otimes \chi_{2i-1}) \\ & & & 1 & 0 & \chi_2 \\ & & & & 1 & \chi_{2i-1} \\ & & & & & 1 \end{pmatrix}$$

does not lift to a continuous homomorphism $G \rightarrow U_6(\mathbb{F}_2)$. Let $P := \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 0 & 0 & 0 \\ & & 1 & 0 & 0 & 0 \\ & & & 1 & 0 & 0 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}$

$Q := \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 0 & 0 \\ & & 1 & 0 & 0 & 0 \\ & & & 1 & 0 & 0 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}$ in $U_6(\mathbb{F}_2)$. Then $P^q = I_6$ and $[P, Q][E_{5,6}, E_{3,5}] = I_6$ in $\bar{U}_6(\mathbb{F}_2)$,

but $[P, Q][E_{5,6}, E_{3,5}] = E_{1,6} \neq I_6$ in $U_6(\mathbb{F}_2)$. Thus, we can define a continuous group homomorphism $\bar{\rho}_4: G \rightarrow \bar{U}_6(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto P, \quad x_2 \mapsto Q, \quad x_{2i-1} \mapsto E_{5,6}, \quad x_{2i} \mapsto E_{3,5},$$

and $\bar{\rho}_4(x_j) := I_6$ for $j \neq 1, 2, 2i-1, 2i$, but $\bar{\rho}_4$ does not lift to a continuous group homomorphism $G \rightarrow U_6(\mathbb{F}_2)$. Here we use Proposition 4.18 to identify $e_{1,4} \circ \bar{\rho}_4$ and $e_{1,4} \circ \rho_4$ with $f_2(\chi_2 \otimes \chi_{2i})$, and $e_{2,5} \circ \bar{\rho}_4$ and $e_{2,5} \circ \rho_4$ with $f_2(\chi_2 \otimes \chi_2)$. This proves

$$\kappa_3(\chi_{2,2,1} + \chi_{2,2i-1,2i}) = \kappa_3(\chi_{1,2,2} + \chi_{2,1,2} + \chi_{2,2i,2i-1}) \neq 0 \text{ for all } i \in \tilde{J}/2,$$

and thus the fourth relation, since there is a unique non-zero element in H^2 . $\square$

Proof of Theorem 4.1. The theorem now follows from Corollary 4.9, and Propositions 4.21, and 4.22. $\square$

5. DEMUSHKIN GROUPS OF TYPE II

The goal of this section is to prove the following theorem.

Theorem 5.1. *Let $d \geq 3$ be an odd number, and let G be the pro-2 group generated by elements $x_1, \dots, x_d$ with the single relation $x_1^2 x_2^{2^f} [x_2, x_3] \cdots [x_{d-1}, x_d] = 1$ with $f \geq 2$ or $f = \infty$ with the convention $2^\infty := 0$. Then the canonical class of G is trivial, and G is A_3 -formal.*

In this section, we let G be a pro-2 group as in Theorem 5.1. We use the same notation for $\mathcal{C}^\bullet$, $\mathcal{Z}^\bullet$, $H^\bullet$, and χ_i as in Section 4. We write again $J := \{1, \dots, d\}$ and now set $J/2 := \{1, \dots, (d-1)/2\}$.

5.1. The Koszul–Hochschild complex - Demushkin groups of type II. Again, by [30, Theorem 5.2], the graded $\mathbb{F}_2$ -cohomology algebra of G is a Koszul algebra. We now provide a basis for R and $K_3^3(H^\bullet)$.

Proposition 5.2. *The $\mathbb{F}_2$ -vector subspace $R \subset H^1 \otimes H^1$ such that $H^\bullet = T(H^1)/(R)$ admits the set $\mathbf{B}_2^\Pi$ as a basis where*

$$\begin{aligned} \mathbf{B}_2^\Pi := & \{\chi_i \otimes \chi_j \text{ for all } (i, j) \notin \{(1, 1), (2r, 2r+1), (2r+1, 2r) : r \in J/2\}; \\ & \chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i}, \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \text{ for all } i \in J/2\}. \end{aligned}$$

Proof. This follows from the description of the cup product on $H^\bullet(G, \mathbb{F}_2)$ in Lemma 3.11. $\square$

Proposition 5.3. *The $\mathbb{F}_2$ -vector space $K_3^3(H^\bullet)$ admits the set $\mathbf{B}_3^\Pi = \mathbf{S}^\Pi \cup \mathbf{D}^\Pi \cup \mathbf{T}^\Pi \cup \widetilde{\mathbf{T}}^\Pi \cup \mathbf{E}^\Pi$ as a basis where*

$$\mathbf{S}^\Pi = \left\{ \chi_i \otimes \chi_j \otimes \chi_k \text{ with } \begin{cases} i, k \neq 1 & \text{if } j = 1 \\ i, k \neq 2r+1 & \text{if } j = 2r \\ i, k \neq 2r & \text{if } j = 2r+1 \end{cases} \right\},$$

$$\begin{aligned} \mathbf{D}^\Pi = & \{ \chi_j \otimes (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}), \\ & (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j \text{ for } i \in J/2, j \neq 2i, 2i+1, \\ & \chi_j \otimes (\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i}), \text{ for } i \in J/2, j \neq 1, 2i \\ & (\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j \text{ for } i \in J/2, j \neq 1, 2i+1 \}, \end{aligned}$$

$$\begin{aligned} \mathbf{T}^\Pi = & \{ \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i}, \\ & \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i} \otimes \chi_{2i+1} \text{ for } i \in J/2 \}, \end{aligned}$$

$$\begin{aligned} \widetilde{\mathbf{T}}^\Pi = & \{ \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_1, \\ & \chi_{2i} \otimes \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i}, \\ & \chi_1 \otimes \chi_1 \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i}, \text{ for } i \in J/2 \}, \end{aligned}$$

$$\mathbf{E}^\Pi = \{ \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+3} \otimes \chi_{2i+2} \otimes \chi_1 \text{ for } 1 \leq i \leq (d-3)/2 \}.$$

Proof. By [34, Lemma 5.10], we know $\dim_{\mathbb{F}_2} K_3^3(H^\bullet) = d^3 - 2d$. Hence it suffices to count the number of elements in $\mathbf{B}_3^\Pi$ and to show that the set is linearly independent. We have $\#\mathbf{S}^\Pi = (d-1)d(d-1)$ since we have d choices for the middle index j and for each choice exactly one value for i and k excluded. We have $\#\mathbf{D}^\Pi = 4(d-2)(d-1)/2 = 2(d-2)(d-1)$ since we have four times $(d-1)/2$ choices for the index i and for each choice exactly two values for j excluded. We have $\#\mathbf{E}^\Pi = (d-3)/2$ since we have $(d-3)/2$ choices for the index i each furnishing one element. We have $\#\mathbf{T}^\Pi = d-1$ as there are $(d-1)/2$ choices for the index i each furnishing two triple sums, and $\#\widetilde{\mathbf{T}}^\Pi = 3(d-1)/2$ since each choice of i furnishes three elements. Thus, in total we have

$$\begin{aligned} \#\mathbf{B}_3^\Pi &= \#\mathbf{S}^\Pi + \#\mathbf{D}^\Pi + \#\mathbf{T}^\Pi + \#\widetilde{\mathbf{T}}^\Pi + \#\mathbf{E}^\Pi \\ &= d(d-1)^2 + 2(d-2)(d-1) + (d-1) + 3(d-1)/2 + (d-3)/2 \\ &= d^3 - 2d. \end{aligned}$$

It remains to show that $\mathbf{B}_3^\Pi$ is linearly independent. Assume we write the zero vector in $K_3^3(H^\bullet)$ as an $\mathbb{F}_2$ -linear combination of elements in $\mathbf{B}_3^\Pi$. Since the terms in $\mathbf{S}^\Pi$ only occur in $\mathbf{S}^\Pi$, their coefficients must be zero. Similarly, for $i \in J/2$, terms of the form $\chi_{2i} \otimes \chi_1 \otimes \chi_1$ and $\chi_1 \otimes \chi_1 \otimes \chi_{2i+1}$ each only occur once in sums in $\widetilde{\mathbf{T}}^\Pi$. Hence the coefficients of $\chi_{2i} \otimes \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i}$ and $\chi_1 \otimes \chi_1 \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i}$ in $\widetilde{\mathbf{T}}^\Pi$ must be zero. Terms of the form $\chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i}$ and $\chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1}$ only occur in sums in $\mathbf{T}^\Pi$ and $\widetilde{\mathbf{T}}^\Pi$. We have shown that the coefficients of the corresponding sums in $\widetilde{\mathbf{T}}^\Pi$ are zero. Hence the coefficients of all the terms in $\mathbf{T}^\Pi$ must be zero. For $i \in J/2$ and $j \neq 2i, 2i+1$, terms of the form $\chi_{2i} \otimes \chi_{2i+1} \otimes \chi_j$ only occur once in sums in $\mathbf{D}^\Pi$. Hence the coefficient of $(\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j$ must be zero. For

$i \in J/2$ and $j \neq 2i, 2i+1$, the only sum which contains the term $\chi_j \otimes \chi_{2i} \otimes \chi_{2i+1}$ is $\chi_j \otimes (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i})$ in $\mathbf{D}^{\text{II}}$. This implies that the coefficient of the latter must be zero. Similarly, for $i \in J/2$ and $j \neq 1, 2i+1$, the only remaining sum which contains the term $\chi_{2i+1} \otimes \chi_{2i} \otimes \chi_j$ is $(\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j$ in $\mathbf{D}^{\text{II}}$. This implies that coefficient of the latter must be zero as well. For $i \in J/2$ and $j \neq 1, 2i$, terms of the form $\chi_j \otimes \chi_{2i+1} \otimes \chi_{2i}$ only occur in the sum $\chi_j \otimes (\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i})$ in $\mathbf{D}^{\text{II}}$. This implies that coefficient of the latter must be zero, too. This shows that the coefficients of all terms in $\mathbf{D}^{\text{II}}$ must be zero. In particular, the coefficient of $(\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_1$ is zero. Thus, the term $\chi_{2i+1} \otimes \chi_{2i} \otimes \chi_1$ now only occurs in sums in $\widetilde{\mathbf{T}}^{\text{II}}$ and in $\mathbf{E}^{\text{II}}$. Now we use induction to show that the remaining coefficients are zero as well. For $i = 1$, the term $\chi_3 \otimes \chi_2 \otimes \chi_1$ only occurs in $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_3 \otimes \chi_2 + \chi_3 \otimes \chi_2 \otimes \chi_1$ in $\widetilde{\mathbf{T}}^{\text{II}}$. Thus, the coefficient of this term is zero. Now the only remaining sum where the term $\chi_1 \otimes \chi_3 \otimes \chi_2$ occurs is the sum $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_3 \otimes \chi_2 + \chi_5 \otimes \chi_4 \otimes \chi_1$ in $\mathbf{E}^{\text{II}}$. Hence the coefficient of the latter is zero. Then the only remaining sum where the term $\chi_5 \otimes \chi_4 \otimes \chi_1$ occurs is the sum $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_5 \otimes \chi_4 + \chi_5 \otimes \chi_4 \otimes \chi_1$ in $\widetilde{\mathbf{T}}^{\text{II}}$, and hence the coefficient of the latter must be zero. Now we proceed by induction to show that the coefficients of all terms in $\mathbf{E}^{\text{II}}$ and of the remaining terms in $\widetilde{\mathbf{T}}^{\text{II}}$ are zero. This proves that all coefficients must be zero and $\mathbf{B}_3^{\text{II}}$ is linearly independent. $\square$

5.2. The differential in the Koszul–Hochschild complex - Demushkin groups of type II. Let $\partial: \text{Hom}_{\mathbb{F}_2}(K_2^2(H^\bullet), H^1) \rightarrow \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ denote the differential in the complex (3) which computes the Hochschild cohomology of $H^\bullet$. We will now compute the effect of ∂ . In the following, we use notation analogous to the one introduced in Notation 4.5 and 4.6 in Section 4.2.

Lemma 5.4. *Let $i \in J/2$. Then we have the identities*

$$(\chi_{2i} \otimes \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i})^* = \partial \sigma_{2i, 2i+1'+2i+1, 2i}^{2i+1},$$

and

$$(\chi_1 \otimes \chi_1 \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i})^* = \partial \sigma_{2i, 2i+1'+2i+1, 2i}^{2i}$$

in $\text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$.

Proof. The proof is similar to the proof of Lemma 4.7. To prove the first identity, let $i \in J/2$. To simplify the notation, we write $\sigma = \sigma_{2i, 2i+1'+2i+1, 2i}^{2i+1}$. First, we compute, using the definition of σ in the final step,

$$\begin{aligned} & \partial \sigma(\chi_{2i, 1, 1} + \chi_{2i, 2i+1, 2i} + \chi_{2i+1, 2i, 2i}) \\ &= \chi_{2i} \cup \sigma(\chi_{1, 1} + \chi_{2i+1, 2i}) + \chi_{2i+1} \cup \sigma(\chi_{2i, 2i}) \\ & \quad + \sigma(\chi_{2i, 2i+1} + \chi_{2i+1, 2i}) \cup \chi_{2i} + \sigma(\chi_{2i, 1}) \cup \chi_1 = \chi_{2i+1} \cup \chi_{2i} \neq 0. \end{aligned}$$

The value of σ on other elements in $\mathbf{B}_3^{\text{II}}$ can only be non-zero if they contain the factor $\chi_{2i, 2i+1} + \chi_{2i+1, 2i}$. This may be the case for elements in $\mathbf{D}^{\text{II}}$, $\mathbf{T}^{\text{II}}$, and $\widetilde{\mathbf{T}}^{\text{II}}$. As we have seen before, $\partial \sigma$ is trivial on elements in $\mathbf{T}^{\text{II}}$ by symmetry. A similar computation as above yields $\partial \sigma(\chi_{1, 1, 2i+1} + \chi_{2i+1, 2i, 2i+1} + \chi_{2i+1, 2i+1, 2i}) = \chi_{2i+1} \cup \chi_{2i+1} = 0$. This proves the claim for $\widetilde{\mathbf{T}}^{\text{II}}$. The only terms we need to consider in $\mathbf{D}^{\text{II}}$ are those of the form $\chi_j \otimes (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i})$ and $(\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j$. However, since $j \neq 2i$, we get $\partial \sigma(\chi_j \otimes (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i})) = \chi_j \cup \chi_{2i+1} = 0$. Similarly, we get that $\partial \sigma$ vanishes on the other term as well. This

completes the proof for the first identity. The second identity follows in a similar way. $\square$

Proposition 5.5. *Let $\kappa \in \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$. Assume that κ vanishes on all elements in $\mathbf{B}_3^\Pi$ except possibly for elements of the form $\chi_{2i} \otimes \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i}$ and $\chi_1 \otimes \chi_1 \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i}$ in $\widetilde{\mathbf{T}}^\Pi$. Then $[\kappa] = 0$ in $\text{HH}^{3,-1}(H^\bullet)$.*

Proof. This follows from Lemma 5.4. $\square$

5.3. Construction of the canonical class - Demushkin groups of type II.

We will now construct the canonical class of G . Let $\text{Hom}(G, \mathbb{F}_2)$ denote the $\mathbb{F}_2$ -vector space of continuous group homomorphisms. We choose f_1 to be the identity $H^1 \rightarrow \mathcal{Z}^1 = \text{Hom}(G, \mathbb{F}_2)$ and omit it from the notation. We construct an $\mathbb{F}_2$ -linear map $f_2: R \rightarrow \mathcal{C}^1$ by defining it on each element of $\mathbf{B}_2^\Pi$ and then extend it $\mathbb{F}_2$ -linearly.

Let $A_{10} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, $A_{01} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$, and $A_{11} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ denote the matrices used in the proof of Lemma 3.11. Let $i \neq j$ and $(i, j) \neq (2k, 2k+1), (2k+1, 2k)$. We define the continuous homomorphism $\rho_{i,j}^\Pi: G \rightarrow U_3(\mathbb{F}_2)$ by the assignment $x_i \mapsto A_{10} = E_{1,2}$, $x_j \mapsto A_{01} = E_{2,3}$, and $x_s \mapsto I_3$ for $s \neq i, j$. We define the continuous map $f_2(\chi_i \otimes \chi_j): G \rightarrow \mathbb{F}_2$ by setting $f_2(\chi_i \otimes \chi_j) := e_{1,3} \circ \rho_{i,j}^\Pi$. For $i \neq 1$, we define the continuous homomorphism $\rho_{i,i}^\Pi: G \rightarrow U_3(\mathbb{F}_2)$ by the assignment $x_i \mapsto A_{11}$, and $x_j \mapsto I_3$ for $j \neq i$. We define the continuous map $f_2(\chi_i \otimes \chi_i): G \rightarrow \mathbb{F}_2$ by setting $f_2(\chi_i \otimes \chi_i) := e_{1,3} \circ \rho_{i,i}^\Pi$. Now let $1 \leq i \leq (d-1)/2$. We would like to define a continuous homomorphism $\rho_{1,1'+2i+1,2i}^\Pi: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the continuous homomorphism $\bar{\rho}_{1,1'+2i+1,2i}^\Pi: G \rightarrow \bar{U}_4(\mathbb{F}_2)$ given by $\bar{\rho}_{1,1'+2i+1,2i}^\Pi = \begin{pmatrix} 1 & \chi_1 & \chi_{2i+1} & * \\ 1 & 0 & \chi_1 & \\ & 1 & \chi_{2i} & \\ & & 1 & \end{pmatrix}$. To construct the desired lift, consider the matrices

$$A := \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad B := \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \text{and } C := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $A^2 = [B, C] = E_{1,4}$ and hence $A^2[B, C] = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{1,1'+2i+1,2i}^\Pi: G \rightarrow U_4(\mathbb{F}_2)$ by the assignment $x_1 \mapsto A$, $x_{2i} \mapsto B$, $x_{2i+1} \mapsto C$, and $x_s \mapsto I_4$ for $s \neq 1, 2i, 2i+1$. We define the desired continuous map $G \rightarrow \mathbb{F}_2$ by setting

$$f_2(\chi_{1,1} + \chi_{2i+1,2i}) := e_{1,4} \circ \rho_{1,1'+2i+1,2i}^\Pi.$$

Next, we would like to define a continuous homomorphism $\rho_{2i,2i+1'+2i+1,2i}^\Pi: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the continuous homomorphism $\bar{\rho}_{2i,2i+1'+2i+1,2i}^\Pi: G \rightarrow \bar{U}_4(\mathbb{F}_2)$

given by $\bar{\rho}_{2i,2i+1'+2i+1,2i}^\Pi = \begin{pmatrix} 1 & \chi_{2i} & \chi_{2i+1} & * \\ 1 & 0 & \chi_{2i+1} & \\ & 1 & \chi_{2i} & \\ & & 1 & \end{pmatrix}$. To construct the desired lift, consider the matrices

$$D := \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \text{and } E := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $[D, E] = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{2i,2i+1'+2i+1,2i}^\Pi$ by the assignment $x_{2i} \mapsto D$, $x_{2i+1} \mapsto E$, and $x_s \mapsto I_4$ for $s \neq 2i, 2i+1$. We define the desired continuous map $G \rightarrow \mathbb{F}_2$ by setting

$$f_2(\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) := e_{1,4} \circ \rho_{2i,2i+1'+2i+1,2i}^\Pi.$$

We now define the map $f_2: R \rightarrow \mathcal{C}^1$ on all of R by extending it $\mathbb{F}_2$ -linearly from $\mathbf{B}_2^{\text{II}}$ to R . We define the map $\Psi_3^{\text{II}}: K_3^3(H^\bullet) \rightarrow \mathcal{Z}^2$ as in (4). Taking the cohomology class of Ψ_3^{II} defines an $\mathbb{F}_2$ -linear map $\kappa_3^{\text{II}}: K_3^3(H^\bullet) \rightarrow H^2$ which represents the canonical class of G in $\text{HH}^{3,-1}(H^\bullet)$.

5.4. One more compatibility result. For the computation of the canonical class we need another compatibility result.

Lemma 5.6. *Let $T^{\text{II}}(\mathbb{F}_2)$ be the subset of $U_5(\mathbb{F}_2)$ consisting of matrices of the form $\begin{pmatrix} 1 & 0 & x & y & t \\ 0 & 1 & v & 0 & w \\ 0 & 0 & 1 & 0 & x \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$ and let $W^{\text{II}}(\mathbb{F}_2)$ denote the subset of matrices of the form $\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & v & 0 & w \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$. Then $T^{\text{II}}(\mathbb{F}_2)$ is a subgroup of $U_5(\mathbb{F}_2)$, and $W^{\text{II}}(\mathbb{F}_2)$ is a normal subgroup of $T^{\text{II}}(\mathbb{F}_2)$. Moreover, the map $\tau_5^{\text{II}}: U_4(\mathbb{F}_2) \rightarrow T^{\text{II}}(\mathbb{F}_2)/W^{\text{II}}(\mathbb{F}_2)$ defined by $\tau_5^{\text{II}}: \begin{pmatrix} 1 & x & y & t \\ 0 & 1 & 0 & x \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & x & y & t \\ 0 & 1 & * & 0 & * \\ 0 & 0 & 1 & 0 & x \\ 0 & 0 & 0 & 1 & z \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$ is an isomorphism of groups.*

Proof. This follows from a direct computation as in the proof of Lemma 4.13. $\square$

Notation 5.7. Let $n \geq 3$. For $1 \leq s_i < t_i \leq n$ for $i = 1, 2$, we let $E_{s_1, t_1}^{s_2, t_2}$ denote the matrix in $U_n(\mathbb{F}_2)$ with only non-zero entries above the diagonal in positions (s_1, t_1) and (s_2, t_2) . We note that it will be clear from the context which n we consider and we therefore omit it from notation for the matrices $E_{s_1, t_1}^{s_2, t_2}$.

Proposition 5.8. *Let $\rho_5^{\text{II}}: G \rightarrow T^{\text{II}}(\mathbb{F}_2)$ be a continuous group homomorphism satisfying $\rho_5^{\text{II}}(x_1) = E_{3,5}^{1,3}$, $\rho_5^{\text{II}}(x_{2i}) = E_{1,4}$, $\rho_5^{\text{II}}(x_{2i+1}) = E_{4,5}$, and $\rho_5^{\text{II}}(x_k) = I_5$ for $k \neq 1, 2i, 2i+1$. Then*

$$e_{1,5} \circ \rho_5^{\text{II}} = f_2(\chi_{1,1} + \chi_{2i+1,2i})$$

as continuous maps $G \rightarrow \mathbb{F}_2$.

Proof. This follows from the definition of f_2 and Lemma 5.6. $\square$

5.5. Computation of the canonical class - Demushkin groups of type II. Now we compute the values of the map representing the canonical class of G .

Proposition 5.9. *The map $\kappa_3^{\text{II}}: K_3^3(H^\bullet) \rightarrow H^2$ vanishes on all elements in $\mathbf{B}_3^{\text{II}}$ except, possibly, for the elements of the form $\chi_{2i} \otimes \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i}$ and $\chi_1 \otimes \chi_1 \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i}$ in $\widetilde{\mathbf{T}}^{\text{II}}$.*

Proof. We use the same strategy and notation as in the proof of Proposition 4.21. In fact, for elements in $\mathbf{S}^{\text{II}}$ and $\mathbf{T}^{\text{II}}$ we can use the same homomorphisms as in the proof of Proposition 4.21 since the required matrix relations hold where we note that $\chi_{1,1,1} \notin \mathbf{S}^{\text{II}}$ and $2i, 2i+1 \geq 2$ for indices in $\mathbf{T}^{\text{II}}$. This shows that κ_3^{II} vanishes on $\mathbf{S}^{\text{II}}$ and $\mathbf{T}^{\text{II}}$.

Now we consider the set $\mathbf{D}^{\text{II}}$. First, let $i \in J/2$ and $j \neq 2i, 2i+1$. We consider elements of the form $\chi_j \otimes (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i})$. We need to show that the continuous homomorphism $\bar{\rho}: G \rightarrow \overline{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_j & f_2(\chi_{j,2i}) & f_2(\chi_{j,2i+1}) & * \\ & 1 & \chi_{2i} & \chi_{2i+1} & f_2(\chi_{2i,2i+1} + \chi_{2i+1,2i}) \\ & & 1 & 0 & \chi_{2i+1} \\ & & & 1 & \chi_{2i} \\ & & & & 1 \end{pmatrix}$$

lifts to a homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$. We have $(E_{1,2})^2[E_{4,5}^{2,3}, E_{3,5}^{2,4}] = I_5$. Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_j \mapsto E_{1,2}, \quad x_{2i} \mapsto E_{4,5}^{2,3}, \quad x_{2i+1} \mapsto E_{3,5}^{2,4},$$

and $\rho(x_k) = I_5$ for $k \neq j, 2i, 2i+1$. We use Proposition 4.15 to identify $e_{1,4} \circ \rho$ with $f_2(\chi_{j,2i+1})$. By Dwyer's Theorem 3.3, the existence of ρ implies $\kappa_3^\Pi(\chi_j \otimes (\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i})) = 0$.

Next, we consider elements of the form $(\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j$. We need to show that the continuous homomorphism $\bar{\rho}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_{2i} & \chi_{2i+1} & f_2(\chi_{2i,2i+1} + \chi_{2i+1,2i}) & * \\ & 1 & 0 & \chi_{2i+1} & f_2(\chi_{2i+1,j}) \\ & & 1 & \chi_{2i} & f_2(\chi_{2i,j}) \\ & & & 1 & \chi_j \\ & & & & 1 \end{pmatrix}$$

lifts to a homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$. We have $(E_{4,5})^2[E_{3,4}^{1,2}, E_{2,4}^{1,3}] = I_5$. Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_j \mapsto E_{4,5}, \quad x_{2i} \mapsto E_{3,4}^{1,2}, \quad x_{2i+1} \mapsto E_{2,4}^{1,3},$$

and $\rho(x_k) = I_5$ for $k \neq j, 2i, 2i+1$. We use Proposition 4.15 to identify $e_{1,4} \circ \rho$ with $f_2(\chi_{2i+1,j})$. By Dwyer's Theorem 3.3, the existence of ρ implies $\kappa_3^\Pi((\chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j) = 0$.

Now, let $i \in J/2$ and $j \neq 1, 2i$. We consider elements of the form $\chi_j \otimes (\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i})$. We want to show that the continuous homomorphism $\bar{\rho}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_j & f_2(\chi_{j,1}) & f_2(\chi_{j,2i+1}) & * \\ & 1 & \chi_1 & \chi_{2i+1} & f_2(\chi_{1,1} + \chi_{2i+1,2i}) \\ & & 1 & 0 & \chi_1 \\ & & & 1 & \chi_{2i} \\ & & & & 1 \end{pmatrix}$$

lifts to a homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$. We have $(E_{2,3}^{3,5})^2[E_{4,5}, E_{2,3}] = I_5$. Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto E_{2,3}^{3,5}, \quad x_{2i} \mapsto E_{4,5}, \quad x_{2i+1} \mapsto E_{2,3}, \quad x_j \mapsto E_{1,2},$$

and $\rho(x_k) = I_5$ for $k \neq 1, 2i, 2i+1, j$. We use Proposition 4.15 to identify $e_{1,4} \circ \rho$ with $f_2(\chi_{j,2i+1})$. By Dwyer's Theorem 3.3, the existence of ρ implies $\kappa_3^\Pi(\chi_j \otimes (\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i})) = 0$.

Now, let $i \in J/2$ and $j \neq 1, 2i+1$. We consider elements of the form $(\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j$. We want to show that the continuous homomorphism $\bar{\rho}: G \rightarrow \bar{U}_5(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_1 & \chi_{2i+1} & f_2(\chi_{1,1} + \chi_{2i+1,2i}) & * \\ & 1 & 0 & \chi_1 & f_2(\chi_{1,j}) \\ & & 1 & \chi_{2i} & f_2(\chi_{2i,j}) \\ & & & 1 & \chi_j \\ & & & & 1 \end{pmatrix}$$

lifts to a homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$. We have $(E_{1,2}^{2,4})^2[E_{3,4}, E_{1,3}] = I_5$. Thus, by Lemma 3.10, we can define the continuous homomorphism $\rho: G \rightarrow U_5(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto E_{1,2}^{2,4}, \quad x_{2i} \mapsto E_{3,4}, \quad x_{2i+1} \mapsto E_{1,3}, \quad x_j \mapsto E_{4,5},$$

and $\rho(x_k) = I_5$ for $k \neq 1, 2i, 2i+1, j$. We use Proposition 4.15 to identify $e_{2,5} \circ \rho$ with $f_2(\chi_{1,j})$. By Dwyer's Theorem 3.3, the existence of ρ implies $\kappa_3^{\text{II}}((\chi_1 \otimes \chi_1 + \chi_{2i+1} \otimes \chi_{2i}) \otimes \chi_j) = 0$. This finishes the proof for the set $\mathbf{D}^{\text{II}}$.

Next, we consider the set $\widetilde{\mathbf{T}}^{\text{II}}$. Let $1 \leq i \leq (d-1)/2$. We only need to consider elements of the form $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_1$. We want to show that the continuous homomorphism $\bar{\rho}: G \rightarrow \bar{U}_6(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_1 & \chi_{2i+1} & f_2(\chi_{1,1} + \chi_{2i+1,2i}) & f_2(\chi_{1,2i+1}) & * \\ & 1 & 0 & \chi_1 & \chi_{2i+1} & f_2(\chi_{1,1} + \chi_{2i+1,2i}) \\ & & 1 & \chi_{2i} & 0 & f_2(\chi_{2i,1}) \\ & & & 1 & 0 & \chi_1 \\ & & & & 1 & \chi_{2i} \\ & & & & & 1 \end{pmatrix}$$

lifts to a homomorphism $\rho: G \rightarrow U_6(\mathbb{F}_2)$. Consider the matrix $T := \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 0 & 0 \\ & & 1 & 0 & 0 & 0 \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}$

in $U_6(\mathbb{F}_2)$. We have $T^2[E_{5,6}^{3,4}, E_{2,5}^{1,3}] = I_6$ in $U_6(\mathbb{F}_2)$. (Note, however, that $T^2 = [E_{5,6}^{3,4}, E_{2,5}^{1,3}] = E_{2,6}^{1,4} \neq I_6$ in $U_6(\mathbb{F}_2)$.) Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho: G \rightarrow U_6(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto T, \quad x_{2i} \mapsto E_{5,6}^{3,4}, \quad x_{2i+1} \mapsto E_{2,5}^{1,3},$$

and $\rho(x_k) = I_6$ for $k \neq 1, 2i, 2i+1$. We use Convention 4.20 and Proposition 4.18 to identify $e_{1,5} \circ \rho$ with $f_2(\chi_{1,2i+1})$, Proposition 4.15 to identify $e_{3,6} \circ \rho$ with $f_2(\chi_{2i,1})$, and Proposition 5.8 to identify $e_{2,6} \circ \rho$ with $f_2(\chi_{1,1} + \chi_{2i+1,2i})$. By Dwyer's Theorem 3.3, the existence of ρ implies $\kappa_3^{\text{II}}(\chi_{1,1,1} + \chi_{1,2i+1,2i} + \chi_{2i+1,2i,1}) = 0$. This proves the assertion for the set $\widetilde{\mathbf{T}}^{\text{II}}$.

Finally, we consider the set $\mathbf{E}^{\text{II}}$. Let $1 \leq i \leq (d-3)/2$. We need to show that the continuous homomorphism $\bar{\rho}: G \rightarrow \bar{U}_6(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_1 & \chi_{2i+3} & f_2(\chi_{1,1} + \chi_{2i+3,2i+2}) & f_2(\chi_{1,2i+1}) & * \\ & 1 & 0 & \chi_1 & \chi_{2i+1} & f_2(\chi_{1,1} + \chi_{2i+1,2i}) \\ & & 1 & \chi_{2i+2} & 0 & f_2(\chi_{2i+2,1}) \\ & & & 1 & 0 & \chi_1 \\ & & & & 1 & \chi_{2i} \\ & & & & & 1 \end{pmatrix}$$

lifts to a homomorphism $\rho: G \rightarrow U_6(\mathbb{F}_2)$. We consider again the matrix $T =$

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 0 & 0 \\ & & 1 & 0 & 0 & 0 \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}$$

in $U_6(\mathbb{F}_2)$. We have $T^2[E_{5,6}, E_{2,5}][E_{3,4}, E_{1,3}] = I_6$ in $U_6(\mathbb{F}_2)$. (Note that $T^2 = E_{2,6}^{1,4}$, $[E_{5,6}, E_{2,5}] = E_{2,6}$, and $[E_{3,4}, E_{1,3}] = E_{1,4}$ in $U_6(\mathbb{F}_2)$.) Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho: G \rightarrow U_6(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto T, \quad x_{2i} \mapsto E_{5,6}, \quad x_{2i+1} \mapsto E_{2,5}, \quad x_{2i+2} \mapsto E_{3,4}, \quad x_{2i+3} \mapsto E_{1,3},$$

and $\rho(x_k) = I_6$ for $k \neq 1, 2i, 2i+1, 2i+2, 2i+3$. We use Convention 4.20 and Proposition 4.18 to identify $e_{1,5} \circ \rho$ with $f_2(\chi_{1,2i+1})$, Proposition 4.15 to identify $e_{3,6} \circ \rho$ with $f_2(\chi_{2i+2,1})$, and Proposition 5.8 to identify $e_{2,6} \circ \rho$ with $f_2(\chi_{1,1} + \chi_{2i+1,2i})$. By Dwyer's Theorem 3.3, the existence of ρ implies

$$\kappa_3^{\text{II}}(\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+3} \otimes \chi_{2i+2} \otimes \chi_1) = 0$$

and proves the assertion for the set $\mathbf{E}^{\text{II}}$. This proves the proposition. $\square$

Remark 5.10. In fact, the map κ_3^{II} also vanishes on elements of the form $\chi_1 \otimes \chi_1 \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i+1} + \chi_{2i+1} \otimes \chi_{2i+1} \otimes \chi_{2i}$ in $\widetilde{\mathbf{T}}^{\text{II}}$. However, κ_3^{II} is non-trivial on elements of the form $\chi_{2i} \otimes \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i+1} \otimes \chi_{2i} + \chi_{2i+1} \otimes \chi_{2i} \otimes \chi_{2i}$ in $\widetilde{\mathbf{T}}^{\text{II}}$. This follows from the fact that the continuous homomorphism $\bar{\rho}: G \rightarrow \overline{U}_6(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_{2i} & \chi_{2i+1} & f_2(\chi_{2i,2i+1} + \chi_{2i+1,2i}) & f_2(\chi_{2i,1}) & * \\ & 1 & 0 & \chi_{2i+1} & \chi_1 & f_2(\chi_{1,1} + \chi_{2i+1,2i}) \\ & & 1 & \chi_{2i} & 0 & f_2(\chi_{2i,2i}) \\ & & & 1 & 0 & \chi_{2i} \\ & & & & 1 & \chi_1 \\ & & & & & 1 \end{pmatrix}$$

does not lift to a continuous homomorphism $G \rightarrow U_6(\mathbb{F}_2)$. For, let $T' := \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 0 & 0 & 0 \\ & & 1 & 1 & 0 & 0 \\ & & & 1 & 0 & 0 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix}$ in $U_6(\mathbb{F}_2)$. We have $(E_{5,6}^{2,5})^2 = E_{2,6}$ and $[T', E_{2,4}^{1,3}] = E_{2,6}^{1,6}$ and hence $(E_{5,6}^{2,5})^2[T', E_{2,4}^{1,3}] = I_6$ in $\overline{U}_6(\mathbb{F}_2)$, but $(E_{5,6}^{2,5})^2[T', E_{2,4}^{1,3}] = E_{1,6} \neq I_6$ in $U_6(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define a continuous group homomorphism $\bar{\rho}: G \rightarrow \overline{U}_6(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto E_{5,6}^{2,5}, \quad x_{2i} \mapsto T', \quad x_{2i+1} \mapsto E_{2,4}^{1,3},$$

and $\bar{\rho}(x_k) = I_6$ for $k \neq 1, 2i, 2i+1$, but $\bar{\rho}$ does not lift to a continuous homomorphism $G \rightarrow U_6(\mathbb{F}_2)$. By Dwyer's Theorem 3.3, this implies that, for all $i \in J/2$, we have $\kappa_3^{\text{II}}(\chi_{2i,1,1} + \chi_{2i,2i+1,2i} + \chi_{2i+1,2i,2i}) \neq 0$.

Proof of Theorem 5.1. The assertion now follows from Propositions 5.5 and 5.9. $\square$

6. DEMUSHKIN GROUPS OF TYPE III AND IV

In this section, we consider the remaining cases of pro-2 Demushkin groups. In fact, the cases of defining relation (7) or (8) can be treated at the same time. Our goal is to prove the following theorem which, by the classification of pro-2 Demushkin groups recalled in Section 3.3, concludes the proof of Theorem 1.2.

Theorem 6.1. *Let $f \geq 2$ and $d \geq 2$ be an even number. Assume that G is the pro-2 group generated by $x_1, \dots, x_d$ subject to either the single relation (7) or the single relation (8), where we allow $f = \infty$ for relation (7) and assume $d \geq 4$ for relation (8). Then the canonical class of G is trivial, and G is A_3 -formal.*

In this section, we let G be a pro-2 group as in Theorem 6.1. We use the same notation for $C^\bullet$, $\mathcal{Z}^\bullet$, $H^\bullet$, and χ_i as in Section 4. We write again $J := \{1, \dots, d\}$, $J/2 := \{1, \dots, d/2\}$, and $\tilde{J} := (J/2) \setminus \{1\} = \{2, \dots, d/2\}$.

6.1. The Koszul–Hochschild complex - type III and IV. Again, by [30, Theorem 5.2], the graded $\mathbb{F}_2$ -cohomology algebra of G is a Koszul algebra. We now provide a basis for R and $K_3^3(H^\bullet)$.

Proposition 6.2. *The $\mathbb{F}_2$ -vector subspace $R \subset H^1 \otimes H^1$ such that $H^\bullet = T(H^1)/(R)$ admits the set $\mathbf{B}_2^{\text{III}}$ as a basis where*

$$\begin{aligned} \mathbf{B}_2^{\text{III}} := & \{\chi_i \otimes \chi_j \text{ for all } (i, j) \notin \{(1, 1), (2r-1, 2r), (2r, 2r-1) : r \in J/2\}; \\ & \chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i-1}, \chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \text{ for all } i \in J/2\}. \end{aligned}$$

Proof. This follows from the description of the cup product on $H^\bullet(G, \mathbb{F}_2)$ in Lemma 3.11. $\square$

Next, we need to find a basis for $K_3^3(H^\bullet)$.

Proposition 6.3. *The $\mathbb{F}_2$ -vector space $K_3^3(H^\bullet)$ admits the set $\mathbf{B}_3^{\text{III}} := \mathbf{S}^{\text{III}} \cup \mathbf{D}^{\text{III}} \cup \widetilde{\mathbf{D}}^{\text{III}} \cup \mathbf{T}^{\text{III}} \cup \widetilde{\mathbf{T}}^{\text{III}} \cup \mathbf{E}^{\text{III}} \cup \mathbf{X}^{\text{III}}$ as a basis where*

$$\mathbf{S}^{\text{III}} = \left\{ \chi_i \otimes \chi_j \otimes \chi_k \text{ with } \begin{cases} i, k \neq 1, 2 & \text{if } j = 1 \\ i, k \neq 2r & \text{if } j = 2r - 1 \\ i, k \neq 2r - 1 & \text{if } j = 2r \end{cases} \right\},$$

$$\begin{aligned} \mathbf{D}^{\text{III}} = & \{ \chi_j \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}), \\ & (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_j \text{ for } i \in J/2, j \neq 2i - 1, 2i; \\ & \chi_j \otimes (\chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i-1}), \text{ for } i \in J/2, j \neq 1, 2, 2i - 1 \\ & (\chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_j \text{ for } i \in J/2, j \neq 1, 2, 2i \}, \end{aligned}$$

$$\begin{aligned} \mathbf{T}^{\text{III}} = & \{ \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i-1} + \chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i-1} \otimes \chi_{2i-1} \otimes \chi_{2i} \text{ for } i \in \tilde{J}/2, \\ & \chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i} \otimes \chi_{2i-1} \text{ for } i \in J/2 \}, \end{aligned}$$

$$\begin{aligned} \widetilde{\mathbf{T}}^{\text{III}} = & \{ \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_1, \\ & \chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_2, \\ & \chi_{2i-1} \otimes \chi_1 \otimes \chi_1 + \chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i-1}, \\ & \chi_1 \otimes \chi_1 \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i} \otimes \chi_{2i-1}, \text{ for } i \in \tilde{J}/2 \}, \end{aligned}$$

$$\begin{aligned} \mathbf{E}^{\text{III}} = & \{ \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1, \\ & \chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_2 \text{ for } 2 \leq i \leq d/2 - 1 \}. \end{aligned}$$

$$\begin{aligned} \mathbf{X}^{\text{III}} = & \{ \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_2 \otimes \chi_1, \\ & \chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_2, \\ & \chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_2 + \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_4 \otimes \chi_3 \otimes \chi_1, \\ & \chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_2 + \chi_2 \otimes \chi_4 \otimes \chi_3 \}. \end{aligned}$$

Proof. By [34, Lemma 5.10], we know $\dim_{\mathbb{F}_2} K_3^3(H^\bullet) = d^3 - 2d$. Hence it suffices to count the number of elements in $\mathbf{B}_3^{\text{III}}$ and to show that the set is linearly independent. To compute the number of elements in $\mathbf{S}^{\text{III}}$, we observe that we have d choices for the middle index j . For $j \geq 2$, we then have $d - 1$ choices for i and k . This yields $(d - 1)^3$ elements. For $j = 1$, we have $d - 2$ choices for i and k , which yields $(d - 2)^2$ elements. Hence we have $\#\mathbf{S}^{\text{III}} = (d - 1)^3 + (d - 2)^2$. We have $\#\mathbf{D}^{\text{III}} = 2[d/2(d - 2) + (d - 2) + (d/2 - 1)(d - 3)] = (2d - 1)(d - 2)$ since we have first $d/2$ choices for the index i with for each choice exactly two values for j excluded, and then for $i = 1$ there are two values for j excluded, and for $i \geq 2$ there are three values for j excluded; for each pair we then get two elements. We have $\#\mathbf{T}^{\text{III}} = d - 1$ as there is one element for $i = 1$ and then there are $d/2 - 1$ choices for the index $i \geq 2$ each furnishing two triple sums. Similarly, we have $\#\widetilde{\mathbf{T}}^{\text{III}} = 4(d/2 - 1) = 2d - 4$, and $\#\mathbf{E}^{\text{III}} = 2(d/2 - 2) = d - 4$ since we have $d/2 - 2$

choices for the index i each furnishing two elements. Thus, in total we have

$$\begin{aligned} \#\mathbf{B}_3^{\text{III}} &= \#\mathbf{S}^{\text{III}} + \#\mathbf{D}^{\text{III}} + \#\mathbf{T}^{\text{III}} + \#\widetilde{\mathbf{T}}^{\text{III}} + \#\mathbf{E}^{\text{III}} + \#\mathbf{X}^{\text{III}} \\ &= [(d-1)^3 + (d-2)^2] + (2d-1)(d-2) \\ &\quad + (d-1) + (2d-4) + (d-4) + 4 = d^3 - 2d. \end{aligned}$$

It remains to show that $\mathbf{B}_3^{\text{III}}$ is linearly independent. Assume we write the zero vector in $K_3^3(H^\bullet)$ as an $\mathbb{F}_2$ -linear combination of elements in $\mathbf{B}_3^{\text{III}}$. Since the terms in $\mathbf{S}^{\text{III}}$ only occur in $\mathbf{S}^{\text{III}}$, their coefficients must be zero. For $i \in \tilde{J}/2$, terms of the form $\chi_{2i-1} \otimes \chi_{2i-1} \otimes \chi_{2i}$ and $\chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i}$ only occur once in sums in $\mathbf{T}^{\text{III}}$. In addition, for $i = 1$, the term $\chi_2 \otimes \chi_2 \otimes \chi_1$ only occurs once in a sum in $\mathbf{T}^{\text{III}}$. For $i \in \tilde{J}/2$, terms of the form $\chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i-1}$ and $\chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i}$ now only occur in sums in $\widetilde{\mathbf{T}}^{\text{III}}$. This implies that the coefficients of the elements $\chi_{2i-1} \otimes \chi_1 \otimes \chi_1 + \chi_{2i-1} \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i-1}$ and $\chi_1 \otimes \chi_1 \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i} \otimes \chi_{2i-1}$ must be zero. For $i \in J/2$, terms of the form $\chi_j \otimes \chi_{2i-1} \otimes \chi_{2i}$ and $\chi_{2i-1} \otimes \chi_{2i} \otimes \chi_j$ only occur in sums in $\mathbf{D}^{\text{III}}$. Hence the coefficients of $\chi_j \otimes (\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1})$ and $(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_j$ for $i \in J/2$ and $j \neq 1, 2$ must be zero. Then, for $i \in J/2$ and $j \neq 1, 2$, terms of the form $\chi_j \otimes \chi_{2i} \otimes \chi_{2i-1}$ and $\chi_{2i} \otimes \chi_{2i-1} \otimes \chi_j$ only occur in the sums $\chi_j \otimes (\chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i-1})$ and $(\chi_1 \otimes \chi_1 + \chi_{2i} \otimes \chi_{2i-1}) \otimes \chi_j$ in $\mathbf{D}^{\text{III}}$. The coefficients of these sums therefore must be zero as well. This shows that the coefficients of all terms in $\mathbf{D}^{\text{III}}$ are zero. Next, we consider elements in $\mathbf{X}^{\text{III}}$. The only sum which contains the term $\chi_2 \otimes \chi_1 \otimes \chi_1$ is the sum $\chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_2$. Thus, the coefficient of the latter must be zero as well. Now the only remaining sum which contains the term $\chi_1 \otimes \chi_1 \otimes \chi_2$ is the sum $\chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_2 + \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_4 \otimes \chi_3 \otimes \chi_1$. Thus, the coefficient of the latter must be zero, too. Then the only remaining sum which contains the term $\chi_1 \otimes \chi_2 \otimes \chi_2$ is the sum $\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_2 + \chi_2 \otimes \chi_4 \otimes \chi_3$. Thus, the coefficient of the latter must be zero. Finally, for $\mathbf{X}^{\text{III}}$, the only remaining sum which contains the term $\chi_1 \otimes \chi_2 \otimes \chi_1$ is the sum $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_2 \otimes \chi_1$. Thus, the coefficient of the latter must be zero. This shows that the coefficients of all terms in $\mathbf{X}^{\text{III}}$ are zero. The only elements with potentially non-zero coefficients remaining are the elements in $\mathbf{E}^{\text{III}}$ and the sums $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_4 \otimes \chi_3 + \chi_4 \otimes \chi_3 \otimes \chi_1$ and $\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_2$ in $\widetilde{\mathbf{T}}^{\text{III}}$. We will now show that the coefficients of these terms also must be zero by induction on i . For $i = 2$, the only remaining sum which contains the term $\chi_4 \otimes \chi_3 \otimes \chi_1$ is the sum $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_4 \otimes \chi_3 + \chi_4 \otimes \chi_3 \otimes \chi_1$ in $\widetilde{\mathbf{T}}^{\text{III}}$. Thus, the coefficient of the latter sum must be zero. Then, for $i \geq 2$, knowing that the coefficient of $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_1$ in $\widetilde{\mathbf{T}}^{\text{III}}$ is zero, implies that the coefficient of $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1$ in $\mathbf{E}^{\text{III}}$ is zero, too. This implies that the coefficient of $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i+2} \otimes \chi_{2i+1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1$ in $\widetilde{\mathbf{T}}^{\text{III}}$ is zero as well. Proceeding by induction, this shows that the coefficients of $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_1$ in $\widetilde{\mathbf{T}}^{\text{III}}$ and of $\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1$ in $\mathbf{E}^{\text{III}}$ are zero. Using a similar induction for $\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_2$ in $\widetilde{\mathbf{T}}^{\text{III}}$ and $\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_2$ in $\mathbf{E}^{\text{III}}$ we then get that the coefficients of all elements in $\widetilde{\mathbf{T}}^{\text{III}}$ and $\mathbf{E}^{\text{III}}$ are zero. This finishes the proof. $\square$

Remark 6.4. For $d = 2$, the sets $\mathbf{D}^{\text{III}}$, $\widetilde{\mathbf{T}}^{\text{III}}$, and $\mathbf{E}^{\text{III}}$ are empty, and the set $\mathbf{B}_3^{\text{III}}$ consists of just four elements

$$\begin{aligned} \mathbf{B}_3^{\text{III}} = \{ & \chi_2 \otimes \chi_2 \otimes \chi_2, \chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_2 \otimes \chi_1, \\ & \chi_2 \otimes \chi_2 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_2, \\ & \chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_2 \}. \end{aligned}$$

6.2. The differential in the Koszul–Hochschild complex - type III and IV.

Next, we compute the effect of $\partial: \text{Hom}_{\mathbb{F}_2}(K_2^2(H^\bullet), H^1) \rightarrow \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ in the complex (3) which computes the Hochschild cohomology of $H^\bullet$. In the following, we use notation analogous to the one introduced in Notation 4.5 and 4.6 in Section 4.2. The following identities of linear maps in $\text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$ follow from direct computations analogous to the arguments in the proofs of Lemmas 4.7 and 5.4.

Lemma 6.5. *For every $i \geq 2$, we have*

$$\partial \sigma_{2i-1, 2i'+2i, 2i-1}^{2i-1} = (\chi_{1,1,2i} + \chi_{2i,2i-1,2i} + \chi_{2i,2i,2i-1})^* \in (\widetilde{\mathbf{T}}^{\text{III}})^*,$$

and

$$\partial \sigma_{2i-1, 2i'+2i, 2i-1}^{2i} = (\chi_{2i-1,1,1} + \chi_{2i-1,2i,2i-1} + \chi_{2i,2i-1,2i-1})^* \in (\widetilde{\mathbf{T}}^{\text{III}})^*.$$

Lemma 6.6. *We have*

$$\begin{aligned} \partial \sigma_{1, 2'+2, 1}^1 &= (\chi_{1,1,1} + \chi_{1,2,1})^* + (\chi_{2,1,2} + \chi_{1,2,2} + \chi_{2,4,3})^* \in (\mathbf{X}^{\text{III}})^* \\ &+ (\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1} + \chi_{2,1,2})^* \in (\mathbf{X}^{\text{III}})^* \\ &+ \sum_{2 \leq i \leq d/2} (\chi_{2,1,2} + \chi_{2,2i,2i-1} + \chi_{2i,2i-1,2})^* \in (\widetilde{\mathbf{T}}^{\text{III}})^* \\ &+ \sum_{2 \leq i \leq d/2-1} (\chi_{2,1,2} + \chi_{2,2i,2i-1} + \chi_{2i+2,2i+1,2})^* \in (\mathbf{E}^{\text{III}})^*, \end{aligned}$$

and

$$\partial \sigma_{1, 2'+2, 1}^2 = (\chi_{1,1,1} + \chi_{1,2,1})^* + (\chi_{1,1,2} + \chi_{1,2,2} + \chi_{1,1,1} + \chi_{4,3,1})^* \in (\mathbf{X}^{\text{III}})^*.$$

Lemma 6.7. *We have*

$$\partial \sigma_{1, 1'+2, 1}^1 = (\chi_{2,1,2} + \chi_{1,2,2} + \chi_{2,4,3})^* \in (\mathbf{X}^{\text{III}})^*,$$

and

$$\partial \sigma_{1, 1'+2, 1}^2 = (\chi_{1,1,2} + \chi_{1,2,2} + \chi_{1,1,1} + \chi_{4,3,1})^* \in (\mathbf{X}^{\text{III}})^*.$$

Lemma 6.8. *We have*

$$\begin{aligned} \partial \sigma_{1, 1'+4, 3}^1 &= (\chi_{1,1,2} + \chi_{1,2,2} + \chi_{1,1,1} + \chi_{4,3,1})^* \in (\mathbf{X}^{\text{III}})^* \\ &+ (\chi_{2,1,2} + \chi_{1,2,2} + \chi_{2,4,3})^* \in (\mathbf{X}^{\text{III}})^* \\ &+ (\chi_{1,1,1} + \chi_{1,4,3} + \chi_{6,5,1})^* \in (\mathbf{E}^{\text{III}})^*, i = 2, \\ &+ (\chi_{2,1,2} + \chi_{2,4,3} + \chi_{6,5,2})^* \in (\mathbf{E}^{\text{III}})^*, i = 2, \end{aligned}$$

and

$$\begin{aligned} \partial \sigma_{1, 1'+4, 3}^2 &= (\chi_{1,1,2} + \chi_{1,2,2} + \chi_{1,1,1} + \chi_{4,3,1})^* \in (\mathbf{X}^{\text{III}})^* \\ &+ (\chi_{1,1,1} + \chi_{1,4,3} + \chi_{6,5,1})^* \in (\mathbf{E}^{\text{III}})^*, i = 2. \end{aligned}$$

Lemma 6.9. *For $i \geq 3$, we have*

$$\begin{aligned} \partial\sigma_{1,1'+2i,2i-1}^1 &= (\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1)^* \in (\mathbf{E}^{\text{III}})^* \\ &\quad + (\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_2)^* \in (\mathbf{E}^{\text{III}})^*, \end{aligned}$$

and

$$\partial\sigma_{1,1'+2i,2i-1}^2 = (\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1)^* \in (\mathbf{E}^{\text{III}})^*.$$

As a consequence of these computations, we can now conclude the following result which will help us computing the canonical class of G in the next sections.

Proposition 6.10. *Let $\kappa \in \text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$. Assume that κ vanishes on all the elements in the sets $\mathbf{S}^{\text{III}}$, $\mathbf{D}^{\text{III}}$, $\mathbf{T}^{\text{III}}$ in $\mathbf{B}_3^{\text{III}}$, that κ vanishes on the elements $\chi_{1,1,1} + \chi_{1,2i,2i-1} + \chi_{2i,2i-1,1}$ in $\widetilde{\mathbf{T}^{\text{III}}}$ for every $i \geq 2$, and, for every $i \geq 2$, the value of κ on the element $\chi_{2,1,2} + \chi_{2,2i,2i-1} + \chi_{2i,2i-1,2}$ in $\widetilde{\mathbf{T}^{\text{III}}}$ agrees with the value of κ on the element $\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1} + \chi_{2,1,2}$ in $\mathbf{X}^{\text{III}}$. Then $[\kappa] = 0$ in $\text{HH}^{3,-1}(H^\bullet)$.*

Proof. By Lemma 6.5, we know that the elements $(\chi_{1,1,2i} + \chi_{2i,2i-1,2i} + \chi_{2i,2i,2i-1})^*$ and $(\chi_{2i-1,1,1} + \chi_{2i-1,2i,2i-1} + \chi_{2i,2i-1,2i-1})^*$ in $(\mathbf{T}^{\text{III}})^*$ are in the image of ∂ . By Lemma 6.9, we have, for $i \geq 3$,

$$(\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_1)^* = \partial\sigma_{1,1'+2i,2i-1}^2,$$

and

$$\begin{aligned} &(\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_{2i} \otimes \chi_{2i-1} + \chi_{2i+2} \otimes \chi_{2i+1} \otimes \chi_2)^* \\ &= \partial\sigma_{1,1'+2i,2i-1}^1 + \partial\sigma_{1,1'+2i,2i-1}^2. \end{aligned}$$

By Lemmas 6.7 and 6.8, we have

$$\begin{aligned} &(\chi_1 \otimes \chi_1 \otimes \chi_1 + \chi_1 \otimes \chi_4 \otimes \chi_3 + \chi_6 \otimes \chi_5 \otimes \chi_1)^* \\ &= \partial\sigma_{1,1'+4,3}^2 + \partial\sigma_{1,1'+2,1}^2, \end{aligned}$$

and

$$\begin{aligned} &(\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_4 \otimes \chi_3 + \chi_6 \otimes \chi_5 \otimes \chi_2)^* \\ &= \partial\sigma_{1,1'+4,3}^1 + \partial\sigma_{1,1'+4,3}^2 + \partial\sigma_{1,1'+2,1}^1. \end{aligned}$$

This shows that $(\mathbf{E}^{\text{III}})^*$ is contained in the image of ∂ . By the second parts of Lemmas 6.6 and 6.7, we have

$$(\chi_{1,1,1} + \chi_{1,2,1})^* = \partial\sigma_{1,2'+2,1}^2 + \partial\sigma_{1,1'+2,1}^2.$$

By Lemma 6.7, we have

$$\begin{aligned} &(\chi_{1,1,2} + \chi_{1,2,2} + \chi_{1,1,1} + \chi_{4,3,1})^* = \partial\sigma_{1,1'+2,1}^2, \text{ and} \\ &(\chi_{2,1,2} + \chi_{1,2,2} + \chi_{2,4,3})^* = \partial\sigma_{1,1'+2,1}^1. \end{aligned}$$

This shows that $(\mathbf{X}^{\text{III}})^*$ is in the image of ∂ , except possibly for the element $(\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1} + \chi_{2,1,2})^*$. As a consequence of the above and Lemma 6.6, we then

get

$$\begin{aligned}
& (\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1} + \chi_{2,1,2})^* + \sum_{2 \leq i \leq d/2} (\chi_{2,1,2} + \chi_{2,2i,2i-1} + \chi_{2i,2i-1,2})^* \\
&= \partial\sigma_{1,2'+2,1}^1 + \partial\sigma_{1,2'+2,1}^2 + \partial\sigma_{1,1'+2,1}^2 + \partial\sigma_{1,1'+4,3}^1 + \partial\sigma_{1,1'+4,3}^2 \\
&+ \sum_{3 \leq i \leq d/2-1} (\partial\sigma_{1,1'+2i,2i-1}^1 + \partial\sigma_{1,1'+2i,2i-1}^2).
\end{aligned}$$

This shows all the required relations. $\square$

Remark 6.11. For $d = 2$, we only get the relation that $\partial\sigma_{1,2'+2,1}^1$ is equal to the sum $(\chi_{1,1,1} + \chi_{1,2,1})^* + (\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1} + \chi_{2,1,2})^*$ in $\text{Hom}_{\mathbb{F}_2}(K_3^3(H^\bullet), H^2)$.

6.3. Construction of the canonical class - type III and IV. We will now construct the canonical class of G . Let $\text{Hom}(G, \mathbb{F}_2)$ denote the $\mathbb{F}_2$ -vector space of continuous group homomorphisms. We choose f_1 to be the identity $H^1 \rightarrow \mathcal{Z}^1 = \text{Hom}(G, \mathbb{F}_2)$ and omit it from the notation. We construct an $\mathbb{F}_2$ -linear map $f_2: R \rightarrow \mathcal{C}^1$ by defining it on each element of $\mathbf{B}_2^{\text{III}}$ and then extend it $\mathbb{F}_2$ -linearly.

Let $A_{10} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, $A_{01} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$, and $A_{11} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ denote the matrices used in the proof of Lemma 3.11. Let $i \neq j$ and $(i, j) \neq (2k-1, 2k), (2k, 2k-1)$. We define the continuous homomorphism $\rho_{i,j}^{\text{III}}: G \rightarrow U_3(\mathbb{F}_2)$ by the assignment $x_i \mapsto A_{10}$, $x_j \mapsto A_{01}$, and $x_s \mapsto I_3$ for $s \neq i, j$. We define the continuous map $f_2(\chi_i \otimes \chi_j): G \rightarrow \mathbb{F}_2$ by setting $f_2(\chi_i \otimes \chi_j) := e_{1,3} \circ \rho_{i,j}^{\text{III}}$. For $i \neq 1$, we define the continuous homomorphism $\rho_{i,i}^{\text{III}}: G \rightarrow U_3(\mathbb{F}_2)$ by the assignment $x_i \mapsto A_{11}$, and $x_j \mapsto I_3$ for $j \neq i$. We define the continuous map $f_2(\chi_i \otimes \chi_i): G \rightarrow \mathbb{F}_2$ by setting $f_2(\chi_i \otimes \chi_i) := e_{1,3} \circ \rho_{i,i}^{\text{III}}$. Next, we would like to define a continuous homomorphism $\rho_{1,1'+2,1}^{\text{III}}: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the continuous homomorphism $\bar{\rho}_{1,1'+2,1}^{\text{III}}: G \rightarrow \bar{U}_4(\mathbb{F}_2)$ given by $\bar{\rho}_{1,1'+2,1}^{\text{III}} = \begin{pmatrix} 1 & \chi_1 & \chi_2 & * \\ & 1 & 0 & \chi_1 \\ & & 1 & \chi_1 \\ & & & 1 \end{pmatrix}$. To construct the desired lift, consider the matrices

$$S := \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \text{ and } T := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $S^2 = [S, T] = E_{1,4}$ and hence $S^2[S, T] = I_4$ in $U_4(\mathbb{F}_2)$. Moreover, $S^4 = T^4 = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{1,1'+2,1}^{\text{III}}: G \rightarrow U_4(\mathbb{F}_2)$ by the assignment $x_1 \mapsto S$, $x_2 \mapsto T$, and $x_s \mapsto I_4$ for $s \neq 1, 2$. We define the desired continuous map $G \rightarrow \mathbb{F}_2$ by setting

$$f_2(\chi_{1,1} + \chi_{2,1}) := e_{1,4} \circ \rho_{1,1'+2,1}^{\text{III}}.$$

Now, let $i \in \tilde{J}/2$. We would like to define a continuous group homomorphism $\rho_{1,1'+2i,2i-1}^{\text{III}}: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the homomorphism $\bar{\rho}_{1,1'+2i,2i-1}^{\text{III}}: G \rightarrow \bar{U}_4(\mathbb{F}_2)$ given by $\bar{\rho}_{1,1'+2i,2i-1}^{\text{III}} = \begin{pmatrix} 1 & \chi_1 & \chi_{2i} & * \\ & 1 & 0 & \chi_1 \\ & & 1 & \chi_{2i-1} \\ & & & 1 \end{pmatrix}$. To construct the desired lift, consider the matrices

$$A := \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, B := \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \text{ and } C := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $A^2 = [B, C] = E_{1,4}$ and hence $A^2[B, C] = I_4$ in $U_4(\mathbb{F}_2)$. Moreover, $A^4 = B^4 = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{1,1'+2i,2i-1}^{\text{III}}: G \rightarrow U_4(\mathbb{F}_2)$

by the assignment $x_1 \mapsto A$, $x_{2i-1} \mapsto B$, $x_{2i} \mapsto C$, and $x_s \mapsto I_4$ for $s \neq 1, 2i-1, 2i$. We define the desired continuous map $G \rightarrow \mathbb{F}_2$ by setting

$$f_2(\chi_{1,1} + \chi_{2i,2i-1}) := e_{1,4} \circ \rho_{1,1'+2i,2i-1}^{\text{III}}.$$

Finally, let $i \in J/2$. We would like to define a continuous homomorphism $\rho_{2i-1,2i'+2i,2i-1}^{\text{III}}: G \rightarrow U_4(\mathbb{F}_2)$ which lifts the continuous homomorphism $\bar{\rho}_{2i-1,2i'+2i,2i-1}^{\text{III}}: G \rightarrow \bar{U}_4(\mathbb{F}_2)$

given by $\bar{\rho}_{2i-1,2i'+2i,2i-1}^{\text{III}} = \begin{pmatrix} 1 & \chi_{2i-1} & \chi_{2i} & * \\ & 1 & 0 & \chi_{2i} \\ & & 1 & \chi_{2i-1} \\ & & & 1 \end{pmatrix}$. To construct the desired lift, consider the matrices

$$D := \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \text{ and } E := \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $D^4 = E^4 = I_4$ and $[D, E] = I_4$ in $U_4(\mathbb{F}_2)$. Thus, by Lemma 3.10, we can define $\rho_{2i-1,2i'+2i,2i-1}^{\text{III}}$ by the assignment $x_{2i-1} \mapsto D$, $x_{2i} \mapsto E$, and $x_s \mapsto I_4$ for $s \neq 2i-1, 2i$. We define the desired continuous map $G \rightarrow \mathbb{F}_2$ by setting

$$f_2(\chi_{2i-1} \otimes \chi_{2i} + \chi_{2i} \otimes \chi_{2i-1}) := e_{1,4} \circ \rho_{2i-1,2i'+2i,2i-1}^{\text{III}}.$$

We now define the map $f_2: R \rightarrow \mathcal{C}^1$ on all of R by extending it $\mathbb{F}_2$ -linearly from $\mathbf{B}_2^{\text{III}}$ to R . We define the map $\Psi_3^{\text{III}}: K_3^3(H^\bullet) \rightarrow \mathcal{Z}^2$ as in (4). Taking the cohomology class of Ψ_3^{III} defines an $\mathbb{F}_2$ -linear map $\kappa_3^{\text{III}}: K_3^3(H^\bullet) \rightarrow H^2$ which represents the canonical class of G in $\text{HH}^{3,-1}(H^\bullet)$.

6.4. One more compatibility result. For the computation of the canonical class we need another compatibility result.

Lemma 6.12. *Let $V^{\text{III}}(\mathbb{F}_2)$ be the subset of $U_7(\mathbb{F}_2)$ consisting of matrices of the*

form $\begin{pmatrix} 1 & x & a & b & c & z & t \\ 0 & 1 & 0 & a & 0 & 0 & x \\ 0 & 0 & 1 & x & 0 & z & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & y \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$ and let $W^{\text{III}}(\mathbb{F}_2)$ denote the subset of matrices of the form

$\begin{pmatrix} 1 & 0 & a & b & 0 & c & 0 \\ 0 & 1 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$. Then $V^{\text{III}}(\mathbb{F}_2)$ is a subgroup of $U_7(\mathbb{F}_2)$, and $W^{\text{III}}(\mathbb{F}_2)$ is a normal

subgroup of $V^{\text{III}}(\mathbb{F}_2)$. Moreover, the map $\tau_7^{\text{III}}: U_4(\mathbb{F}_2) \rightarrow V^{\text{III}}(\mathbb{F}_2)/W^{\text{III}}(\mathbb{F}_2)$ defined

*by $\tau_7^{\text{III}}: \begin{pmatrix} 1 & x & y & t \\ 0 & 1 & 0 & x \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & x & * & * & z & * & t \\ 0 & 1 & 0 & * & 0 & 0 & x \\ 0 & 0 & 1 & x & 0 & z & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & y \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$ is an isomorphism of groups.*

Proof. This follows from a direct computation as in the proof of Lemma 4.13. $\square$

Notation 6.13. Let $n \geq 3$. For $1 \leq s_i < t_i \leq n$ and $i = 1, 2, 3, 4$, let $E_{(s_1, t_1); (s_2, t_2)}^{(s_3, t_3); (s_4, t_4)}$ denote the unique matrix in $U_n(\mathbb{F}_2)$ with non-zero entries above the diagonal in positions (s_i, t_i) . When one of the (s_i, t_i) is omitted in the notation, then this entry is zero. For example, $E_{(1,2); (2,7)}^{(3,4); (4,8)}$ is the matrix in $U_n(\mathbb{F}_2)$ with non-zero entries above the diagonal in positions $(1, 2)$, $(2, 7)$, $(3, 4)$, and $(4, 8)$, and $E_{(1,3); (2,4)}^{(7,8)}$ is the matrix in $U_n(\mathbb{F}_2)$ with non-zero entries above the diagonal in positions $(1, 3)$, $(2, 4)$, and $(7, 8)$.

Proposition 6.14. *Let $\rho_7^{\text{III}}: G \rightarrow V^{\text{III}}(\mathbb{F}_2)$ be a continuous group homomorphism satisfying $\rho_7^{\text{III}}(x_1) = E_{(1,2);(3,4)}^{(2,7)}$, $\rho_7^{\text{III}}(x_{2i-1}) = E_{(5,7)}$, $\rho_7^{\text{III}}(x_{2i}) = E_{(1,5)}^{(3,6)}$, and $\rho_7^{\text{III}}(x_k) = I_7$ for $k \neq 1, 2i-1, 2i$. Then*

$$e_{1,7} \circ \rho_7^{\text{III}} = f_2(\chi_{1,1} + \chi_{2i,2i-1})$$

as continuous maps $G \rightarrow \mathbb{F}_2$.

Proof. This follows from the definition of f_2 and Lemma 6.12. $\square$

6.5. Computation of the canonical class - type III and IV. Now we determine the values of the map representing the canonical class of G .

Proposition 6.15. *For every $i \geq 2$, the values of κ_3^{III} on the element $\chi_2 \otimes \chi_1 \otimes \chi_2 + \chi_2 \otimes \chi_2 \otimes \chi_{2i-1} + \chi_{2i} \otimes \chi_{2i-1} \otimes \chi_2$ in $\widetilde{\mathbf{T}}^{\text{III}}$ and on the element $\chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_2$ in $\mathbf{X}^{\text{III}}$ agree.*

Proof. Let $i \geq 2$. It suffices to show that κ_3^{III} vanishes on the sum of the two elements. That is, after removing $\chi_{2,1,2}$ which appears twice, we want to show that

$$(14) \quad \kappa_3^{\text{III}}(\chi_{2,1,1} + \chi_{2,2i,2i-1} + \chi_{1,2,1} + \chi_{1,1,2} + \chi_{2i,2i-1,2}) = 0.$$

To do so, we will show that the continuous group homomorphism $\bar{\rho}: G \rightarrow \overline{U}_8(\mathbb{F}_2)$ given by

$$\bar{\rho} = \begin{pmatrix} 1 & \chi_1 & \chi_2 & f_2(\chi_{1,2} + \chi_{2,1}) & \chi_{2i} & f_2(\chi_{2,2i}) & f_2(\chi_{1,1} + \chi_{2i,2i-1}) & * \\ & 1 & 0 & \chi_2 & 0 & 0 & \chi_1 & f_2(\chi_{1,2} + \chi_{2,1}) \\ & & 1 & \chi_1 & 0 & \chi_{2i} & 0 & f_2(\chi_{1,1} + \chi_{2i,2i-1}) \\ & & & 1 & 0 & 0 & 0 & \chi_1 \\ & & & & 1 & 0 & \chi_{2i-1} & f_2(\chi_{2i-1,2}) \\ & & & & & 1 & 0 & \chi_{2i-1} \\ & & & & & & 1 & \chi_2 \\ & & & & & & & 1 \end{pmatrix}$$

lifts to a continuous group homomorphism $\rho: G \rightarrow U_8(\mathbb{F}_2)$. Let $A := E_{(1,2);(2,7)}^{(3,4);(4,8)}$, and $B := E_{(1,3);(2,4)}^{(7,8)}$. Let $C := E_{5,7}^{6,8}$ and $D := E_{1,5}^{3,6}$. We have $A^4 = C^4 = I_8$ and $A^2[A, B][C, D] = I_8$ in $U_8(\mathbb{F}_2)$. (Note, however, that $A^2 = [C, D] = E_{1,7}^{3,8} \neq I_8$ and $[A, B] = I_8$ in $U_8(\mathbb{F}_2)$.) Thus, by Lemma 3.10, we can define a continuous group homomorphism $\rho: G \rightarrow U_8(\mathbb{F}_2)$ by the assignment

$$x_1 \mapsto A, \quad x_2 \mapsto B, \quad x_{2i-1} \mapsto C, \quad x_{2i} \mapsto D,$$

and $\rho(x_k) = I_8$ for $k \neq 1, 2, 2i-1, 2i$. We use Convention 4.20 and Proposition 6.14 to identify $e_{1,7} \circ \rho$ with $f_2(\chi_{1,1} + \chi_{2i,2i-1})$, and we use an analogue of Proposition 6.14 to identify $e_{3,8} \circ \rho$ with $f_2(\chi_{1,1} + \chi_{2i,2i-1})$. Moreover, we use analogues of Propositions 4.18 and 5.8 to identify $e_{1,5} \circ \rho$ with $f_2(\chi_{2,2i})$, $e_{2,8} \circ \rho$ with $f_2(\chi_{1,2} + \chi_{2,1})$, and $e_{5,8} \circ \rho$ with $f_2(\chi_{2i-1,2})$. By Dwyer's Theorem 3.3, the existence of ρ implies (14). $\square$

Remark 6.16. We can determine the value of κ_3^{III} on $\chi_1 \otimes \chi_1 \otimes \chi_2 + \chi_1 \otimes \chi_2 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_1 + \chi_2 \otimes \chi_1 \otimes \chi_2$ as follows. We define the matrices

$$A = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ & 1 & 0 & 1 & 1 & 0 \\ & & 1 & 0 & 1 & 0 \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix} \text{ and } B = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ & 1 & 0 & 0 & 0 & 0 \\ & & 1 & 1 & 0 & 0 \\ & & & 1 & 0 & 0 \\ & & & & 1 & 1 \\ & & & & & 1 \end{pmatrix}.$$

We have $A^4 = I_6$ and $A^2 \cdot [A, B] = E_{1,6}$ in $U_6(\mathbb{F}_2)$, i.e., we have $A^2 \cdot [A, B] = I_6$ in $\overline{U}_6(\mathbb{F}_2)$, but $A^2 \cdot [A, B] \neq I_6$ in $U_6(\mathbb{F}_2)$. This shows that the assignment $x_1 \mapsto A$ and $x_2 \mapsto B$ defines a continuous group homomorphism $\bar{\rho}: G \rightarrow \overline{U}_6(\mathbb{F}_2)$ which does not

lift to a continuous group homomorphism $\rho: G \rightarrow U_6(\mathbb{F}_2)$. By Dwyer's Theorem 3.3, this implies $\kappa_3^{\text{III}}(\chi_{1,1,2} + \chi_{1,2,1} + \chi_{2,1,1} + \chi_{2,1,2}) \neq 0$. In fact, we can also use Dwyer's Theorem to show $\kappa_3^{\text{III}}(\chi_{1,1,1} + \chi_{1,2,1}) \neq 0$.

Proposition 6.17. *The canonical class of G is trivial.*

Proof. By Proposition 6.10 and Proposition 6.15, it suffices to show that κ_3^{III} vanishes on the elements $\chi_{1,1,1} + \chi_{1,2i,2i-1} + \chi_{2i,2i-1,1}$ in $\widetilde{\mathbf{T}}^{\text{III}}$ for every $i \geq 2$, and that κ_3^{III} vanishes on $\mathbf{S}^{\text{III}}$, $\mathbf{D}^{\text{III}}$, and $\mathbf{T}^{\text{III}}$. Both claims follow from the arguments used in the proof of Propositions 4.21 and 5.9, where we only need to replace the assignments of x_{2i} and x_{2i+1} with assignments of x_{2i-1} and x_{2i} , respectively. Here we note that every matrix $M \in U_n(\mathbb{F}_2)$ we use in Sections 4, 5, and 6 satisfies $M^4 = I_n$ in $U_n(\mathbb{F}_2)$. Hence an additional factor M^{2^f} in a matrix relation of the form (9), corresponding to the additional factor $x_1^{2^f}$ or $x_3^{2^f}$ in the defining relation for G , does not impose an obstruction for defining a continuous group homomorphism $G \rightarrow U_n(\mathbb{F}_2)$. $\square$

Proof of Theorem 6.1. The assertion follows from Propositions 2.15 and 6.17. $\square$

REFERENCES

- [1] D. Benson, H. Krause and S. Schwede, *Realizability of modules over Tate cohomology*, Trans. Amer. Math. Soc. **356** (2004), no. 9, 3621–3668.
- [2] U. Buijs, J. M. Moreno-Fernández and A. Murillo, *A_∞ -structures and Massey products*, Mediterr. J. Math. **17**, 31 (2020).
- [3] S. P. Demuškin, *The group of a maximal p -extension of a local field*, Izv. Akad. Nauk SSSR Ser. Mat. **25** (1961), 329–346.
- [4] S. P. Demuškin, *On 2-extensions of a local field*, Sibirsk. Mat. Ž. **4** (1963), 951–955.
- [5] E. X. Djurhuus and G. Quick, *Non-realizability of a triple Massey product for the algebra $\mathbb{F}_2[a, b, c]/(ab, bc)$* , New York J. Math. **32** (2026), 349–360.
- [6] W. G. Dwyer, *Homology, Massey products and maps between groups*, J. Pure Appl. Algebra **6** (1975), 177–190.
- [7] I. Efrat, *Small maximal pro- p Galois groups*, Manuscripta Math. **95** (1998), no. 2, 237–249.
- [8] I. Efrat and E. Matzri, *Triple Massey products and absolute Galois groups*, J. Eur. Math. Soc. **19** (2017), no. 12, 3629–3640.
- [9] P. Guillot, J. Mináč and A. Topaz, *Four-fold Massey products in Galois cohomology*, Compos. Math. **154** (2018), no. 9, 1921–1959, with an appendix by O. Wittenberg.
- [10] Y. Harpaz and O. Wittenberg, *The Massey vanishing conjecture for number fields*, Duke Math. J. **172** (2023), no. 1, 1–41.
- [11] M. Hopkins and K. Wickelgren, *Splitting varieties for triple Massey products*, J. Pure Appl. Algebra **219** (2015), 1304–1319.
- [12] B. Jacob, R. Ware, *A recursive description of the maximal pro-2 Galois group via Witt rings*, Math. Z. **200** (1989), no. 3, 379–396.
- [13] T. V. Kadeishvili, *The algebraic structure in the homology of an A_∞ -algebra* (Russian, English summary) Soobshch. Akad. Nauk Gruz. SSR **108** (1982), no. 2, 249–252 (1983).
- [14] T. V. Kadeishvili, *The structure of the A_∞ -algebra, and the Hochschild and Harrison cohomologies* (Russian), Trudy Tbiliss. Mat. Inst. Razmadze Akad. Nauk Gruz. SSR **91** (1988), 19–27 (an English version is available at arXiv:math/0210331).
- [15] T. V. Kadeishvili, *A_∞ -algebra Structure in Cohomology and its Applications*, Lecture notes 2023, available at <https://doi.org/10.48550/arXiv.2307.10300>.
- [16] B. Keller, *Introduction to A -infinity algebras and modules*, Homology Homotopy Appl. **3** (2001), 1–35.
- [17] J. P. Labute, *Classification of Demushkin groups*, Canadian J. Math. **19** (1967), 106–132.
- [18] Y. H. J. Lam, Y. Liu, R. Sharifi, P. Wake and J. Wang, *Generalized Bockstein maps and Massey products*, Forum Math. Sigma **11** (2023), Paper No. e5, 41 pp.

- [19] C. Maire, J. Mináč, R. Ramakrishna and N. D. Tân, *On the strong Massey property for number fields*, preprint, arXiv:2409.01028v1, 2024.
- [20] E. Matzri, *Triple Massey products and Galois cohomology*, preprint, arXiv:1411.4146 (2014).
- [21] E. Matzri, *Higher triple Massey products and symbols*, J. Algebra **527** (2019), 136–146.
- [22] S. A. Merkulov, *Strong homotopy algebras of a Kähler manifold*, Internat. Math. Res. Notices (3) (1999), 153–164.
- [23] A. Merkurjev and F. Scavia, *Degenerate fourfold Massey products over arbitrary fields*, J. Eur. Math. Soc. **28** (2026), no. 8, 3499–3537.
- [24] A. Merkurjev and F. Scavia, *The Massey Vanishing Conjecture for fourfold Massey products modulo 2*, Ann. Sci. Éc. Norm. Supér. (4) **58** (2025), no. 3, 589–606.
- [25] A. Merkurjev and F. Scavia, *Non-formality of Galois cohomology modulo all primes*, Compos. Math. **161** (2025), no. 4, 831–858.
- [26] J. Mináč and N. D. Tân, *The kernel unipotent conjecture and Massey products on an odd rigid field* (with an appendix by I. Efrat, J. Mináč and N. D. Tân), Adv. Math. **273** (2015), 242–270.
- [27] J. Mináč and N. D. Tân, *Counting Galois $\mathbb{U}_4(\mathbb{F}_p)$ -extensions using Massey products*, J. Number Theory **176** (2017), 76–112.
- [28] J. Mináč and N. D. Tân, *Triple Massey products vanish over all fields*, J. Lond. Math. Soc. (2) **94** (2016), no. 3, 909–932.
- [29] J. Mináč and N. D. Tân, *Triple Massey products and Galois theory*, J. Eur. Math. Soc. **19** (2017), 255–284.
- [30] J. Mináč, F. W. Pasini, C. Quadrelli and N. D. Tân, *Koszul algebras and quadratic duals in Galois cohomology*, Adv. Math. **380** (2021), Paper No. 107569, 49 pp.
- [31] F. Muro, *Derived universal Massey products*, Homology Homotopy Appl. **25** (2023), no. 1, 189–218.
- [32] J. Neukirch, A. Schmidt and K. Wingberg, *Cohomology of number fields*, Second edition, Grundlehren Math. Wiss., vol. **323**, Springer-Verlag, Berlin, 2008.
- [33] A. Pál and G. Quick, *Real projective groups are formal*, Math. Ann. **392** (2025), 1833–1876.
- [34] A. Pál and G. Quick, *A₃-formality for Demushkin groups at odd primes*, preprint, arXiv:2601.07551 (2026).
- [35] A. Pál and E. Szabó, *The strong Massey vanishing conjecture for fields with virtual cohomological dimension at most 1*, Israel J. Math. **272** (2026), no. 1, 153–190.
- [36] A. Polishchuk and L. Positselski, *Quadratic algebras*, University Lecture Series, 37. American Mathematical Society, Providence, RI, 2005. xii+159 pp.
- [37] L. Positselski, *Mixed Artin–Tate motives with finite coefficients*, Moscow Math. J. **11** (2011), 317–402.
- [38] L. Positselski, *Koszulity of cohomology = $K(\pi, 1)$ -ness + quasi-formality*, J. Algebra **483** (2017), 188–229.
- [39] C. Quadrelli, *Massey products in Galois cohomology and the Elementary Type Conjecture*, J. Number Theory **258** (2024), 40–65.
- [40] P. Seidel and R. Thomas, *Braid group actions on derived categories of coherent sheaves*, Duke Math. J. **108** (2001), no. 1, 37–108.
- [41] J.-P. Serre, *Structure de certains pro- p -groupes (d’après Demuškin)*, Séminaire Bourbaki, vol. **8**, Soc. Math. France, Paris, 1995, pp. 145–155, Exp. No. 252.
- [42] J.-P. Serre, *Galois cohomology*, Springer Monographs in Mathematics, Springer, Berlin-New York, 1997.

MATHEMATICAL INSTITUTE, EÖTVÖS LORÁND UNIVERSITY, H-1117 BUDAPEST, HUNGARY
 Email address: ambrus.pal@ttk.elte.hu

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU, NO-7491 TRONDHEIM, NORWAY
 Email address: gereon.quick@ntnu.no