



Stable étale realization and étale cobordism

Gereon Quick

Mathematisches Institut, WWU Münster, Einsteinstr. 62, 48149 Münster, Germany

Received 14 August 2006; accepted 20 March 2007

Available online 27 March 2007

Communicated by Michael J. Hopkins

Abstract

We show that there is a stable homotopy theory of profinite spaces and use it for two main applications. On the one hand we construct an étale topological realization of the stable \mathbb{A}^1 -homotopy theory of smooth schemes over a base field of arbitrary characteristic in analogy to the complex realization functor for fields of characteristic zero.

On the other hand we get a natural setting for étale cohomology theories. In particular, we define and discuss an étale topological cobordism theory for schemes. It is equipped with an Atiyah–Hirzebruch spectral sequence starting from étale cohomology. Finally, we construct maps from algebraic to étale cobordism and discuss algebraic cobordism with finite coefficients over an algebraically closed field after inverting a Bott element.

© 2007 Elsevier Inc. All rights reserved.

MSC: 14F42; 14F35

Keywords: \mathbb{A}^1 -homotopy theory of schemes; Étale topological type; Algebraic cobordism; Étale cohomology; Profinite spaces; Stable model structure


1. Introduction

For the proof of the Milnor Conjecture Voevodsky invented algebraic cobordism, a new cohomology theory for schemes represented by the Thom spectrum in his new framework of \mathbb{A}^1 -homotopy theory. Using the known topological realization functor for schemes over a field of characteristic zero he linked his theory to the classical one by showing that the complex topological realization of the algebraic cobordism theory yields a map to complex cobordism.

E-mail address: gquick@math.uni-muenster.de.

Independently, Levine and Morel constructed in a geometric way an algebraic cobordism theory as the universal oriented cohomology theory on smooth schemes and used it to prove Rost’s Degree Formula in characteristic zero.

Looking at the following diagram of cohomology theories



gives rise to the following questions. Is there an étale cobordism theory that fits into this picture? Can the étale theories be described in some kind of stable homotopy theory? Can one replace the complex realization functor in characteristic zero by a topological realization functor on the stable motivic category for arbitrary characteristics? Is there a statement similar to the result of Thomason that algebraic and étale cobordism with finite coefficients agree after inverting a Bott element? In this paper we give an answer to these questions.

Sending a scheme to its étale topological type, first constructed by Artin–Mazur and Friedlander, gives naturally rise to cohomology theories. The idea is due to Eric Friedlander who constructed in [13] a first version of an étale topological K-theory for schemes, which had been developed further by Dwyer and Friedlander in [11]. This theory turned out to be a powerful tool for the study of algebraic K-theory with finite coefficients. In particular, Thomason proved in his famous paper [34] that algebraic K-theory with finite coefficients agrees with étale K-theory after inverting a Bott element.

Čech and étale homotopy theory of schemes had also been studied by Cox [7] and Edwards–Hastings [12] in the 1970s. At the end of the 1970s, in [33] Victor P. Snaith had constructed a p -adic cobordism theory for schemes. His approach is close to the definition of algebraic K-theory by Quillen. He defines for every scheme V over \mathbb{F}_q , a topological cobordism spectrum $\underline{AF}_{q,V}$. The homotopy groups of this spectrum are the p -adic cobordism groups of V . He has calculated these groups for projective bundles, Severi–Brauer schemes and other examples.

In the beginning of the 1980s, Roy Joshua had already studied a version of étale (co-)bordism as the generalized (co-)homology theory represented by a mod ℓ -variant of MU on the étale homotopy type as in Friedlander’s book [14], where Friedlander also shows the existence of an Atiyah–Hirzebruch spectral sequence for such theories. Joshua had also defined an algebraic bordism theory in analogy to the topological bordism relation and constructed a map from algebraic to étale bordism using the Thom–Pontrjagin construction and tubular neighbourhoods for geometrically unbranched projective varieties over an algebraically closed field of arbitrary characteristic. Joshua did not publish these ideas, since the interest in algebraic cobordism only arose about ten years later. But his geometric constructions and his application to a Spanier–Whitehead duality in étale homotopy theory can be found in [23].

In the present paper, we also follow Friedlander’s idea, but in a slightly different setting. We consider the profinite completion $\hat{\text{Ét}}$ of Friedlander’s étale topological type functor of [14] with values in the category of simplicial profinite sets $\hat{\mathcal{S}}$. Based on the work of Hovey [16] and Morel [27], we construct a stable homotopy category $\hat{\mathcal{SH}}$ for simplicial profinite sets for any

fixed prime number ℓ . This is one possible category for the study of étale topological cohomology theories. We define general cohomology theories such as étale K-theory, étale Morava K-theory and étale cobordism as the theories represented by profinite spectra such as $K\hat{U}$, $K\hat{(n)}$ and $M\hat{U}$, respectively. This approach unifies different étale theories and coincides with the well-known theories of étale cohomology and étale K-theory with finite coefficients over a separably closed field, while étale cobordism is a new theory, which will be discussed in this paper.

An advantage of this approach is the immediate construction of an Atiyah–Hirzebruch spectral sequence for smooth schemes over a field starting from étale cohomology and converging to étale cobordism with finite coefficients.

For a base field without ℓ -torsion and with finite ℓ -cohomological dimension, for example a separably closed, a finite or a local field, étale cobordism in even degrees is an oriented cohomology theory, i.e. it has Chern classes. Since the algebraic cobordism of Levine and Morel [25] is universal for such functors, there is a canonical map from this algebraic to étale cobordism in even degrees.

For the comparison with Voevodsky’s theory [35], we have to extend the étale topological type functor to the stable homotopy category of smooth schemes over an arbitrary base field. Isaksen [18] and Schmidt [32] extended independently the étale topological type functor to the unstable \mathbb{A}^1 -homotopy category. One of our main results is the existence of an étale realization on the *stable* \mathbb{A}^1 -homotopy category.

Theorem 1. *Let k be an arbitrary base field. The functor $\hat{\text{Ét}}$ induces an étale realization of the stable motivic homotopy category of \mathbb{P}^1 -spectra:*

$$L\hat{\text{Ét}}: \mathcal{SH}(k) \rightarrow \hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Ét}}k.$$

In the theorem $\hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Ét}}k$ denotes the stable homotopy category of profinite S^2 -spectra over $\hat{\text{Ét}}k$. In particular, the functor $L\hat{\text{Ét}}$ is equal to $\hat{\text{Ét}}$ on the suspension spectra of smooth schemes and sends stable \mathbb{A}^1 -equivalences between suspension spectra of smooth schemes to isomorphisms in $\hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Ét}}k$. Over a separably closed base field the image of the motivic Thom spectrum is even canonically isomorphic in the stable profinite homotopy category to the profinite $M\hat{U}$ -spectrum.

If k has no ℓ -torsion and has finite ℓ -cohomological dimension, this realization yields a canonical map from algebraic to étale cobordism. The above map fits into a commutative diagram with the map from the Levine–Morel theory.

Furthermore, we show that the absolute Galois group of the base field acts trivially on the cobordism of the separable closure of the field. Together with the Atiyah–Hirzebruch spectral sequence this allows us to determine the étale cobordism of local fields.

Finally, there is a candidate for a Bott element in Voevodsky’s theory that is mapped to an invertible element in étale cobordism. Together with the Atiyah–Hirzebruch spectral sequence for étale cobordism and the announced motivic Atiyah–Hirzebruch spectral sequence for algebraic cobordism of Hopkins and Morel, the results of Levine [24] can be used to show that, at least over an algebraically closed field, algebraic and étale cobordism with finite coefficients agree after inverting a Bott element.

Conditional Theorem 2. *Let X be a smooth scheme of finite type over an algebraically closed field k with $\text{char} k \neq \ell$. Let $\beta \in MGL^{0,1}(k; \mathbb{Z}/\ell^v)$ be the Bott element. If we assume the exis-*

tence and convergence of an Atiyah–Hirzebruch spectral sequence from motivic cohomology to algebraic cobordism, then there is an isomorphism

$$\phi : \left(\bigoplus_{p,q} MGL^{p,q}(X; \mathbb{Z}/\ell^v) \right) [\beta^{-1}] \xrightarrow{\cong} \bigoplus_{p,q} \hat{M}U_{\acute{e}t}^p(X; \mathbb{Z}/\ell^v).$$

A similar result should hold over an arbitrary base field. But over a base field which is not separably closed, the proposed definition of étale cobordism might not be the best possible. The problem is that this homotopical construction does not reflect the twists in étale cohomology. In order to solve this problem, Morel suggested to the author to study maps in the homotopy theory relative over the base $\hat{E}tk$ and to define étale cobordism via $\hat{E}t(MGL)$ instead of $\hat{M}U$. Over a separably closed field both approaches agree. The author hopes to be able to discuss this in future work.

The content of the paper is as follows. In the first section we construct the stable homotopy category $\hat{\mathcal{S}}\mathcal{H}$ of profinite spaces and discuss generalized cohomology theories of profinite spaces. In the next section, $\hat{\mathcal{S}}\mathcal{H}$ will turn out to be the target category for the stable étale topological realization functor. It also yields a natural frame work for the study of generalized étale cohomology theories such as étale cobordism, which we discuss in the fourth section. In the last part we will compare algebraic and étale cobordism and will discuss the conditional theorem above.

2. Profinite spectra

2.1. Profinite spaces

We want to construct the target category for the stable étale realization, the stable homotopy category of profinite spaces. Therefore, we recall some facts about profinite spaces and their homotopy groups.

Let \mathcal{E} denote the category of sets and let \mathcal{F} be the full subcategory of \mathcal{E} whose objects are finite sets. Let $\hat{\mathcal{E}}$ be the category of compact and totally disconnected topological spaces. We may identify \mathcal{F} with a full subcategory of $\hat{\mathcal{E}}$ in the obvious way. The limit functor $\lim : \text{pro-}\mathcal{F} \rightarrow \hat{\mathcal{E}}$, which sends a pro-object X of \mathcal{F} to the limit in $\hat{\mathcal{E}}$ of the diagram corresponding to X , is an equivalence of categories.

We denote by $\hat{\mathcal{S}}$ (respectively \mathcal{S}) the category of simplicial profinite sets (respectively simplicial sets). The objects of $\hat{\mathcal{S}}$ (respectively \mathcal{S}) will be called *profinite spaces* (respectively *spaces*). The forgetful functor $\hat{\mathcal{E}} \rightarrow \mathcal{E}$ admits a left adjoint $(\hat{\cdot}) : \mathcal{E} \rightarrow \hat{\mathcal{E}}$. It induces a functor $(\hat{\cdot}) : \mathcal{S} \rightarrow \hat{\mathcal{S}}$, which is called *profinite completion*. It is left adjoint to the forgetful functor $|\cdot| : \hat{\mathcal{S}} \rightarrow \mathcal{S}$ which sends a profinite space to its underlying simplicial set.

For a profinite space X we define the set $\mathcal{R}(X)$ of simplicial open equivalence relations on X . An element R of $\mathcal{R}(X)$ is a simplicial profinite subset of the product $X \times X$ such that, in each degree n , R_n is an equivalence relation on X_n and an open subset of $X_n \times X_n$. It is ordered by inclusion. For every element R of $\mathcal{R}(X)$, the quotient X/R is a simplicial finite set and the map $X \rightarrow X/R$ is a map of profinite spaces. When we consider the limit $\lim_{R \in \mathcal{R}(X)} X/R$ in $\hat{\mathcal{S}}$, the map $X \rightarrow \lim_{R \in \mathcal{R}(X)} X/R$ is an isomorphism, cf. [27, Lemme 1].

Let X be a profinite space. The continuous cohomology $H^*(X; \pi)$ of X with coefficients in the profinite abelian group π is defined as the cohomology of the complex $C^*(X; \pi)$ of continuous cochains of X with values in π , i.e. $C^n(X; \pi)$ denotes the set $\text{Hom}_{\hat{\mathcal{E}}}(X_n, \pi)$ of continuous

maps $\alpha : X_n \rightarrow \pi$ and the differentials $\delta^n : C^n(X; \pi) \rightarrow C^{n+1}(X; \pi)$ are the morphisms associating to α the map $\sum_{i=0}^{n+1} \alpha \circ d_i$, where d_i denotes the i th face map of X . If π is a finite abelian group and Z a simplicial set, then the cohomologies $H^*(Z; \pi)$ and $H^*(\hat{Z}; \pi)$ are canonically isomorphic.

Let ℓ be a fixed prime number. Fabien Morel has shown in [27] that the category $\hat{\mathcal{S}}$ can be given the structure of a closed model category. The weak equivalences are the maps inducing isomorphisms in continuous cohomology with coefficients \mathbb{Z}/ℓ ; the cofibrations are the degree-wise monomorphisms. The homotopy category is denoted by $\hat{\mathcal{H}}$. Morel has also given an explicit construction of fibrant replacements in $\hat{\mathcal{S}}$, cf. [27, 2.1]. It is based on the \mathbb{Z}/ℓ -completion functor of [5]. We denote the fibrant replacement in $\hat{\mathcal{S}}$ of a profinite space X by \hat{X}^ℓ and call it the ℓ -completion of X . If Y is a simplicial set, we also denote by \hat{Y}^ℓ the ℓ -completion of its profinite completion \hat{Y} .

The category $\hat{\mathcal{S}}$ has a pointed analogue $\hat{\mathcal{S}}_*$ whose objects are maps $* \rightarrow X$ in $\hat{\mathcal{S}}$, where $*$ denotes the constant simplicial set equal to a point. Its morphisms are maps in $\hat{\mathcal{S}}$ that respect the base points. The forgetful functor $\hat{\mathcal{S}}_* \rightarrow \hat{\mathcal{S}}$ has a left adjoint, which consists in adding a disjoint base point $X \mapsto X_+$. The category $\hat{\mathcal{S}}_*$ has the obvious induced closed model category structure. The product for two pointed profinite spaces X and Y in $\hat{\mathcal{S}}_*$ is the smash product $X \wedge Y \in \hat{\mathcal{S}}_*$, defined in the usual way as the quotient $(X \times Y)/(X \vee Y)$ in $\hat{\mathcal{S}}_*$. One can show that (smash) products are compatible with ℓ -completion in $\hat{\mathcal{H}}$ (respectively $\hat{\mathcal{H}}_*$).

Morel has shown that the categories $\hat{\mathcal{S}}$ and $\hat{\mathcal{S}}_*$ have natural simplicial structures, see [27]. In particular, for a pointed profinite space Y , there are profinite mapping spaces $\text{hom}_*(W, Y)$ for every pointed simplicial finite set W . The simplicial profinite set $\text{hom}_*(W, Y)$ is defined in degree n by the set $\text{Hom}_{\hat{\mathcal{S}}}(W \wedge \Delta[n]_+, Y)$, whose profinite structure is inherited from Y by considering it as the limit over the integers s and the elements R of $\mathcal{R}(Y)$ of the finite sets $\text{Hom}_{\hat{\mathcal{S}}}(\text{sk}_s W \wedge \Delta[n]_+, Y/R)$. Here, $\text{sk}_s W$ denotes the s -skeleton of W . As an application, let S^1 be the simplicial finite set $\Delta[1]/\partial\Delta[1]$. As usual we set $\hat{\Omega}X := \text{hom}_*(S^1, X)$. In particular, $\hat{\Omega}X$ is fibrant if X is fibrant. This adjunction may be extended to the homotopy category $\hat{\mathcal{H}}_*$. For every profinite space X and every profinite space Y , there is a natural bijection $\text{Hom}_{\hat{\mathcal{H}}_*}(S^1 \wedge X, Y) \cong \text{Hom}_{\hat{\mathcal{H}}_*}(X, \hat{\Omega}Y)$. Since $S^1 \wedge X$ is a cogroup object and $\hat{\Omega}X$ a group object in $\hat{\mathcal{H}}_*$, we conclude that the previous bijection of maps in $\hat{\mathcal{H}}_*$ is in fact an isomorphism of groups, cf. [8].

There are several possibilities to define the homotopy groups of a pointed profinite space. One could define $\pi_k X$ to be the homotopy group $\pi_k|\hat{X}^\ell|$ of the underlying simplicial set of the fibrant replacement in $\hat{\mathcal{S}}_*$ of X . Dehon suggests the following approach in [8, Section 1.3]. If X is a pointed fibrant profinite space, we define $\pi_0 X$ to be the coequalizer in $\hat{\mathcal{E}}$ of the diagram $d_0, d_1 : X_1 \rightrightarrows X_0$; and $\pi_k X$ for $k \geq 1$ is defined to be the group $\pi_0 \hat{\Omega}^k X$. One can easily see that $\pi_k X$ has a natural structure of a pro- ℓ -group for every $k \geq 1$. Both definitions agree since $\pi_k(\hat{\Omega}X) = \pi_{k+1}X$.

We conclude this discussion with a collection of results on the homotopy groups of the ℓ -completion $\hat{X}^\ell \in \hat{\mathcal{S}}$ of a simplicial set X , which can be deduced from the methods of [5, VI, §5].

Proposition 3. *Let X be a pointed connected simplicial set.*

(1) *We have an isomorphism*

$$\pi_1(\hat{X}^\ell) \cong \pi_1(\hat{X})^\ell.$$

In particular, if $\pi_1 X$ is a finitely generated abelian group, there is an isomorphism $\pi_1(\hat{X}^\ell) \cong \mathbb{Z}_\ell \otimes_{\mathbb{Z}} \pi_1 X$, where \mathbb{Z}_ℓ denotes the ℓ -adic integers.

(2) If X is in addition simply connected, and its higher homotopy groups are finitely generated, then we have for all $n \geq 2$

$$\pi_n \hat{X}^\ell \cong \pi_n \hat{X} \cong \mathbb{Z}_\ell \otimes_{\mathbb{Z}} \pi_n X.$$

2.1.1. The model structure on \hat{S} is fibrantly generated

The main technical result of this subsection is that the model structure on \hat{S} , hence also on \hat{S}_* , is fibrantly generated. We recall some notations and constructions from [27]. Let $n \geq 0$ be a non-negative integer and S be a profinite set. The functor $\hat{S}^{\text{op}} \rightarrow \mathcal{E}, X \mapsto \text{Hom}_{\hat{S}}(X_n, S)$ is representable by a simplicial profinite set, which is denoted by $L(S, n)$. It is given by the formula

$$L(S, n) : \Delta^{\text{op}} \rightarrow \hat{\mathcal{E}}, [k] \mapsto S^{\text{Hom}_{\Delta}([n], [k])}.$$

If M is a profinite abelian group, then $L(M, n)$ has a natural structure of a simplicial profinite abelian group and the abelian group $\text{Hom}_{\hat{S}}(X, L(M, n))$ can be identified with the group $C^n(X; M)$ of continuous n -cochains with values in M . Furthermore, for every $k, L(M, *)([k])$ may be considered in the usual way as an abelian cochain complex.

For a profinite abelian group M , let $Z^n(X; M)$ denote the abelian group of n -cocycles of the complex $C^*(X; M)$. The functor $\hat{S}^{\text{op}} \rightarrow \mathcal{E}, X \mapsto Z^n(X; M)$ is also representable by a simplicial profinite abelian group, which is denoted by $K(M, n)$, called the profinite Eilenberg–Mac Lane space of type (M, n) . The homomorphism $C^n(X; M) \rightarrow Z^{n+1}(X; M)$ given by the differential defines a natural map of simplicial profinite abelian groups $L(M, n) \rightarrow K(M, n + 1)$. Consider the two sets of morphisms of Lemme 2 in [27]:

$$P := \{L(M, n) \rightarrow K(M, n + 1), K(M, n) \rightarrow * \mid M \text{ is an abelian pro-}\ell\text{-group, } n \geq 0\}$$

and

$$Q := \{L(M, n) \rightarrow * \mid M \text{ is an abelian pro-}\ell\text{-group, } n \geq 0\}.$$

Theorem 4. *The simplicial model structure on \hat{S} , in which the weak equivalences are the \mathbb{Z}/ℓ -cohomological isomorphisms and the cofibrations are the dimensionwise monomorphisms, is left proper and fibrantly generated with P as the set of generating fibrations and Q as the set of generating trivial fibrations.*

Proof. The left properness is due to the fact that all objects in \hat{S} are cofibrant, see for example [15, Corollary 13.1.3]. The assertion of the theorem now follows from the methods of the proofs of Lemme 2 and Proposition 3 in [27]. One can check for example the list of dual criteria for a cofibrantly generated model structure of Kan’s Theorem 11.3.1 in [15], where we only need the existence of small limits since we use cosmallness. One can find a detailed proof in [31]. \square

2.2. Profinite spectra

2.2.1. The stable model structure

The well-known stabilization of the category of simplicial spectra of [4] uses the fact that S is proper in an essential way in order to construct functorial factorizations. Since the category

$\hat{\mathcal{S}}$ is only left but not right proper, we have to find a different method. Hovey [16] has pointed out a general way to stabilize a left proper cellular model category with respect to a left Quillen endofunctor T .

We have to start with a result on the localization of model categories. We recall the definition of local objects and local equivalences of [15].

Definition 5. Let \mathcal{C} be a simplicial model category. Let K be a set of objects in \mathcal{C} .

- (1) A map $f : A \rightarrow B$ is called a *K-local equivalence* if for every element X of K the induced map of simplicial mapping spaces $f^* : \text{Map}(B, X) \rightarrow \text{Map}(A, X)$ is a weak equivalence of simplicial sets.
- (2) Let \mathcal{K} denote the class of K -local equivalences. An object X is called *K-local* if it is \mathcal{K} -local, i.e. if X is fibrant and for every K -local equivalence $f : A \rightarrow B$ the induced map $f^* : \text{Map}(B, X) \rightarrow \text{Map}(A, X)$ is a weak equivalence.

Theorem 6. Let \mathcal{C} be a left proper fibrantly generated simplicial model category with all small limits. Let K be a set of fibrant objects in \mathcal{C} and let \mathcal{K} be the class of K -local equivalences.

- (1) The left Bousfield localization of \mathcal{C} with respect to \mathcal{K} exists, i.e. there is a model category structure $L_{\mathcal{K}}\mathcal{C}$ on the underlying category \mathcal{C} in which:
 - (a) the class of weak equivalences of $L_{\mathcal{K}}\mathcal{C}$ equals the class of K -local equivalences of \mathcal{C} ,
 - (b) the class of cofibrations of $L_{\mathcal{K}}\mathcal{C}$ equals the class of cofibrations of \mathcal{C} ,
 - (c) the class of fibrations of $L_{\mathcal{K}}\mathcal{C}$ is the class of maps with the right lifting property with respect to those maps that are both cofibrations and K -local equivalences.
- (2) The fibrant objects of $L_{\mathcal{K}}\mathcal{C}$ are the K -local objects of \mathcal{C} .
- (3) $L_{\mathcal{K}}\mathcal{C}$ is left proper. It is fibrantly generated if every object of \mathcal{C} is cofibrant.
- (4) The simplicial structure of \mathcal{C} gives $L_{\mathcal{K}}\mathcal{C}$ the structure of a simplicial model category.

Proof. The proof is a dual version of the proof of Theorem 5.1.1 of [15]. But since we do not assume \mathcal{C} to be cellular, we have to use a dual version of Lemma 2.5 of [6], in order to avoid the use of Proposition 5.2.3 of [15]. One may look at [31] for more details. \square

We return to spectra. Let \mathcal{C} be a left proper fibrantly generated simplicial model category with all small limits and finite colimits. Let $T : \mathcal{C} \rightarrow \mathcal{C}$ be a left Quillen endofunctor on \mathcal{C} . Let $U : \mathcal{C} \rightarrow \mathcal{C}$ be its right adjoint.

Definition 7. A *spectrum* X is a sequence $(X_n)_{n \geq 0}$ of objects of \mathcal{C} together with structure maps $\sigma : TX_n \rightarrow X_{n+1}$ for all $n \geq 0$. A *map of spectra* $f : X \rightarrow Y$ is a collection of maps $f_n : X_n \rightarrow Y_n$ commuting with the structure maps. We denote the category of spectra by $\text{Sp}(\mathcal{C}, T)$.

We begin by defining an intermediate strict structure, see [4] or [16]. As usual we have the following functors.

Definition 8. Given $n \geq 0$, the *evaluation functor* $\text{Ev}_n : \text{Sp}(\mathcal{C}, T) \rightarrow \mathcal{C}$ takes X to X_n . It has a left adjoint $F_n : \mathcal{C} \rightarrow \text{Sp}(\mathcal{C}, T)$ defined by $(F_n A)_m = T^{m-n} A$ if $m \geq n$ and $(F_n A)_m = *$ otherwise. The structure maps are the obvious ones.

The evaluation functor also has a right adjoint $R_n : \mathcal{C} \rightarrow \text{Sp}(\mathcal{C}, T)$ defined by $(R_n A)_i = U^{n-i} A$ if $i \leq n$ and $(R_n A)_i = *$ otherwise. The structure map $TU^{n-i} A \rightarrow U^{n-i} A$ is adjoint to the identity map of U^{n-i} when $i < n$.

Definition 9. A map f in $\text{Sp}(\mathcal{C}, T)$ is a *projective weak equivalence* (respectively *projective fibration*) if each map f_n is a weak equivalence (respectively fibration). A map i is a *projective cofibration* if it has the left lifting property with respect to all projective trivial fibrations.

The following proposition is proved in the standard way [16, Proposition 1.14].

Proposition 10. A map $i : A \rightarrow B$ in $\text{Sp}(\mathcal{C}, T)$ is a projective (trivial) cofibration if and only if $i_0 : A_0 \rightarrow B_0$ and the induced maps $j_n : A_n \sqcup_{TA_{n-1}} TB_{n-1}$ for $n \geq 1$ are (trivial) cofibrations in \mathcal{C} .

We will show that the projective structure is in fact a fibrantly generated model structure. We denote the set of generating fibrations of \mathcal{C} by P and the set of generating trivial fibrations by Q . Inspired by Proposition 10, we set

$$\tilde{P} := \{g_n : R_n R \rightarrow R_n S \times_{R_{n-1} U S} R_{n-1} U R, \text{ for all } f : R \rightarrow S \text{ in } P, n \geq 0\}$$

and

$$\tilde{Q} := \{g_n : R_n R \rightarrow R_n S \times_{R_{n-1} U S} R_{n-1} U R, \text{ for all } f : R \rightarrow S \text{ in } Q, n \geq 0\}$$

where g_n is the map induced by the commutative diagram

$$\begin{array}{ccc} R_n R & \longrightarrow & R_n S \\ \downarrow & & \downarrow \\ R_{n-1} U R & \longrightarrow & R_{n-1} U S. \end{array} \tag{1}$$

Lemma 11. If $f : X \rightarrow Y$ is a fibration (respectively trivial fibration) in \mathcal{C} , then the maps $g_n : R_n X \rightarrow R_n Y \times_{R_{n-1} U Y} R_{n-1} U X$ are fibrations (respectively trivial fibrations) in $\text{Sp}(\mathcal{C}, T)$.

Proof. Since R_n is defined via the right Quillen functor U which preserves fibrations and trivial fibrations, it is clear that the map $(R_n f)_i = U^{n-i}(f)$ on the i th level is a fibration (respectively trivial fibration). In diagram (1), the maps on the i th level are either identities or equal to $U^{n-i}(f)$. Hence the induced map g_n is also a fibration (respectively trivial fibration). \square

Theorem 12. Let \mathcal{C} be a left proper fibrantly generated simplicial model category with all small limits. The projective weak equivalences, projective fibrations and projective cofibrations define a left proper fibrantly generated simplicial model structure on $\text{Sp}(\mathcal{C}, T)$ with set of generating fibrations \tilde{P} and set of generating trivial fibrations \tilde{Q} .

Proof. This intermediate result can be proven in essentially the same way as Theorem 1.14 in [16]. The hard point is the factorization axiom which follows from a cosmall object argument and the fact that \mathcal{C} is fibrantly generated. Since we use a cosmall instead of small object argument the existence of small limits suffices. \square

It remains to modify this structure in order to get a stable structure, i.e. one in which the prolongation of T is a Quillen equivalence. We will do this by applying the localization theorem, Theorem 6, to the projective model structure on spectra. We want the stable weak equivalences to be the maps that induce isomorphisms on all generalized cohomology theories. A generalized cohomology theory is represented by the analogue of an Ω -spectrum.

Definition 13. A spectrum $E \in \text{Sp}(\mathcal{C}, T)$ is defined to be a U -spectrum if each E_n is fibrant and the adjoint structure maps $E_n \rightarrow U E_{n+1}$ are weak equivalences for all $n \geq 0$.

By Corollary 9.7.5 in [15], since each E_n is fibrant and since the right Quillen functor U preserves fibrations, we know that $E_n \rightarrow U E_{n+1}$ is a weak equivalence in \mathcal{C} if and only if the induced map $\text{Map}(A, E_n) \rightarrow \text{Map}(A, U E_{n+1})$ of mapping spaces is a weak equivalence in \mathcal{S} for every cofibrant object A . By adjunction this is equivalent to $\text{Map}(F_n A, E) \rightarrow \text{Map}(F_{n+1} T A, E)$ being a weak equivalence in \mathcal{S} for every cofibrant object A . So we have to define the set K to consist of all U -spectra such that the maps $F_{n+1} T A \rightarrow F_n A$ adjoint to the identity map of $T A$ are the K -local equivalences. By the fact that the projective model structure on $\text{Sp}(\mathcal{C}, T)$ is fibrantly generated, left proper and simplicial, we may deduce the following result from Theorem 6.

Theorem 14. Let \mathcal{C} be a left proper fibrantly generated simplicial model category with all small limits. Let K be the set of U -spectra. There is a stable model structure on $\text{Sp}(\mathcal{C}, T)$ which is defined to be the K -localized model structure $L_{\mathcal{K}} \text{Sp}(\mathcal{C}, T)$ of Theorem 6 where \mathcal{K} is the class of all K -local equivalences.

We apply this to the category $\mathcal{C} = \hat{\mathcal{S}}_*$ and $T = S^1 \wedge \cdot$. By Theorem 4 its model structure is simplicial, left proper and fibrantly generated. Note that $S^1 \wedge \cdot$ is a left Quillen endofunctor since it takes monomorphisms to monomorphisms and preserves cohomological equivalences by the suspension axiom. Its right adjoint is $\hat{\Omega}$.

Definition 15. A profinite spectrum X consists of a sequence $X_n \in \hat{\mathcal{S}}_*$ of pointed profinite spaces for $n \geq 0$ and maps $\sigma_n : S^1 \wedge X_n \rightarrow X_{n+1}$ in $\hat{\mathcal{S}}_*$.

A morphism $f : X \rightarrow Y$ of spectra consists of maps $f_n : X_n \rightarrow Y_n$ in $\hat{\mathcal{S}}_*$ for $n \geq 0$ such that $\sigma_n(1 \wedge f_n) = f_{n+1}\sigma_n$.

We denote by $\text{Sp}(\hat{\mathcal{S}}_*)$ the corresponding category and call it the category of profinite spectra.

Corollary 16. There is a stable model structure on $\text{Sp}(\hat{\mathcal{S}}_*)$ for which the prolongation $S^1 \wedge \cdot : \text{Sp}(\hat{\mathcal{S}}_*) \rightarrow \text{Sp}(\hat{\mathcal{S}}_*)$ is a Quillen equivalence.

In particular, the stable equivalences are the maps that induce an isomorphism on all generalized cohomology theories, represented by profinite $\hat{\Omega}$ -spectra; the stable cofibrations are the maps $i : A \rightarrow B$ such that i_0 and the induced maps $j_n : A_n \amalg_{S^1 \wedge A_{n-1}} S^1 \wedge B_{n-1} \rightarrow B_n$ are monomorphisms for all n ; the stable fibrations are the maps with the right lifting property with respect to all maps that are both stable equivalences and stable cofibrations.

Remark 17. There is also a stable model structure on symmetric profinite spectra in the sense of [17], see [31].

2.2.2. Profinite completion of spectra

Let $\text{Sp}(\mathcal{S}_*)$ be the stable model structure of simplicial spectra defined in [4]. Let $(\hat{\cdot}) : \text{Sp}(\mathcal{S}_*) \rightarrow \text{Sp}(\hat{\mathcal{S}}_*)$ be the profinite completion applied levelwise that takes the spectrum X to the profinite spectrum \hat{X} whose structure maps are given by

$$S^1 \wedge \hat{X}_n \cong S^1 \wedge X_n \xrightarrow{\hat{\sigma}} \hat{X}_{n+1}.$$

Let $|\cdot| : \text{Sp}(\hat{\mathcal{S}}_*) \rightarrow \text{Sp}(\mathcal{S}_*)$ be the levelwise applied forgetful functor. Since the two functors $(\hat{\cdot})$ and $|\cdot|$ on $\hat{\mathcal{S}}_*$ commute with smash products, they form an adjoint pair of functors. The following result may be easily deduced from [27, §2, Proposition 1].

Proposition 18. *The functor $(\hat{\cdot}) : \text{Sp}(\mathcal{S}_*) \rightarrow \text{Sp}(\hat{\mathcal{S}}_*)$ preserves stable equivalences and cofibrations.*

The functor $|\cdot| : \text{Sp}(\hat{\mathcal{S}}_) \rightarrow \text{Sp}(\mathcal{S}_*)$ preserves fibrations and stable equivalences between fibrant objects.*

In particular, $(\hat{\cdot})$ induces a functor on the homotopy categories and the pair $((\hat{\cdot}), |\cdot|)$ is a Quillen pair of adjoint functors.

Similar versions of the following facts have also been proved by Dehon [8]. We deduce the following result from Proposition 3.

Proposition 19. *Let E be a (-1) -connected spectrum and suppose that the \mathbb{Z}/ℓ -cohomology of each E_n , $n \geq 1$, is finite-dimensional in each degree. Then the stable homotopy groups of the profinite completion \hat{E} are given by the following isomorphism for all n :*

$$\pi_n \hat{E} \cong \pi_n \hat{E}^\ell \cong \mathbb{Z}_\ell \otimes_{\mathbb{Z}} \pi_n E.$$

Example 20. 1. Let MU be the simplicial spectrum representing complex cobordism and let MU/ℓ^v be the cofibre spectrum defined by multiplication with ℓ^v . For the profinite completions there are isomorphisms

$$\pi_{2*} \hat{M}U \cong \mathbb{Z}_\ell \otimes_{\mathbb{Z}} \pi_{2*} MU \quad \text{and} \quad \pi_{2*} \hat{M}U/\ell^v \cong \mathbb{Z}/\ell^v \otimes \pi_{2*} MU,$$

where $\pi_{2*} \hat{M}U$ is a polynomial ring in infinitely many variables, cf. [1].

2. Let $\hat{K}U$ be the profinite completion of the Ω -spectrum representing complex K-theory: $\hat{K}U_{2i} = \hat{B}U \times \mathbb{Z}$ and $\hat{K}U_{2i+1} = \hat{U}$ for all $i \geq 0$. Although KU does not satisfy the hypothesis of the proposition, we get by Proposition 3 $\pi_{2i}(\hat{K}U) \cong \mathbb{Z}_\ell$ and $\pi_{2i+1}(\hat{K}U) = 0$ for all i .

2.2.3. Generalized cohomology theories on profinite spaces

It is the main feature of the stable homotopy category of profinite spectra that it provides a general setting for cohomology theories on profinite spaces. We define generalized cohomology theories on $\hat{\mathcal{S}}$ to be the functors represented by profinite spectra. Profinutely completed cohomology theories have already been studied by Dehon in [8].

Let E, X be spectra in $\text{Sp}(\hat{\mathcal{S}}_*)$, we set $E^n(X) := \text{Hom}_{\hat{\mathcal{S}}\mathcal{H}}(X, E[n])$, where $\text{Hom}_{\hat{\mathcal{S}}\mathcal{H}}(X, E[n])$ denotes the set of maps that lower the dimension by n , and call this the n th cohomology group

of X with values in E . We set $E^*(X) := \bigoplus_n E^n(X)$. For a pointed profinite space X we define its n th cohomology groups with values in E by

$$E^n(X) := \text{Hom}_{\mathcal{SH}}(\Sigma^\infty(X), E[n]). \tag{2}$$

For a pair (X, A) of profinite spaces we define the relative cohomology by $E^n(X, A) := E^n(X/A)$.

2.2.4. *The Atiyah–Hirzebruch spectral sequence*

Let X be a profinite space. For every integer p , we denote by $\text{sk}_p X$ the profinite subspace of X which is generated by the simplices of degree less or equal p . We call $\text{sk}_p X$ the p -skeleton of X . For every k the set of k -simplices of $\text{sk}_p X$ is closed in X_k such that $\text{sk}_p X$ is a simplicial profinite subset of X , cf. [8]. Let E be a profinite spectrum. We consider the skeletal filtration $\text{sk}_0 X \subset \dots \subset \text{sk}_p X \subset \text{sk}_{p+1} X \subset \dots \subset X$. It yields a filtration on $E^* X$ defined by $F^p E^* X := \text{Ker}(E^* X \rightarrow E^* \text{sk}_{p-1} X)$.

Since the coefficient groups E^q have a natural profinite structure, we may consider continuous cohomology with coefficients in E^q . The proof of the following theorem is essentially the one given by Adams in [1]. Dehon has translated it to the profinite setting in [8, Proposition 2.1.9]. Although Dehon proves the following assertion only for special cohomology theories, the proof works in the category of profinite spectra for any profinite cohomology theory, see [31].

Proposition 21. *For any profinite spectrum E and for any profinite space X there is a spectral sequence $\{E_r^{p,q}\}$ with $E_2^{p,q} \cong H^p(X; E^q) \Rightarrow E^{p+q}(X)$. The spectral sequence converges strongly, in the sense of [1, III, §8.2], to the graded term $E_\infty^{p,q} = F^p E^* X / F^{p+1} E^* X$ of the filtration on $E^* X$ if $\lim_r^1 E_r^{p,q} = 0$. In particular, the spectral sequence converges if $H^p(X; E^q) = 0$ for p sufficiently large or if $H^p(X; E^q)$ is finite for all p . We call it the profinite Atiyah–Hirzebruch spectral sequence.*

2.2.5. *Comparison with generalized cohomology theories of pro-spectra*

As a corollary, we compare the cohomology of profinite spectra with cohomology theories of pro-spectra. Isaksen constructs in [19] a stable model structure on the category of pro-spectra. If E is a pro-spectrum then the r th cohomology $E_{\text{pro}}^r(X)$ of a pro-spectrum X with coefficients in E is the set $[X, E]_{\text{pro}}^{-r}$ of maps that lower the degree by r in the stable homotopy category of pro-spectra. In addition, for two pro-spectra X and E he shows the existence of an Atiyah–Hirzebruch spectral sequence $E_2^{p,q} = H^{-p}(X; \pi_{-q} E) \Rightarrow [X, E]_{\text{pro}}^{p+q}$ which is in particular convergent if E is a constant pro-spectrum and X is the suspension spectrum of a finite-dimensional pro-space. Taking completion and homotopy limits defines a functor $(\hat{\cdot}) : \text{pro-Sp}(\mathcal{S}_*) \rightarrow \text{Sp}(\hat{\mathcal{S}}_*)$. One can check that it also induces maps $E_{\text{pro}}^*(X) \rightarrow \hat{E}^*(\hat{X})$, which yield morphisms of Atiyah–Hirzebruch spectral sequences. Since the E_2 -terms agree, we get

Corollary 22. *Let $\{X_s\}_{s \in I}$ be a finite-dimensional pro-space. Then there is an isomorphism*

$$MU_{\text{pro}}^* (\{X_s\}; \mathbb{Z}/\ell^v) \cong \hat{M}U^* (\hat{X}; \mathbb{Z}/\ell^v).$$

3. Stable étale realization on the \mathbb{A}^1 -homotopy category

The construction of the étale topological type functor Et from locally Noetherian schemes to pro-spaces is due to Artin–Mazur and Friedlander. The construction of the \mathbb{A}^1 -homotopy category of schemes gave rise to the question if this functor may be enlarged to the category of spaces. This has been answered independently by Isaksen and Schmidt. The latter one constructed in [32] a geometric functor to the category of pro-objects in \mathcal{H} . Isaksen gave a rigid, but less intuitive construction in [18]. We follow Isaksen’s approach. The first step in this direction was the construction of a model structure on $\text{pro-}\mathcal{S}$.

3.0.1. The functor $\hat{\text{Et}}$

Let $\text{pro-}\mathcal{S}$ be the category whose objects are cofiltered diagrams $X(-) : I \rightarrow \mathcal{S}$ and its morphisms are defined by

$$\text{Hom}_{\text{pro-}\mathcal{S}}(X(-), Y(-)) := \lim_{t \in J} \text{colim}_{s \in I} \text{Hom}_{\mathcal{S}}(X(s), Y(t)).$$

The cohomology with \mathbb{Z}/ℓ -coefficients is defined to be

$$H^*(X(-), \mathbb{Z}/\ell) := \text{colim}_s H^*(X(s), \mathbb{Z}/\ell).$$

Isaksen has constructed several model structures on $\text{pro-}\mathcal{S}$; in particular, the \mathbb{Z}/ℓ -cohomological model structure of [20] in which the weak equivalences are morphisms inducing isomorphisms in \mathbb{Z}/ℓ -cohomology and the cofibrations are levelwise monomorphisms.

We define a completion functor $(\hat{\cdot}) : \text{pro-}\mathcal{S} \rightarrow \hat{\mathcal{S}}$ as the composite of two functors. First we apply $(\hat{\cdot}) : \mathcal{S} \rightarrow \hat{\mathcal{S}}$ levelwise, then we take the limit in $\hat{\mathcal{S}}$ of the underlying diagram. The next lemma shows that the functor $(\hat{\cdot})$ respects the model structures on $\text{pro-}\mathcal{S}$ and $\hat{\mathcal{S}}$. Its proof is clear using Lemme 1.1.1 of [8]. Using the functor $\hat{\mathcal{S}} \xrightarrow{\mathcal{G}} \text{pro-}\mathcal{S} \xrightarrow{\mathcal{I}} \text{pro-}\mathcal{S}$ of Isaksen’s it follows as in [20] from this lemma that the homotopy categories $\text{Ho}(\text{pro-}\mathcal{S})$ and $\hat{\mathcal{H}}$ are in fact equivalent via completion.

Lemma 23.

(1) *Let $X \in \text{pro-}\mathcal{S}$ be a pro-simplicial set. Then we have a natural isomorphism of cohomology groups*

$$H^*(X; \mathbb{Z}/\ell) \xrightarrow{\cong} H^*(\hat{X}; \mathbb{Z}/\ell).$$

(2) *A morphism $f : X \rightarrow Y$ of pro-simplicial sets induces an isomorphism in \mathbb{Z}/ℓ -cohomology if and only if the morphism $\hat{f} : \hat{X} \rightarrow \hat{Y}$ in $\hat{\mathcal{S}}$ induces an isomorphism in continuous \mathbb{Z}/ℓ -cohomology.*

(3) *The functor $(\hat{\cdot}) : \text{pro-}\mathcal{S} \rightarrow \hat{\mathcal{S}}$ preserves monomorphisms.*

Now we turn to the applications in algebraic geometry. We refer the reader to [14,18] for a detailed discussion of the category of rigid hypercoverings and rigid pullbacks. The following definition is taken from [18].

Definition 24. Let X be a locally Noetherian scheme. The *étale topological type* of X is defined to be the pro-simplicial set

$$\text{Et } X := \text{Re} \circ \pi : \text{HRR}(X) \rightarrow \mathcal{S}$$

sending a rigid hypercovering U of X to the simplicial set of connected components of U . If $f : X \rightarrow Y$ is a map of locally Noetherian schemes, then the strict map $\text{Et } f : \text{Et } X \rightarrow \text{Et } Y$ is given by the functor $f^* : \text{HRR}(Y) \rightarrow \text{HRR}(X)$ and the natural transformation $\text{Et } X \circ \text{Et } f \rightarrow \text{Et } Y$.

Isaksen uses the insight of Dugger [9] that one can construct the unstable \mathbb{A}^1 -homotopy category in a universal way. Starting from an almost arbitrary category \mathcal{C} , Dugger constructs an enlargement of \mathcal{C} that carries a model structure and is universal for this property. Isaksen extends in [18] the functor Et via this general method to the \mathbb{A}^1 -homotopy category. The idea is that $\text{Et } X$ should be the above $\text{Et } X$ on a representable presheaf X and should preserve colimits and the simplicial structure. For our purpose, we would like to define $\hat{\text{Et}}$ directly in the way Dugger suggests. But the problem is that $(\hat{\cdot}) : \text{pro-}\mathcal{S} \rightarrow \hat{\mathcal{S}}$ is not a left adjoint functor and does not preserve all small colimits. Therefore, we define $\hat{\text{Et}}$ to be the composition of Et followed by completion.

Let Sm/k be the category of smooth quasi-projective schemes of finite type over k and $\Delta^{\text{op}} \text{PreShv}(\text{Sm}/k)$ the category of simplicial presheaves on Sm/k . We make the following definition whose first part is due to Isaksen [18].

Definition 25. If X is a representable presheaf, then $\text{Et } X$ is the étale topological type of X . If P is a discrete presheaf, i.e. P is just a presheaf of sets, then P can be written as a colimit $\text{colim}_i X_i$ of representables and we define $\text{Et } P := \text{colim}_i \text{Et } X_i$. Finally, an arbitrary simplicial presheaf can be written as the coequalizer of the diagram

$$\coprod_{[m] \rightarrow [n]} P_m \otimes \Delta^n \rightrightarrows \coprod_{[n]} P_n \otimes \Delta^n,$$

where each P_n is discrete. Define $\text{Et } P$ to be the coequalizer of the diagram

$$\coprod_{[m] \rightarrow [n]} \text{Et } P_m \otimes \Delta^n \rightrightarrows \coprod_{[n]} \text{Et } P_n \otimes \Delta^n.$$

We define the profinite étale topological type functor $\hat{\text{Et}}$ to be the composition of Et and the profinite completion functor $\text{pro-}\mathcal{S} \rightarrow \hat{\mathcal{S}}$:

$$\hat{\text{Et}} := (\hat{\cdot}) \circ \text{Et} : \Delta^{\text{op}} \text{PreShv}(\text{Sm}/k) \rightarrow \hat{\mathcal{S}}.$$

For computations the following remark is crucial.

Remark 26. Let M be a locally constant sheaf on the locally Noetherian scheme X . By [14, Proposition 5.9], we know that there is an isomorphism for étale cohomology $H_{\text{ét}}^*(X; M) \cong H^*(\text{Et } X, M)$ where $H^*(\text{Et } X; M)$ denotes the cohomology of a pro-simplicial set $\text{Et } X$ with coefficients in the local coefficient system M corresponding to the sheaf M . For a finite abelian

group π , we have in addition a natural isomorphism by Lemma 23: $H^*(Z; \pi) \cong H^*(\hat{Z}; \pi)$ for every pro-simplicial set Z . Hence we get as well

$$H_{\text{ét}}^*(X; \pi) \cong H^*(\hat{\text{Ét}} X, \pi)$$

for every locally Noetherian scheme X and every finite abelian group π .

Furthermore, the pro-fundamental group $\pi_1(\text{Ét } X)$ is isomorphic as a profinite group to the étale fundamental group $\pi_1^{\text{ét}}(X)$ of X . Hence $\pi_1(\hat{\text{Ét}} X)$ is equal to $\pi_1^{\text{ét}}(X)^{\wedge \ell}$ by Proposition 3.

Examples. 1. Let R be a strict local Henselian ring, i.e. a local Henselian ring with separably closed residue field. Then $\text{Spec } R$ has no nontrivial étale covers and the étale topological type of $\text{Spec } R$ is a contractible space.

2. Let k be a separably closed field with $\text{char}(k) \neq \ell$. The space \mathbb{G}_m is connected and its ℓ -completed étale fundamental group is \mathbb{Z}_ℓ , the ℓ -adic integers. This implies $\hat{\text{Ét}} \mathbb{G}_m \cong K(\mathbb{Z}_\ell, 1)$ in $\hat{\mathcal{S}}$, where $K(\mathbb{Z}_\ell, 1)$ is naturally a profinite space since \mathbb{Z}_ℓ is a profinite group.

3. Let k be a separably closed field with $\text{char}(k) \neq \ell$. Let \mathbb{P}_k^1 be the projective line over k . Since \mathbb{P}_k^1 is connected and its étale fundamental group $\pi_1^{\text{ét}}(\mathbb{P}_k^1) = 0$ is trivial, see e.g. [26, I, Example 5.2 f)], $\hat{\text{Ét}} \mathbb{P}_k^1$ is a simply connected space. Apart from H^0 its only nonzero étale cohomology group is $H_{\text{ét}}^2(\mathbb{P}_k^1, \mathbb{Z}/\ell) \cong \mathbb{Z}/\ell$, with a chosen isomorphism $\mathbb{Z}/\ell \rightarrow \mathbb{Z}/\ell(1)$. Hence $\hat{\text{Ét}} \mathbb{P}_k^1$ is isomorphic in $\hat{\mathcal{S}}$ to the simplicial finite set S^2 .

4. Let $k = \mathbb{F}_q$ be a finite field with $\text{char}(k) = p \neq \ell$. The étale topological type of k is isomorphic to S^1 in $\hat{\mathcal{H}}$. For, it is connected and its ℓ -completed fundamental group is the ℓ -completion of the absolute Galois group of k , i.e. $\pi_{1, \text{ét}}^\ell(k) = \mathbb{Z}_\ell$. Since the cohomology groups $H^i(k; \mathbb{Z}/\ell)$ vanish for $i > 1$, $\hat{\text{Ét}} k$ is a space of dimension one and weakly equivalent to S^1 .

3.1. Profinite étale realization of motivic spaces

We consider as in [9] the category $U(\text{Sm}/k) := \Delta^{\text{op}} \text{PreShv}(\text{Sm}/k)$ of simplicial presheaves on Sm/k with the projective model structure: the weak equivalences (fibrations) are objectwise weak equivalences (fibrations) of simplicial sets, the cofibrations are the maps having the left lifting property with respect to all trivial fibrations. Then one takes the left Bousfield localization of this model structure at the set S of maps:

- (1) for every finite collection $\{X^a\}$ of schemes with disjoint union X , the map $\coprod X^a \rightarrow X$ from the coproduct of the presheaves represented by X^a to the presheaf represented by X ;
- (2) every Nisnevich (étale) hypercover $U. \rightarrow X$;
- (3) $X \times \mathbb{A}^1 \rightarrow X$ for every scheme X .

We call this the *Nisnevich (étale) \mathbb{A}^1 -local projective model structure* according to [18], and denote it by $LU(k)$ (respectively $L_{\text{ét}}U(k)$).

Proposition 8.1 of [9] states that $LU(k)$ is Quillen equivalent to the Nisnevich \mathbb{A}^1 -localized model category $\mathcal{M}\mathcal{V}_k$ of [29] and the analogue holds for the étale case. For an étale realization over a field of positive characteristic, we have to consider the following point. Since the étale fundamental group of the affine line \mathbb{A}_k^1 over a field k of positive characteristic is non-trivial, one has to complete away from the characteristic of k in this case, using the fact that the projection $X \times \mathbb{A}^1 \rightarrow X$ induces an isomorphism in étale cohomology $H_{\text{ét}}^*(X; \mathbb{Z}/\ell) \rightarrow H_{\text{ét}}^*(X \times \mathbb{A}^1; \mathbb{Z}/\ell)$

for every prime $\ell \neq p$. This means that we have to consider the \mathbb{Z}/ℓ -cohomological model structure on $\text{pro-}\mathcal{S}$, respectively $\hat{\mathcal{S}}$. The following theorem is due to Isaksen [18].

Theorem 27. *Let ℓ be a prime different from the characteristic of k . With respect to the Nisnevich (étale) \mathbb{A}^1 -local projective model structure on simplicial presheaves on Sm/k , the functor $\hat{\text{Et}}$ induces a functor $\text{L}\hat{\text{Et}}$ from the Nisnevich (étale) \mathbb{A}^1 -homotopy category to the \mathbb{Z}/ℓ -cohomological homotopy category of $\hat{\mathcal{S}}$. In particular, $\text{L}\hat{\text{Et}} X$ is just $\hat{\text{Et}} X$ for every scheme in Sm/k , and hence $\hat{\text{Et}}$ preserves \mathbb{A}^1 -weak equivalences between smooth schemes over k .*

Proof. Since every Nisnevich hypercover is also an étale hypercover it suffices to check this for the étale case. The first part of the proof is the one of [18, Theorem 6].

By Lemma 23 completion preserves weak equivalences and cofibrations. Hence the composition $\hat{\text{Et}}$ sends weak equivalences between cofibrant objects into weak equivalences and the total left derived functor $\text{L}\hat{\text{Et}}: \mathcal{H}_{\mathbb{A}^1}^{\text{ét}}(k) \rightarrow \hat{\mathcal{H}}$ from the étale \mathbb{A}^1 -homotopy category $\mathcal{H}_{\mathbb{A}^1}^{\text{ét}}(k)$ of schemes over k of [29] exists. \square

3.2. Étale realization of motivic spectra

We extend the results of the previous section to the stable \mathbb{A}^1 -homotopy category. This étale realization of the stable motivic category is the technical key point for the construction of a transformation from algebraic cobordism given by the *MGL*-spectrum to the profinite étale cobordism of the next section. One of the reasons why we consider the model $\hat{\mathcal{S}}$ of $\text{Ho}(\hat{\mathcal{S}})$, instead of $\text{pro-}\mathcal{S}$, is that it seems to be easier to extend the functor $\hat{\text{Et}}$ to the category of profinite spectra rather than pro-spectra . I am especially grateful to Fabien Morel for helpful discussions on the problems in this section.

Let k be the base field of characteristic different from ℓ and let \bar{k} be its separable closure. For the category $\text{Sp}^{\mathbb{P}^1}(k)$ of motivic \mathbb{P}^1_k -spectra over k , in the sense of [28] for example, we have to consider presheaves \mathcal{X} pointed by a morphism $\text{Spec } k \rightarrow \mathcal{X}$. This forces us to consider also the category $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$ of pointed profinite spaces over $\hat{\text{Et}}k$. Its objects (X, p, s) are pointed profinite spaces X together with a projection morphism $p: X \rightarrow \hat{\text{Et}}k$ and a section morphism $s: \hat{\text{Et}}k \rightarrow X$. The morphisms in this category are commutative diagrams in the obvious sense. The étale realization of a pointed presheaf \mathcal{X} is naturally an object of $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$ via the images of the projection and section morphisms of \mathcal{X} . For the contractible space $\hat{\text{Et}}\bar{k}$, we choose and fix a base point. Since all étale covers of $\text{Spec } \bar{k}$ are trivial, any choice is ok. The point is that this allows us to consider every $\hat{\text{Et}} \mathcal{X}$ as a pointed space compatible with base extensions of k in the following way. The space $\hat{\text{Et}} \mathcal{X}$ is pointed by the composite $* \rightarrow \hat{\text{Et}}\bar{k} \rightarrow \hat{\text{Et}}k \rightarrow \mathcal{X}$. Furthermore, via the canonical maps $X \rightarrow * \rightarrow \hat{\text{Et}}\bar{k} \rightarrow \hat{\text{Et}}k$ and $\hat{\text{Et}}k \rightarrow * \rightarrow X$ we may view every space X in $\hat{\mathcal{S}}_*$ as an object in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$. In particular, the 2-sphere S^2 is naturally an object in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$. We consider the usual model structure on $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$ where weak equivalences (respectively cofibrations, fibrations) are those maps which are \mathbb{Z}/ℓ -weak equivalences (respectively cofibrations, fibrations) in $\hat{\mathcal{S}}$ after forgetting the projection and section maps. This model structure on $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$ is left proper and fibrantly generated and we construct a stable model structure on the category $\text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k, S^2 \wedge \cdot)$ of profinite S^2 -spectra over $\hat{\text{Et}}k$ exactly in the same way as for $\text{Sp}(\hat{\mathcal{S}}_*, S^2 \wedge \cdot)$. Its homotopy category will be denoted by $\hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Et}}k$.

3.2.1. An intermediate category

The problem for the stable motivic version is that Et and hence also $\hat{\text{Et}}$ do not commute with products in general. We have to construct an intermediate category and show by a zig-zag of

functors that we get a functor on the homotopy level. However, the projections to each factor induce a canonical map

$$\hat{\text{Et}}(\mathbb{P}_k^1 \wedge_k X) \rightarrow \hat{\text{Et}}(\mathbb{P}_k^1) \wedge_{\hat{\text{Et}}k} \hat{\text{Et}}(X).$$

Lemma 28. *For every pointed presheaf \mathcal{X} on Sm/k , the sequence of canonical maps in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$*

$$S^2 \wedge \hat{\text{Et}} \mathcal{X} \xrightarrow{\cong} \hat{\text{Et}}(\mathbb{P}_k^1) \wedge_{\hat{\text{Et}}k} \hat{\text{Et}}(\mathcal{X}) \xleftarrow{\cong} \hat{\text{Et}}(\mathbb{P}_k^1 \wedge_k \mathcal{X})$$

is a sequence of \mathbb{Z}/ℓ -weak equivalences.

Proof. For $X \in \text{Sm}/k$, the assertion may easily be deduced from the projective bundle formula for étale cohomology and the isomorphism $\hat{\text{Et}}\mathbb{P}_k^1 \cong S^2$ in $\hat{\mathcal{S}}_*$.

If \mathcal{X} denotes a presheaf on Sm/k , \mathcal{X} is isomorphic to the colimit of representable presheaves $\mathcal{X} = \text{colim}_s X_s$. Since each X_s is a smooth scheme over k , the étale cohomology groups $H^i(\text{Et } X_s; \mathbb{Z}/\ell) = H_{\text{ét}}^i(X_s; \mathbb{Z}/\ell)$ are finite \mathbb{Z}/ℓ -vector spaces in each degree. Hence, since Et commutes with colimits, we get $H^*(\text{Et } \mathcal{X}; \mathbb{Z}/\ell) \cong \lim_s H^*(\text{Et } X_s; \mathbb{Z}/\ell)$, as the \lim^1 of these finite groups vanishes, and the limit over all s of $H^*(\text{Et } X_s; \mathbb{Z}/\ell)$ commutes with the functor $H^*(\text{Et } \mathbb{P}_k^1; \mathbb{Z}/\ell) \otimes_{H^*(\text{Et } k; \mathbb{Z}/\ell)} -$. \square

Hence if $\sigma_n: \mathbb{P}_k^1 \wedge_k E_n \rightarrow E_{n+1}$ is the structure map of a motivic \mathbb{P}_k^1 -spectrum, then $\hat{\text{Et}}$ yields a sequence of maps

$$S^2 \wedge \hat{\text{Et}} E_n \xrightarrow{\cong} \hat{\text{Et}}(\mathbb{P}_k^1) \wedge_{\hat{\text{Et}}k} \hat{\text{Et}}(E_n) \xleftarrow{\cong} \hat{\text{Et}}(\mathbb{P}_k^1 \wedge_k E_n) \xrightarrow{\hat{\text{Et}}\sigma_n} E_{n+1} \tag{3}$$

where the first two maps are weak equivalences in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$. Since there is no natural inverse map $\hat{\text{Et}}(\mathbb{P}_k^1) \wedge_{\hat{\text{Et}}k} \hat{\text{Et}}(\mathcal{X}) \rightarrow \hat{\text{Et}}(\mathbb{P}_k^1 \wedge_k \mathcal{X})$ in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$, we may only construct an étale realization on the level of homotopy categories. Therefore, we consider an intermediate category \mathcal{C} and deduce from a zig-zag of functors

$$\text{Sp}^{\mathbb{P}^1}(k) \xrightarrow{\hat{\text{Et}}} \mathcal{C}/\hat{\text{Et}}k \xleftarrow{i} \text{Sp}(\hat{\mathcal{S}}/\hat{\text{Et}}k, S^2 \wedge \cdot)$$

the existence of a stable realization functor $\mathcal{SH}(k) \rightarrow \hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Et}}k$.

In view of Lemma 28, it is natural to consider the following definition. The objects of the category $\mathcal{C}/\hat{\text{Et}}k$ are sequences

$$\{F_n, F'_n, F''_n; S^2 \wedge F_n \xrightarrow{\cong p_n} F'_n \xleftarrow{\cong q_n} F''_n \xrightarrow{r_n} F_{n+1}\}_{n \in \mathbb{N}}$$

where F_n, F'_n, F''_n are pointed profinite spaces over $\hat{\text{Et}}k$ and p_n, q_n and r_n are maps in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$; furthermore the maps p_n and q_n are weak equivalences in $\hat{\mathcal{S}}_*$.

The morphisms of $\mathcal{C}/\hat{\text{Et}}k$ are levelwise morphisms of $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$ which make the obvious diagrams commutative, where the map $S^2 \wedge E_n \rightarrow S^2 \wedge F_n$ is the map induced by $E_n \rightarrow F_n$. The functor $\text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k, S^2 \wedge \cdot) \xrightarrow{i} \mathcal{C}/\hat{\text{Et}}k$ denotes the full embedding which sends $\{F_n, S^2 \wedge F_n \xrightarrow{\sigma_n} F_{n+1}\}$ to $\{F_n, S^2 \wedge F_n, S^2 \wedge F_n; S^2 \wedge F_n \xrightarrow{\text{id}} S^2 \wedge F_n \xleftarrow{\text{id}} S^2 \wedge F_n \xrightarrow{\sigma_n} F_{n+1}\}$.

On the other hand, we get a functor $\text{Sp}^{\mathbb{P}^1}(k) \xrightarrow{\hat{\text{Et}}} \mathcal{C}/\hat{\text{Et}}k$, when we apply $\hat{\text{Et}}$ levelwise.

We define a class W of maps in $\mathcal{C}/\hat{\text{Et}}k$ as the image of the stable equivalences of $\text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k, S^2 \wedge \cdot)$ under the embedding i . Since the maps in W are the images of weak equivalences in a model structure and since i is a full embedding, it follows that W admits a calculus of fractions and we may form the localized category $\text{Ho}(\mathcal{C}/\hat{\text{Et}}k) := \mathcal{C}[W^{-1}]$. We call the maps in W weak equivalences or stable equivalences, by abuse of notation. We will call a map in W a level equivalence if it is in the image of the level equivalences of $\text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k, S^2 \wedge \cdot)$ under i .

Proposition 29. *The induced embedding $i : \hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Et}}k \rightarrow \text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$ is an equivalence of categories.*

The corresponding inverse equivalence is denoted by $j : \text{Ho}(\mathcal{C}/\hat{\text{Et}}k) \rightarrow \hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Et}}k$. Because of the necessary choices in the proof, one should note that j is not quite constructible in practice.

Proof. Since i is a full embedding, it suffices to show that for every $F \in \text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$ there is a spectrum $E \in \hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Et}}k$ such that $i(E) \cong F$ in $\text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$. The crucial point is to construct the structure map of a spectrum from the given data of F .

Let R be a fixed fibrant replacement functor in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$. We consider the category $\text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k, RS^2 \wedge \cdot)$ as a Quillen equivalent model for $\hat{\mathcal{S}}\mathcal{H}_2/\hat{\text{Et}}k$. Since R commutes with products, applying R on each level yields the following sequence:

$$RS^2 \wedge RF_n \xrightarrow{\simeq Rp_n} RF'_n \xleftarrow{\simeq Rq_n} RF''_n \xrightarrow{Rr_n} RF_{n+1}.$$

In addition, the functor R can be chosen such that the map Rq_n is a trivial fibration between fibrant and cofibrant objects, see Proposition 8.1.23 of [15]. The sequence we get is still isomorphic in $\text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$ to the initial one since they are even level equivalent. By Proposition 9.6.4 of [15], there is a right inverse s_n of Rq_n in $\hat{\mathcal{S}}_*/\hat{\text{Et}}k$ such that $Rq_n s_n = \text{id}_{RF'_n}$ and a homotopy $s_n Rq_n \sim \text{id}_{RF''_n}$. We denote by E the resulting spectrum with structure maps $\sigma_n := Rr_n \circ s_n \circ Rp_n$.

Now it is easy to check that $i(E)$ is isomorphic to F in $\text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$. \square

3.2.2. The main result

We want to show that $\tilde{\text{Et}}_{\text{Sp}} : \text{Sp}^{\mathbb{P}^1}(k) \rightarrow \mathcal{C}/\hat{\text{Et}}k$ has a total left derived functor. Therefore, we have to choose a good model for the stable motivic category $\mathcal{SH}(k)$. By [9] we know that $LU(k)$ is a left proper cellular simplicial model category. Hence, in order to be able to apply the methods of [16], we only need an appropriate left Quillen endofunctor. Unfortunately, as the referee pointed out, the map $* \rightarrow \mathbb{P}^1$ is not a projective cofibration in $LU(k)$. One should note that this problem does not appear when we use the injective model structure on presheaves, in which every monomorphism is a cofibration, see Jardine’s work on motivic spectra [22]. But Et is not a Quillen functor on the injective structure since there are too many injective cofibrations, see [18, Remark 3].

Nevertheless, there are at least two possible approaches to solve this problem and we are indebted to the referee for calling our attention to them. The probably more elegant solution is to show that Et is a left Quillen functor on the flasque model structure on $\Delta^{\text{op}} \text{PreShv}(\text{Sm}/k)$ constructed by Isaksen in [21], for which $* \rightarrow \mathbb{P}^1$ is a flasque cofibration. The easier and more obvious way is to factor $* \rightarrow \mathbb{P}^1$ as $* \rightarrow Q \rightarrow \mathbb{P}^1$, where $* \rightarrow Q$ is a projective cofibration and $p : Q \rightarrow \mathbb{P}^1$ is a trivial fibration in $LU(k)$. We follow the latter approach, which is in the spirit of [10, Section 2.2], where the sphere $\mathbb{A}^1/\mathbb{A}^1 - \{0\}$ is replaced by a projective cofibrant motivic space. Up to weak equivalence the choice of the motivic sphere is irrelevant for the

stable motivic homotopy category, see [16, Theorem 5.7]. Since $* \rightarrow Q$ is a cofibration, the functor $Q \wedge \cdot : LU(k) \rightarrow LU(k)$ is a left Quillen functor and we are able to apply the methods of [16, Sections 3 and 5]. Furthermore, via the map $p : Q \rightarrow \mathbb{P}^1$, every \mathbb{P}^1 -spectrum induces a Q -spectrum.

Proposition 30. *The canonical functors*

$$Sp^{\mathbb{P}^1}(k) \rightarrow Sp(\mathcal{MV}_k, \mathbb{P}^1 \wedge \cdot) \rightarrow Sp(LU(k), Q \wedge \cdot)$$

are Quillen equivalences.

Proof. The first equivalence follows as in the proof of [16, Corollary 3.5], taking into account that cofibrations are sent to cofibrations and that the fibrant objects agree in both model structures.

The second equivalence follows from [16, Theorem 5.7], taking into account that $LU(k)$ is Quillen equivalent to \mathcal{MV}_k and that all objects in \mathcal{MV}_k are cofibrant. The functor $\mathcal{MV}_k \rightarrow LU(k)$ is just the full embedding of Nisnevich sheaves into presheaves, its inverse is sheafification, see [9]. \square

Since p is a trivial fibration between cofibrant objects, it is even a simplicial homotopy equivalence. Hence $\hat{E}t p : \hat{E}t Q \rightarrow \hat{E}t \mathbb{P}^1$ is a weak equivalence in \hat{S}_* . The base change $\cdot \wedge_{\text{Spec } k} \text{Spec } \bar{k}$ preserves homotopy equivalences. Hence, by the example above, $\hat{E}t Q_{\bar{k}}$ is a simply connected space whose cohomology is equal to the one of S^2 . We conclude that $\hat{E}t Q_{\bar{k}}$ is even isomorphic to the simplicial finite set S^2 and to $\hat{E}t \mathbb{P}_{\bar{k}}^1$ in \hat{S}_* and the sequence of maps in Lemma 28 yields a sequence of \mathbb{Z}/ℓ -weak equivalences in \hat{S}_* for every simplicial presheaf \mathcal{X} :

$$S^2 \wedge \hat{E}t \mathcal{X} \xrightarrow{\cong} \hat{E}t(Q) \wedge_{\hat{E}t k} \hat{E}t(\mathcal{X}) \xleftarrow{\cong} \hat{E}t(Q \wedge_k \mathcal{X}). \tag{4}$$

Finally, we can prove that there is a stable étale realization functor on the \mathbb{A}^1 -homotopy theory of schemes over an arbitrary base field of characteristic different from ℓ .

Theorem 31. *The functor $\hat{E}t$ induces an étale realization of the stable motivic homotopy category of \mathbb{P}^1 -spectra:*

$$L\hat{E}t : \mathcal{SH}(k) \rightarrow \hat{S}\mathcal{H}_2/\hat{E}t k.$$

Proof. We use $Sp(LU(k), Q \wedge \cdot)$ as a model for $\mathcal{SH}(k)$. The desired functor is defined to be the composite $L\hat{E}t : \mathcal{SH}(k) \xrightarrow{L\hat{E}t} \text{Ho}(\mathcal{C}/\hat{E}t k) \xrightarrow{j} \hat{S}\mathcal{H}_2/\hat{E}t k$. For the existence of $L\hat{E}t$ it remains to show that stable equivalences in $Sp(LU(k), Q \wedge \cdot)$ are sent to isomorphisms in $\text{Ho}(\mathcal{C}/\hat{E}t k)$.

We know that $\hat{E}t_{\text{Sp}}$ sends level equivalences between cofibrant objects in $LU(k)$ to weak equivalences in $\mathcal{C}/\hat{E}t k$, since $\hat{E}t$ sends weak equivalences between cofibrant objects to weak equivalences in $\hat{S}_*/\hat{E}t k$. Hence it induces a total left derived functor on the projective model structure of $Sp(LU(k), Q \wedge \cdot)$.

We use the notation $\Sigma_n^Q : LU(k) \rightarrow Sp(LU(k), Q \wedge \cdot)$ for the left adjoint to the n th evaluation functor. It is given by $(\Sigma_n^Q \mathcal{X})_m = Q^{m-n} \mathcal{X}$ if $m \geq n$ and $(\Sigma_n^Q \mathcal{X})_m = \text{Spec } k$ otherwise, where

$Q^{m-n} \mathcal{X}$ denotes the smash product of \mathcal{X} with $m - n$ copies of Q . We denote by $F_n : \hat{\mathcal{S}}_*/\hat{\text{E}t}k \rightarrow \mathcal{C}/\hat{\text{E}t}k$ the composition of the corresponding functor $\hat{\Sigma}_n : \hat{\mathcal{S}}_*/\hat{\text{E}t}k \rightarrow \text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{E}t}k, S^2 \wedge \cdot)$ followed by the embedding i .

In order to show that there exists a derived functor on the stable structure it suffices to show that $\hat{\text{E}t}_{\text{Sp}}(\zeta_n^{\mathcal{X}})$ is a stable equivalence for maps $\zeta_n^{\mathcal{X}} : \Sigma_{n+1}Q \wedge \mathcal{X} \rightarrow \Sigma_n \mathcal{X}$ in $\text{Sp}(LU(k), Q \wedge \cdot)$ for all cofibrant presheaves $\mathcal{X} \in LU(k)$, since these are the maps at which we localize to get the stable model structure, cf. [15,16].

We consider the commutative diagram in $\text{Ho}(\mathcal{C}/\hat{\text{E}t}k)$:

$$\begin{array}{ccc}
 \zeta_n^{\hat{\text{E}t}\mathcal{X}} : F_{n+1}(S^2 \wedge \hat{\text{E}t}\mathcal{X}) & \xrightarrow{\cong} & F_n(\hat{\text{E}t}\mathcal{X}) \\
 \updownarrow \cong & & \parallel \\
 F_{n+1}(\hat{\text{E}t}Q \wedge_{\hat{\text{E}t}k} \hat{\text{E}t}\mathcal{X}) & & F_n(\hat{\text{E}t}\mathcal{X}) \\
 \updownarrow \cong & & \parallel \\
 F_{n+1}(\hat{\text{E}t}(Q \wedge \mathcal{X})) & & F_n(\hat{\text{E}t}\mathcal{X}) \\
 \updownarrow \cong & & \updownarrow \cong \\
 \hat{\text{E}t}_{\text{Sp}}\zeta_n^{\mathcal{X}} : \hat{\text{E}t}_{\text{Sp}}(\Sigma_{n+1}^Q(Q \wedge \mathcal{X})) & \longrightarrow & \hat{\text{E}t}_{\text{Sp}}(\Sigma_n^Q \mathcal{X}).
 \end{array}$$

The upper and middle vertical isomorphisms on the left-hand side are the obvious level equivalences deduced from the canonical sequence of weak equivalences (4). The lower vertical isomorphisms are given by the following lemma applied to \mathcal{X} and $Q \wedge \mathcal{X}$, respectively.

Lemma 32. *For every integer $n \geq 0$ and every simplicial presheaf \mathcal{X} there is an isomorphism $F_n(\hat{\text{E}t}\mathcal{X}) \cong \hat{\text{E}t}_{\text{Sp}}(\Sigma_n^Q \mathcal{X})$ in $\text{Ho}(\mathcal{C}/\hat{\text{E}t}k)$.*

Proof. We define an intermediate object $E^n \in \mathcal{C}/\hat{\text{E}t}k$ given in degree $m \geq n$ by

$$E_m^n = (\hat{\text{E}t}Q)^{m-n} \wedge_{\hat{\text{E}t}k} \hat{\text{E}t}\mathcal{X}, (E^n)'_m = (\hat{\text{E}t}Q)^{m+1-n} \wedge_{\hat{\text{E}t}k} \hat{\text{E}t}\mathcal{X}, (E^n)''_m = (E^n)'_m$$

with the obvious structure maps induced by $S^2 \rightarrow \hat{\text{E}t}Q$ respectively the identity; in degree $m < n$ it is defined by $E_m^n = (E^n)'_m = (E^n)''_m = \hat{\text{E}t}k$ with identity maps and the map to the terminal object $\hat{\text{E}t}k$ of $\hat{\mathcal{S}}_*/\hat{\text{E}t}k$.

The object E^n is defined such that there are canonical maps

$$F_n(\hat{\text{E}t}\mathcal{X}) \xrightarrow{\alpha} E^n \xleftarrow{\beta} \hat{\text{E}t}_{\text{Sp}}(\Sigma_n^Q \mathcal{X})$$

induced in degree $m \geq n$ by the canonical weak equivalences of (4)

$$(S^2)^{m-n} \wedge \hat{\text{E}t}\mathcal{X} \xrightarrow{\cong} (\hat{\text{E}t}Q)^{m-n} \wedge_{\hat{\text{E}t}k} \hat{\text{E}t}\mathcal{X} \xleftarrow{\cong} \hat{\text{E}t}(Q^{m-n} \wedge_k \mathcal{X}).$$

It follows that the maps α and β are both level equivalences in $\mathcal{C}/\hat{\text{E}t}k$. Hence α and β are isomorphisms in $\text{Ho}(\mathcal{C}/\hat{\text{E}t}k)$. Their composition is the isomorphism $F_n(\hat{\text{E}t}\mathcal{X}) \cong \hat{\text{E}t}_{\text{Sp}}(\Sigma_n^Q \mathcal{X})$. \square

We deduce from the diagram that $\zeta_n^{\hat{\text{Et}}\mathcal{X}}$ and $\hat{\text{Et}}_{\text{Sp}}\zeta_n^{\mathcal{X}}$ differ only by an isomorphism in $\text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$ given by level equivalences. Since $\zeta_n^{\hat{\text{Et}}\mathcal{X}}$ is a stable equivalence in $\mathcal{C}/\hat{\text{Et}}k$ by definition, the map $\hat{\text{Et}}_{\text{Sp}}\zeta_n^{\mathcal{X}}$ is in fact an isomorphism in $\text{Ho}(\mathcal{C}/\hat{\text{Et}}k)$, which finishes the proof of the theorem. \square

Let MGL denote the motivic spectrum defined in [35] representing algebraic cobordism.

Proposition 33. *Let k be a separably closed field. There is an isomorphism in $\hat{\mathcal{S}}\mathcal{H}_2$*

$$\text{L}\hat{\text{Et}}(MGL) \cong \hat{M}U.$$

Proof. Let $G_k(n, N)$ be the Grassmannian over k and $G_k(n)$ be the colimit over N . By the Thom isomorphisms in étale and singular cohomology, it suffices to show that there is a weak equivalence between $\hat{\text{Et}}(G_k(n))$ and $\hat{G}_{\mathbb{C}}(n)$, the profinite completion of the simplicial set corresponding to the complex Grassmannian manifold.

If $\text{char } k > 0$, let R be the ring of Witt vectors of k , otherwise set $R = k$. By choosing a common embedding of R and \mathbb{C} into an algebraically closed field, Friedlander proved in [13, 3.2.2], that there is a natural sequence of \mathbb{Z}/ℓ -weak equivalences in $\text{pro-}\mathcal{S}_*$

$$\text{Sing}(G_{\mathbb{C}}(n, N)) \rightarrow \text{Et}(G_{\mathbb{C}}(n, N)) \rightarrow \text{Et}(G_R(n, N)) \leftarrow \text{Et}(G_k(n, N)).$$

We remark that weak equivalences in $\text{pro-}\mathcal{H}_*$ in the sense of [13] correspond to \mathbb{Z}/ℓ -weak equivalences in $\text{pro-}\mathcal{S}_*$ in the sense of [20]. By taking colimits with respect to N we get a sequence of \mathbb{Z}/ℓ -weak equivalences in $\text{pro-}\mathcal{S}_*$

$$\text{Sing}(G_{\mathbb{C}}(n)) \rightarrow \text{Et}(G_{\mathbb{C}}(n)) \rightarrow \text{Et}(G_R(n)) \leftarrow \text{Et}(G_k(n)).$$

Since \mathbb{Z}/ℓ -weak equivalences are preserved under completion, this shows that there is an isomorphism $\hat{G}_{\mathbb{C}}(n) \cong \hat{\text{Et}}(G_k(n))$ in $\hat{\mathcal{H}}_*$. \square

3.2.3. Étale realization of S^1 -spectra

The étale realization of S^1 -spectra is essentially simpler. The structure maps $\sigma_n : S_k^1 \wedge_k E_n \rightarrow E_{n+1}$ induce by base extension canonical maps $S_k^1 \wedge E_n \rightarrow S_k^1 \wedge_k E_n \xrightarrow{\sigma_n} E_{n+1}$ in $LU(k)$. Together with the isomorphism $\hat{\text{Et}} S_k^1 = S^1$ in $\hat{\mathcal{S}}_*$, we conclude that $\hat{\text{Et}}$ defines a functor on S^1 -spectra $\hat{\text{Et}} : \text{Sp}^{S^1}(k) \rightarrow \text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k)$.

Theorem 34. *The functor $\hat{\text{Et}}_{\text{Sp}} : \text{Sp}^{S^1}(k) \rightarrow \text{Sp}(\hat{\mathcal{S}}_*/\hat{\text{Et}}k)$ induces a functor $\text{L}\hat{\text{Et}}_{\text{Sp}} : \mathcal{SH}^{S^1}(k) \rightarrow \hat{\mathcal{S}}\mathcal{H}/\hat{\text{Et}}k$ on the stable \mathbb{A}^1 -homotopy category of S^1 -spectra.*

4. Profinite étale cohomology theories

4.1. General theory

Let k be a fixed base field and let ℓ be a fixed prime different from the characteristic of k . The first main application of the previous discussion is a canonical setting for étale cohomology theories for schemes, in particular étale topological cobordism.

Definition 35. Let $E \in \text{Sp}(\hat{\mathcal{S}}_*)$ be a profinite spectrum and let $X \in \text{Sm}/k$.

- (1) We define the *profinite étale cohomology of X in E* to be the profinite cohomology theory represented by E applied to the profinite space $\hat{\text{Et}} X$, i.e.

$$E_{\text{ét}}^n(X) := E^n(\hat{\text{Et}} X) = \text{Hom}_{\mathcal{S}\hat{\mathcal{H}}}(\Sigma^\infty(\hat{\text{Et}} X), E[n]),$$

where we add a base point if X is not already pointed.

- (2) We define relative étale cohomology groups $E_{\text{ét}}^n(X, U)$ for an open subscheme $U \subset X$ by $E_{\text{ét}}^n(X, U) := E^n(\hat{\text{Et}}(X)/\hat{\text{Et}}(U))$.
- (3) Similarly, if \mathcal{X} is a simplicial presheaf on Sm/k , we define the étale cohomology group of \mathcal{X} to be $E_{\text{ét}}^n(\mathcal{X}) := \text{Hom}_{\mathcal{S}\hat{\mathcal{H}}}(\Sigma^\infty(\hat{\text{Et}} \mathcal{X}), E[n])$.

The following fact may be shown in the standard way using Theorem 2.10 of [18].

Proposition 36. Let $U \xrightarrow{i} X \xleftarrow{p} V$ induce an elementary distinguished square [35]. Let E be a profinite spectrum. Then there is a Mayer–Vietoris long exact sequence of graded groups

$$\dots \rightarrow E_{\text{ét}}^n(X) \rightarrow E_{\text{ét}}^n(U) \oplus E_{\text{ét}}^n(V) \rightarrow E_{\text{ét}}^n(U \times_X V) \rightarrow E_{\text{ét}}^{n+1}(X) \rightarrow \dots$$

Proposition 37. Let E be a profinite spectrum. Let $U \subset X$ be an open subscheme of X . We get a long exact sequence of cohomology groups

$$\dots \rightarrow E_{\text{ét}}^n(X) \xrightarrow{j^*} E_{\text{ét}}^n(U) \xrightarrow{i^*} E_{\text{ét}}^n(X, U) \xrightarrow{i^*} E_{\text{ét}}^{n+1}(X) \xrightarrow{j^*} E_{\text{ét}}^{n+1}(U) \rightarrow \dots$$

where $j : \hat{\text{Et}} U \hookrightarrow \hat{\text{Et}} X$ and $i : (\hat{\text{Et}} X, \emptyset) \hookrightarrow (\hat{\text{Et}} X, \hat{\text{Et}} U)$ denote the natural induced inclusions.

Proof. Since $\hat{\text{Et}} U \hookrightarrow \hat{\text{Et}} X \rightarrow \hat{\text{Et}}(X)/\hat{\text{Et}}(U)$ is isomorphic to a cofiber sequence in $\hat{\mathcal{S}}_*$, this is just the usual long exact sequence of Hom-groups induced by a cofiber sequence in a simplicial model category. \square

Proposition 38. Let E be a profinite spectrum. Let $X \in \text{Sm}/k$. The projection $p : X \times \mathbb{A}^1 \rightarrow X$ induces an isomorphism $p^* : E_{\text{ét}}^*(X) \xrightarrow{\cong} E_{\text{ét}}^*(X \times \mathbb{A}^1)$.

Proof. This is clear since p induces a weak equivalence in $\hat{\mathcal{S}}$, see Theorem 27, and hence induces isomorphisms on cohomology theories. \square

Corollary 39. Let E be a profinite spectrum. Let $V \rightarrow X$ be an \mathbb{A}^n -bundle over X in Sm/k . Then $p^* : E_{\text{ét}}^*(X) \xrightarrow{\cong} E_{\text{ét}}^*(V)$ is an isomorphism.

We may summarize these results in the following theorem.

Theorem 40. Let E be a profinite spectrum. The étale cohomology theory $E_{\text{ét}}^*(-)$ represented by E satisfies the axioms of a cohomology theory on Sm/k of [30].

Proof. We check the axioms of a cohomology theory in the sense of [30, Definition 2.0.1].

1. *Localization:* This is clear from Proposition 37.

2. *Excision:* Let $e : (X', U') \rightarrow (X, U)$ be a morphism of pairs of schemes in Sm/k such that e is étale and for $Z = X - U, Z' = X' - U'$ one has $e^{-1}(Z) = Z'$ and $e : Z' \rightarrow Z$ is an isomorphism. By [26, III, Proposition 1.27], we know that the morphism e induces an isomorphism in étale cohomology $H^*(\hat{\text{Et}}(X)/\hat{\text{Et}}(U); \mathbb{Z}/\ell) \cong H^*(\hat{\text{Et}}(X')/\hat{\text{Et}}(U'); \mathbb{Z}/\ell)$. Hence the map $\hat{\text{Et}}(X')/\hat{\text{Et}}(U') \rightarrow \hat{\text{Et}}(X)/\hat{\text{Et}}(U)$ is an isomorphism in $\hat{\mathcal{H}}_*$. Therefore, it induces the desired isomorphism $E^*(\hat{\text{Et}}(X)/\hat{\text{Et}}(U)) \cong E^*(\hat{\text{Et}}(X')/\hat{\text{Et}}(U'))$ for every étale cohomology theory.

3. *Homotopy invariance:* This is the content of Proposition 38. \square

Theorem 41. *Let k be a field and let E be profinite spectrum such that each coefficient group E^q is finitely generated abelian. For every scheme X in Sm/k , there is a convergent spectral sequence*

$$E_2^{p,q} = H_{\text{ét}}^p(X; \mathbb{Z}/\ell^v \otimes E^q) \implies E_{\text{ét}}^{p+q}(X; \mathbb{Z}/\ell^v).$$

Proof. By the hypothesis on E^q the groups $\mathbb{Z}/\ell^v \otimes E^q$ are finite. The spectral sequence above is hence the one of Proposition 21 together with the isomorphisms $H_{\text{ét}}^*(X; \mathbb{Z}/\ell^v \otimes E^q) \cong H^*(\hat{\text{Et}} X; \mathbb{Z}/\ell^v \otimes E^q)$ of Remark 26. The condition for strong convergence in Proposition 21 is satisfied since $H_{\text{ét}}^p(X; \mathbb{Z}/\ell^v \otimes E^q)$ is a finite group for all p by [26, VI, Corollary 5.5]. \square

4.2. Examples

4.2.1. Profinite étale K-theory

Definition 42. We define the profinite étale K-theory of a smooth scheme X to be the cohomology theory represented by the profinitely completed spectrum $\hat{K}U$.

As a first application, we consider a comparison statement for profinite étale K-theory and for Friedlander’s étale K-theory in [13]. Let $\{X_s\} \in \text{pro-}S$ be a pro-object in S and let BU be the simplicial set representing complex K-theory. Friedlander defines the K-theory of $\{X_s\}$ for $\epsilon = 0, 1$ and $k > 0$ by

$$K^\epsilon(\{X_s\}; \mathbb{Z}/\ell^v) = \text{Hom}_{\text{pro-}\mathcal{H}}(\{\Sigma^\epsilon X_s\}, \{P^n BU \wedge C(\ell^v)\})$$

where $\{P^n BU\}$ denotes the Postnikov tower of BU considered as a pro-object in S and $C(\ell^v)$ is the cofibre of the multiplication by ℓ^v map $S^1 \rightarrow S^1$. For a locally Noetherian scheme X , Friedlander defines the étale K-theory by $K_{\text{ét}}^\epsilon(X; \mathbb{Z}/\ell^v) := K^\epsilon(\text{Et } X; \mathbb{Z}/\ell^v)$. By the Atiyah–Hirzebruch spectral sequences for $\hat{K}U$ and $K_{\text{ét}}$ of Theorem 41 and of [13], respectively, the following result is an immediate consequence.

Proposition 43. *For every scheme X of finite type over a separably closed field k the étale K-theory groups $K_{\text{ét}}^*(X; \mathbb{Z}/\ell^v)$ of [13] are isomorphic to the profinite étale K-theory groups $\hat{K}U_{\text{ét}}^*(X; \mathbb{Z}/\ell^v)$ defined above.*

4.2.2. *Profinite étale Morava K-theory*

Definition 44. We define the profinite étale Morava K-theory of a smooth scheme X to be the cohomology theory represented by the profinitely completed Morava K-theory spectrum $\hat{K}(n)$ for $n \geq 0$.

This theory satisfies in particular the conditions of Theorem 41. For $n \geq 1$, the coefficients are given by $\hat{K}(n)^* = \mathbb{Z}/\ell[v_n, v_n^{-1}]$, where $\deg v_n = -2(\ell^n - 1)$. In particular, for $n \geq 1$, the groups $\hat{K}(n)^q$ are finite for each q . Hence there is a converging spectral sequence from étale cohomology to étale Morava K-theory for $n \geq 1$:

$$E_2^{p,q} = H_{\hat{\text{ét}}}^p(X; \hat{K}(n)^q) \implies K(\hat{n})_{\hat{\text{ét}}}^{p+q}(X).$$

For schemes over \mathbb{C} , it agrees with the usual topological Morava K-theory. This follows as for the étale cobordism below from the generalized Riemann Existence Theorem of [2].

4.2.3. *Profinite étale cobordism*

Definition 45. We define the *profinite étale cobordism* of a smooth scheme $X \in \text{Sm}/k$, to be the étale cohomology theory represented by the profinite cobordism spectrum $\hat{M}U$.

Proposition 46. *Let R be a strict local Henselian ring. Then*

$$\hat{M}U_{\hat{\text{ét}}}^*(\text{Spec } R) \cong MU^* \otimes_{\mathbb{Z}} \mathbb{Z}_{\ell} \quad \text{and} \quad \hat{M}U_{\hat{\text{ét}}}^*(\text{Spec } R; \mathbb{Z}/\ell^v) \cong MU^* \otimes_{\mathbb{Z}} \mathbb{Z}/\ell^v.$$

Proof. This may be easily deduced from Example 20, since $\hat{\text{Et}} \text{Spec } R$ is contractible. The last assertion follows since MU^* has no torsion. \square

Proposition 47. *The (reduced) étale cobordism of a finite field k , $\text{char } k \neq \ell$, is given by the isomorphism $\hat{M}U_{\hat{\text{ét}}}^n(k) = \hat{M}U^{n-1}$ and similarly for $\hat{M}U_{\hat{\text{ét}}}^*(k; \mathbb{Z}/\ell^v)$.*

Proof. The profinite space $\hat{\text{Et}} k$ has the homotopy type of the circle S^1 in $\hat{\mathcal{S}}$. \square

The generalized Riemann Existence Theorem of [2, Theorem 12.9], or the comparison theorem for étale cohomology and the Atiyah–Hirzebruch spectral sequences for complex and étale cobordism imply the following proposition.

Proposition 48. *Let X be an algebraic variety over \mathbb{C} . Let $X(\mathbb{C})$ be the topological space of complex points. For every v , there is an isomorphism*

$$\hat{M}U_{\hat{\text{ét}}}^*(X; \mathbb{Z}/\ell^v) \cong MU^*(X(\mathbb{C}); \mathbb{Z}/\ell^v).$$

There are several other obvious applications of the Riemann Existence Theorem and the Atiyah–Hirzebruch spectral sequence. For example, we can deduce a proper base change theorem or an isomorphism for separably closed field extensions etc. from the corresponding theorems for étale cohomology. Furthermore, we can calculate the étale cobordism of a smooth projective curve over a separably closed field.

Finally, we remark that the morphism of profinite spectra $\hat{M}U \rightarrow H\mathbb{Z}/\ell^v$ induced by the orientation yields a unique map of profinite étale cohomology theories $\hat{M}U_{\hat{\text{ét}}}^*(X; \mathbb{Z}/\ell^v) \rightarrow H_{\hat{\text{ét}}}^*(X; \mathbb{Z}/\ell^v)$ for every X in Sm/k .

5. Algebraic versus étale cobordism

The main application of the étale realization functor is existence of a canonical map from algebraic to étale cobordism. We have to show that étale cobordism is an oriented theory on Sm/k .

5.1. Étale cobordism is an oriented cohomology theory

We prove that $\hat{M}U_{\text{ét}}^{2*}(-; \mathbb{Z}/\ell)$ is an oriented cohomology theory in the sense of [25] on the category Sm/k of smooth quasi-projective schemes over a suitable base field. The key point is the projective bundle formula.

Let k be a field and let G_k be its absolute Galois-group. We have to restrict our attention to the fields with finite ℓ -cohomological dimension satisfying the following property.

Definition 49.

- (1) A profinite space X is said to be *without ℓ -torsion* if the canonical map $H^*(X; \mathbb{Z}/\ell^\nu) \rightarrow H^*(X; \mathbb{Z}/\ell)$ is surjective for all integers $\nu \geq 0$.
- (2) A field k is said to be *without ℓ -torsion* if the canonical map in Galois-cohomology $H^*(G_k; \mathbb{Z}/\ell^\nu) \rightarrow H^*(G_k; \mathbb{Z}/\ell)$ is surjective for all integers $\nu \geq 0$.

Example 50. 1. Every separably closed field is without ℓ -torsion.

2. A finite field has no ℓ -torsion.

3. Local fields are without ℓ -torsion if ℓ is different from the residue characteristic of k .

4. Let \mathbb{P}_k^n be the projective space of dimension n over k . If k is a field without ℓ -torsion then the profinite space $\hat{\text{Ét}}\mathbb{P}_k^n$ has no ℓ -torsion in $\hat{\mathcal{S}}$ for every n . This follows from the projective bundle formula for étale cohomology.

In the rest of this chapter we will always assume that k is a field without ℓ -torsion and with finite ℓ -cohomological dimension.

Proposition 51. *The étale cobordism of \mathbb{P}_k^n is given by the direct sum*

$$\hat{M}U_{\text{ét}}^*(\mathbb{P}_k^n; \mathbb{Z}/\ell) \cong \bigoplus_{i=0}^{n-1} \hat{M}U_{\text{ét}}^*(k; \mathbb{Z}/\ell).$$

Proof. Since $\hat{\text{Ét}}\mathbb{P}_k^n$ is a space without ℓ -torsion, the corresponding Atiyah–Hirzebruch spectral sequence collapses at the E_2 -term and gives an isomorphism

$$\hat{M}U^*(\hat{\text{Ét}}\mathbb{P}_k^n; \mathbb{Z}/\ell) \xrightarrow{\cong} \bigoplus_s H^s(\hat{\text{Ét}}\mathbb{P}_k^n; \mathbb{Z}/\ell \otimes MU^{*-s}), \tag{5}$$

as described in [8, Proposition 2.1.9]. The assertion now follows from the projective bundle formula for étale cohomology. \square

I am grateful to Francois-Xavier Dehon for an explanation of the arguments proving the following proposition.

Proposition 52. *For the projective n -space and every smooth scheme X over k the canonical map*

$$\hat{M}U_{\text{ét}}^*(\mathbb{P}_k^n; \mathbb{Z}/\ell) \otimes_{\hat{M}U_{\text{ét}}^*(k; \mathbb{Z}/\ell)} \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell) \xrightarrow{\cong} \hat{M}U_{\text{ét}}^*(\mathbb{P}_k^n \times_k X; \mathbb{Z}/\ell)$$

is an isomorphism.

Proof. This is a classical argument. In the profinite setting, this has been worked out by Dehon. The case with \mathbb{Z}/ℓ -coefficients can be done in a similar way. It is deduced from the Künneth formula for the \mathbb{Z}/ℓ -cohomology of $\hat{\text{E}}t \mathbb{P}_k^n \wedge_{\hat{\text{E}}t k} \hat{\text{E}}t X$. The point is that $\hat{M}U_{\text{ét}}^*(\mathbb{P}_k^n; \mathbb{Z}/\ell)$ (respectively $H_{\text{ét}}^*(\mathbb{P}_k^n; \mathbb{Z}/\ell)$) is a free $\hat{M}U_{\text{ét}}^*(k; \mathbb{Z}/\ell)$ -module (respectively $H_{\text{ét}}^*(k; \mathbb{Z}/\ell)$ -module). \square

Let $\mathcal{O}(1) \rightarrow \mathbb{P}_k^n$ be the canonical quotient line bundle. We conclude from the proof of Proposition 51 that the orientation map $\hat{M}U^*(\hat{\text{E}}t \mathbb{P}_k^n) \rightarrow H^*(\hat{\text{E}}t \mathbb{P}_k^n; \mathbb{Z}/\ell)$ factors through the isomorphism (5). In particular, this implies that the element $\xi_H = c_1(\mathcal{O}(1)) \in H^2(\hat{\text{E}}t \mathbb{P}_k^n; \mathbb{Z}/\ell)$ induces an element $\xi_{\hat{M}U} \in \hat{M}U^2(\hat{\text{E}}t \mathbb{P}_k^n; \mathbb{Z}/\ell)$, which corresponds to a morphism in $\hat{\mathcal{S}}\mathcal{H}$:

$$\xi_{\hat{M}U}: \hat{\Sigma}^\infty(\hat{\text{E}}t \mathbb{P}_k^n) \longrightarrow \hat{M}U/\ell \wedge S^2.$$

Now let $E \rightarrow X$ be a vector bundle over X in Sm/k and let $\mathcal{O}(1)$ be the canonical quotient line bundle over $\mathbb{P}(E)$. This bundle determines a morphism $\mathbb{P}(E) \rightarrow \mathbb{P}_k^N$ for some sufficiently large N . Together with the morphism $\xi_{\hat{M}U}$ we get an element $\xi_{\hat{M}U} \in \hat{M}U_{\text{ét}}^2(\mathbb{P}(E); \mathbb{Z}/\ell)$.

Theorem 53 (Projective Bundle Formula). *Let $E \rightarrow X$ be a rank n vector bundle over X in Sm/k . Then $\hat{M}U_{\text{ét}}^*(\mathbb{P}(E); \mathbb{Z}/\ell)$ is a free $\hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell)$ -module with basis $(1, \xi, \xi^2, \dots, \xi^{n-1})$.*

Proof. We prove the assertion first for the case of a trivial bundle on X . As in Lemma 28, one proves that the canonical morphism of profinite spaces $\hat{\text{E}}t(X) \times_{\hat{\text{E}}t k} \hat{\text{E}}t(\mathbb{P}_k^n) \rightarrow \hat{\text{E}}t(X \times_k \mathbb{P}_k^n)$ is a weak equivalence in $\hat{\mathcal{S}}$. Hence this case follows from Propositions 51 and 52. For the general case, since E is locally trivial for the Zariski topology on X , it suffices to show that the theorem holds for X if it holds for open subsets X_0, X_1 and $X_0 \cap X_1$, with $X = X_0 \cup X_1$. This may be checked by a standard argument using the Mayer–Vietoris sequence. \square

We use Grothendieck’s idea to introduce higher Chern classes for vector bundles.

Definition 54. Let E be a vector bundle of rank n on X in Sm/k and let $\xi_{\hat{M}U}$ be as above. Then we define the i th Chern class of E to be the unique element $c_i(E) \in \hat{M}U_{\text{ét}}^{2i}(X; \mathbb{Z}/\ell)$ such that $c_0(E) = 1, c_i(E) = 0$ for $i > n$ and $\sum_{i=0}^n (-1)^i c_i(E) \xi_{\hat{M}U}^{n-i} = 0$.

The results of Panin in [30] then imply the following theorem.

Theorem 55. *The étale cobordism $\hat{M}U_{\text{ét}}^{2*}(-; \mathbb{Z}/\ell)$ is an oriented ring cohomology theory on Sm/k in the sense of [30, Definition 3.1.1]. This implies that for every projective morphism $f : Y \rightarrow X$ of codimension d there are natural transfer maps*

$$f_* : \hat{M}U_{\text{ét}}^*(Y; \mathbb{Z}/\ell) \rightarrow \hat{M}U_{\text{ét}}^{*+2d}(X; \mathbb{Z}/\ell)$$

which satisfy the axioms of a push-forward in an oriented cohomology theory in the sense of [25].

Remark 56. 1. If k is separably closed, there are several simplifications. First of all, the above statements all hold without \mathbb{Z}/ℓ -coefficients, i.e. the Projective Bundle and the Künneth Formula hold for $\hat{M}U$ and $\hat{M}U/\ell^v$ as well.

Secondly, the choice of an orientation is canonical. Using the isomorphism $\text{L}\hat{\text{E}}tMGL \cong \hat{M}U$, we choose $x_{\hat{M}U} \in \hat{M}U^2(\hat{\text{E}}t\mathbb{P}_k^\infty)$ as the image of the orientation $x_{MGL} : \mathbb{P}^\infty \rightarrow MGL \wedge \mathbb{P}^1$ under $\hat{\text{E}}t$.

2. A priori, the definition of $E(\mathbb{P}^\infty) := \lim_n E(\mathbb{P}_k^n)$ for a cohomology theory E in [30, 1.1] might differ from our definition. But since $\hat{M}U_{\text{ét}}^i(\mathbb{P}_k^n; \mathbb{Z}/\ell)$ is a finite group for every i and n by Proposition 51, the \lim_n^1 of these groups vanishes. Hence there is a canonical isomorphism $\hat{M}U_{\text{ét}}^i(\mathbb{P}_k^\infty; \mathbb{Z}/\ell) \cong \lim_n \hat{M}U_{\text{ét}}^i(\mathbb{P}_k^n; \mathbb{Z}/\ell)$ for $\mathbb{P}_k^\infty = \text{colim}_n \mathbb{P}_k^n$. Hence, in the case of étale cobordism with \mathbb{Z}/ℓ -coefficients, Panin’s and our definitions are compatible.

5.2. Comparison with Ω^*

We consider the algebraic cobordism theory $\Omega^*(-)$ of [25]. As a corollary of Theorem 55, using the universality of Ω^* we get the following theorem.

Theorem 57. *There is a canonical morphism of oriented cohomology theories*

$$\theta : \Omega^*(X) \rightarrow \Omega^*(X; \mathbb{Z}/\ell) \rightarrow \hat{M}U_{\text{ét}}^{2*}(X; \mathbb{Z}/\ell)$$

defined by sending a generator $[f : Y \rightarrow X] \in \Omega^*(X)$, $f : Y \rightarrow X$ a projective morphism between smooth schemes, to the element $f_*(1_Y) \in \hat{M}U_{\text{ét}}^{2*}(X; \mathbb{Z}/\ell)$.

If k is a field of characteristic zero, Theorem 4 of [25] states that $\Omega^*(k)$ is isomorphic to the Lazard ring \mathbb{L}^* . It is conjectured that this isomorphism holds for every field, see the conjecture of [25, §4.3.2].

Proposition 58. *For every separably closed field k , the morphism*

$$\theta : \Omega^*(k; \mathbb{Z}/\ell^v) \rightarrow \hat{M}U_{\text{ét}}^{2*}(k; \mathbb{Z}/\ell^v)$$

is surjective.

If we suppose in addition that the conjecture of [25] is true for k , then θ is an isomorphism.

Proof. Consider the canonical map $\Phi : \mathbb{L}^* \otimes \mathbb{Z}/\ell^v \rightarrow \Omega^*(k; \mathbb{Z}/\ell^v)$. It is split injective for every field, see [25, Corollary 4.3.3]. When we compose this map with $\theta : \Omega^*(k; \mathbb{Z}/\ell^v) \rightarrow \hat{M}U_{\text{ét}}^{2*}(k; \mathbb{Z}/\ell^v)$, we get the canonical map $\mathbb{L}^* \otimes \mathbb{Z}/\ell^v \xrightarrow{\Phi} \Omega^*(k; \mathbb{Z}/\ell^v) \xrightarrow{\theta} \hat{M}U_{\text{ét}}^{2*}(k; \mathbb{Z}/\ell^v)$.

Since this map is unique, it must be the canonical isomorphism. Hence $\theta : \Omega^*(k; \mathbb{Z}/\ell^v) \rightarrow \hat{M}U_{\text{ét}}^{2*}(k; \mathbb{Z}/\ell^v)$ is surjective. The last assertion follows in the same way if we assume the conjecture to be true. \square

5.3. Comparison with $MGL^{*,*}$

Let k be a field as above. Let V be a vector bundle of rank d over X in Sm/k . We recall that the Thom space $\text{Th}(V) \in LU(k)$ of V is defined to be the quotient $\text{Th}(V) = V/(V - i(X))$, where $i : X \rightarrow V$ denotes the zero section of V . We reformulate a lemma from \mathbb{A}^1 -homotopy theory.

Proposition 59. *Let V be a vector bundle over X and $\mathbb{P}(V) \rightarrow \mathbb{P}(V \oplus \mathcal{O})$ be the closed embedding at infinity. Then the canonical morphism of pointed sheaves $\mathbb{P}(V \oplus \mathcal{O})/\mathbb{P}(V) \rightarrow \text{Th}(V)$ induces a weak equivalence in $\hat{\mathcal{S}}_*$ via $\hat{\text{Ét}}$.*

Proof. This is the same proof as for Proposition 3.2.17 of [29] where we use the fact that $\hat{\text{Ét}}$ preserves \mathbb{A}^1 -weak equivalences between smooth schemes by Theorem 27 and commutes with quotients. \square

Now we define the Thom class of V in $\hat{M}U_{\text{ét}}^{2d}(\text{Th}(V); \mathbb{Z}/\ell)$. From the isomorphism $\mathbb{P}(V \oplus \mathcal{O})/\mathbb{P}(V) \cong \text{Th}(V)$ we deduce an exact sequence induced by the corresponding cofiber sequence

$$\hat{M}U_{\text{ét}}^*(\text{Th}(V); \mathbb{Z}/\ell) \rightarrow \hat{M}U_{\text{ét}}^*(\mathbb{P}(V \oplus \mathcal{O}); \mathbb{Z}/\ell) \rightarrow \hat{M}U_{\text{ét}}^*(\mathbb{P}(V); \mathbb{Z}/\ell).$$

Using the projective bundle formula of Theorem 53 this sequence is isomorphic to the exact sequence

$$\hat{M}U_{\text{ét}}^*(\text{Th}(V); \mathbb{Z}/\ell) \rightarrow \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell)[1, u, \dots, u^d] \rightarrow \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell)[1, \dots, u^{d-1}].$$

The element $x := u^d - c_1(V)u^{d-1} + \dots + (-1)^d c_d(V) \in \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell)[1, u, \dots, u^d]$ is sent to $u^d - c_1(V)u^{d-1} + \dots + (-1)^d c_d(V) \in \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell)[1, u, \dots, u^{d-1}]$ which is 0 by the definition of Chern classes. By exactness, there is an element $\text{th}_{\text{ét}}(V) \in \hat{M}U_{\text{ét}}^{2d}(\text{Th}(V); \mathbb{Z}/\ell)$ that is sent to $x \in \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell)[1, u, \dots, u^d]$. We call $\text{th}_{\text{ét}}(V)$ the *étale Thom class of V* . It corresponds to a morphism in $\hat{\mathcal{S}}\mathcal{H}$

$$\text{th}_{\text{ét}}(V) : \hat{\Sigma}^\infty(\hat{\text{Ét}}\text{Th}(V)) \rightarrow \hat{M}U/\ell \wedge S^{2d}.$$

We apply this argument to the tautological n -bundle γ_n over the infinite Grassmannian. Since $MGL_n = \text{Th}(\gamma_n)$, we get a morphism in $\hat{\mathcal{S}}\mathcal{H}_2$

$$\phi : \text{L}\hat{\text{Ét}}(MGL) \longrightarrow \hat{M}U/\ell.$$

Recall the definition of algebraic cobordism for a scheme X in Sm/k

$$MGL^{p,q}(X) := \text{Hom}_{\mathcal{S}\mathcal{H}(k)}(\Sigma_{\mathbb{P}^1}^\infty(X), MGL[p - 2q] \wedge (\mathbb{P}^1)^{\wedge q}),$$

where $MGL[n] := MGL \wedge (S_s^1)^{\wedge n}$, S_s^1 denoting the simplicial circle. One sees that the elements in degree p, q are sent to elements in degree p via ϕ . We summarize this discussion in the following theorem.

Theorem 60. *There is a natural map for every X in Sm/k*

$$\phi : MGL^{p,q}(X; \mathbb{Z}/\ell) \rightarrow \hat{M}U_{\text{ét}}^p(X; \mathbb{Z}/\ell)$$

which defines a morphism of oriented cohomology theories.

By the universality of $\Omega^*(-)$, we get the following corollary.

Corollary 61. *For every $X \in \text{Sm}/k$, there is a canonical commutative diagram of morphisms of oriented cohomology theories*

$$\begin{array}{ccc} \Omega^*(X; \mathbb{Z}/\ell) & \xrightarrow{\theta_{MGL}} & MGL^{2*,*}(X; \mathbb{Z}/\ell) \\ & \searrow \theta_{\hat{M}U} & \swarrow \phi \\ & \hat{M}U_{\text{ét}}^{2*}(X; \mathbb{Z}/\ell) & \end{array}$$

5.4. The Galois action on étale cobordism

Let k be a field of characteristic $p \neq \ell$ and let \bar{k} be a separable closure of k . If X is a scheme over k , there is a natural action on $X_{\bar{k}} = X \otimes_k \bar{k}$ of the Galois group $G_k := \text{Gal}(\bar{k}/k)$ of k . By naturality G_k also acts on $\hat{\text{Ét}} X_{\bar{k}}$.

Now $\Omega^*(\bar{k})$ is generated by classes $[\pi : Y \rightarrow \text{Spec } \bar{k}]$ of smooth projective schemes Y over \bar{k} , see [25, §2.5.4]. Hence in order to determine the action of the absolute Galois group on étale cobordism, it suffices to know the action on each such Y . Since \bar{k} is separably closed, the induced morphism $\sigma : Y \rightarrow Y^\sigma$ induces an isomorphism on étale cohomology with \mathbb{Z}/ℓ -coefficients, where Y^σ is the twisted \bar{k} -scheme with structure morphism $\sigma \circ \pi$. Hence σ induces an isomorphism of homotopy classes $\sigma : \hat{\text{Ét}} Y_{\bar{k}} \xrightarrow{\cong} \hat{\text{Ét}} Y^\sigma$ in $\hat{\mathcal{H}}$. This implies the following proposition.

Proposition 62. *The action of G_k on $\hat{M}U_{\text{ét}}^*(\bar{k}; \mathbb{Z}/\ell^v)$ is trivial.*

Next, we calculate the profinite étale cobordism groups of a local field, i.e. either a finite extension of the field \mathbb{Q}_p or a finite extension of the field of formal power series $\mathbb{F}((t))$ over a finite field of characteristic p . We assume $p \neq \ell$ and we denote by $q = p^f$ the number of elements in the residue class field of k . Let $\ell^{v_0} = (q - 1, \ell^v)$ be the greatest common divisor of $q - 1$ and ℓ^v .

Corollary 63. *Let k be a local field. For all n , the (reduced) profinite étale cobordism groups with \mathbb{Z}/ℓ^v -coefficients of k are given by*

$$\hat{M}U_{\text{ét}}^n(k; \mathbb{Z}/\ell^v) = \begin{cases} \mathbb{Z}/\ell^{v_0} \otimes MU^{n-2}, & n \text{ even,} \\ \mathbb{Z}/\ell^{v_0} \otimes MU^{n-1}, & n \text{ odd.} \end{cases}$$

Proof. Since $\mathbb{Z}/\ell^v \otimes MU^t$ is a finitely generated free \mathbb{Z}/ℓ^v -module with trivial G_k -action, we may identify the Galois cohomology groups $H^i(k; \mathbb{Z}/\ell^v \otimes MU^t)$ with $H^i(k; \mathbb{Z}/\ell^v) \otimes MU^t$. The assertion follows from the local Tate Duality for Galois cohomology and the Atiyah–Hirzebruch spectral sequence. \square

5.5. *Inverting the Bott element*

Let $H^p(X; \mathbb{Z}/n(q))$ denote the motivic cohomology of a smooth scheme X over a field k . For $\text{Spec } k$ there is an isomorphism $H^0(\text{Spec } k; \mathbb{Z}/n(1)) \cong \mu_n(k)$ with the group of n th roots of unity in k . Assuming that k contains an n th root of unity ζ , we have a corresponding motivic Bott element $\beta_n \in H^0(\text{Spec } k; \mathbb{Z}/n(1))$. Levine has shown in [24] that motivic \mathbb{Z}/n -cohomology of a smooth scheme over k agrees with étale \mathbb{Z}/n -cohomology after inverting the Bott element. I am grateful to Marc Levine for an explanation of his ideas.

Furthermore, Hopkins and Morel announced the construction of a motivic Atiyah–Hirzebruch spectral sequence. It is the slice filtration spectral sequence conjectured by Voevodsky in [36] from motivic cohomology with coefficients in the ring MU^* to algebraic cobordism

$$E_2^{p,q,2i} = H^{p,q}(X, MU^{2i}) \implies MGL^{p+2i,q+i}(X) \tag{6}$$

with differentials $d_{2r+1} : E_{2r+1}^{p,q,2i} \rightarrow E_{2r+1}^{p+2i+1,q+i,2i-2r}$.

We consider the spectral sequence for $\text{Spec } k$. Since $E_2^{p,1,q}(\text{Spec } k)$ is concentrated in degrees $p = 0$ and $p = 1$, we deduce $MGL^{0,1}(k) \cong k^\times$. For \mathbb{Z}/n -coefficients, the exact sequence for coefficients implies that we get an isomorphism $MGL^{0,1}(k; \mathbb{Z}/n) \cong \mu_n(k)$ and, via the spectral sequence, the motivic Bott element defined above, is sent to an induced *Bott element* $\beta_n \in MGL^{0,1}(k; \mathbb{Z}/n)$.

Let us now suppose that k is algebraically closed, $n = \ell^v$ and $\text{char } k \neq \ell$. In particular, k contains an ℓ^v th root of unity ζ . It defines an element $\zeta \in H_{\text{ét}}^0(\text{Spec } k; \mu_{\ell^v})$. The element $\zeta \cdot 1_{\hat{M}U/\ell^v}$ induces via the Atiyah–Hirzebruch spectral sequence an element $\zeta_{\hat{M}U/\ell^v} \in \hat{M}U_{\text{ét}}^0(\text{Spec } k; \mathbb{Z}/\ell^v)$. The multiplication with ζ yields an isomorphism of spectral sequences and hence multiplication with $\zeta_{\hat{M}U/\ell^v}$ is an isomorphism on étale cobordism. Furthermore, the map ϕ sends β_{ℓ^v} to $\zeta_{\hat{M}U/\ell^v}$. This implies that ϕ induces a localized map $\phi : MGL^{*,*}(X; \mathbb{Z}/\ell^v)[\beta^{-1}] \rightarrow \hat{M}U_{\text{ét}}^*(X; \mathbb{Z}/\ell^v)$.

Finally, motivic cohomology may be represented in $\mathcal{SH}(k)$ by the Eilenberg–Mac Lane spectrum $H_{\mathbb{Z}/\ell^v}$, defined in degree n by the group completion of the symmetric products

$$K(\mathbb{Z}/\ell^v(n), 2n) = \left(\coprod_{d \geq 0} S^d((\mathbb{P}^1)^{\wedge n}) \right)^+ \otimes \mathbb{Z}/\ell^v.$$

Using the simplicial version of the Dold–Thom theorem and the Künneth isomorphism for symmetric products in étale cohomology over an algebraically closed field of [3, Exposé XVII, Théorème 5.5.21], one shows as before that the étale realization of $H_{\mathbb{Z}/\ell^v}$ is isomorphic in \mathcal{SH} to the profinite Eilenberg–Mac Lane spectrum representing \mathbb{Z}/ℓ^v -cohomology. Hence the étale realization defines a map from motivic \mathbb{Z}/ℓ^v -cohomology to étale \mathbb{Z}/ℓ^v -cohomology. With a fixed isomorphism $\mathbb{Z}/\ell^v(1) \cong \mathbb{Z}/\ell^v$, this map is the unique map of oriented cohomology theories. Furthermore, localization at β is exact. Consequently, the étale realization yields a map

of Atiyah–Hirzebruch spectral sequences, whose E_2 -terms agree by Theorem 1.1 of [24]. This implies the following theorem.

Conditional Theorem 64. *Let X be a smooth scheme of finite type over an algebraically closed field k with $\text{char } k \neq \ell$. If we assume the existence and convergence of the above Atiyah–Hirzebruch spectral sequence (6) from motivic cohomology to algebraic cobordism, then ϕ is an isomorphism*

$$\phi : \left(\bigoplus_{p,q} MGL^{p,q}(X; \mathbb{Z}/\ell^v) \right) [\beta^{-1}] \xrightarrow{\cong} \bigoplus_{p,q} \hat{M}U_{\text{ét}}^p(X; \mathbb{Z}/\ell^v).$$

Acknowledgments

I would like to use this opportunity to thank first of all Christopher Deninger for the suggestion to work on this topic. He helped me with important suggestions and gave me the motivating impression to work on something that captured his interest. I am grateful to Marc Levine, Fabien Morel and Alexander Schmidt, who answered all my questions and explained to me both the great picture and the little details. I thank Francois-Xavier Dehon for an answer to a specific problem on Künneth isomorphisms for cobordism with coefficients. Finally, I would like to thank the kind referee who made several detailed and helpful suggestions.

References

- [1] J.F. Adams, Stable Homotopy and Generalised Homology, Chicago Lecture Notes in Math., Chicago Univ. Press, 1974.
- [2] M. Artin, B. Mazur, Etale Homotopy, Lecture Notes in Math., vol. 100, Springer-Verlag, 1969.
- [3] M. Artin, A. Grothendieck, J.-L. Verdier, Séminaire de Géométrie Algébrique, SGA 4, T. 3, Lecture Notes in Math., vol. 305, Springer-Verlag, 1973.
- [4] A.K. Bousfield, E.M. Friedlander, Homotopy of Γ -spaces, spectra and bisimplicial sets, in: Proc. of the Conference Geometric Applications of Homotopy Theory, vol. II, Evanston, IL, 1977, in: Lecture Notes in Math., vol. 658, Springer-Verlag, 1978, pp. 80–130.
- [5] A.K. Bousfield, D.M. Kan, Homotopy Limits, Completions and Localizations, Lecture Notes in Math., vol. 304, Springer-Verlag, 1972.
- [6] J.D. Christensen, D.C. Isaksen, Duality and pro-spectra, *Algebr. Geom. Topol.* 4 (2004) 781–812.
- [7] D. Cox, Homotopy theory of simplicial schemes, *Compos. Math.* 39 (1979) 263–295.
- [8] F.-X. Dehon, Cobordisme complexe des espaces profinis et foncteur T de Lannes, *Mém. Soc. Math. France* 98 (2004).
- [9] D. Dugger, Universal homotopy theories, *Adv. Math.* 164 (1) (2001) 144–176.
- [10] B.I. Dundas, O. Röndigs, P.A. Østvær, Motivic functors, *Doc. Math.* 8 (2003) 489–525.
- [11] W. Dwyer, E.M. Friedlander, Algebraic and étale K-theory, *Trans. Amer. Math. Soc.* 292 (1) (1985) 247–280.
- [12] D.A. Edwards, H.M. Hastings, Čech and Steenrod Homotopy Theory with Applications to Geometric Topology, Lecture Notes in Math., vol. 542, Springer-Verlag, 1976.
- [13] E.M. Friedlander, Etale K-theory I: Connections with étale cohomology and algebraic vector bundles, *Invent. Math.* 60 (1980) 105–134.
- [14] E.M. Friedlander, Etale Homotopy of Simplicial Schemes, *Ann. of Math. Stud.*, vol. 104, Princeton Univ. Press, 1982.
- [15] P.S. Hirschhorn, Model Categories and Their Localizations, *Math. Surveys Monogr.*, vol. 99, Amer. Math. Soc., 2003.
- [16] M. Hovey, Spectra and symmetric spectra in general model categories, *J. Pure Appl. Algebra* 165 (2001) 63–127.
- [17] M. Hovey, B. Shipley, J. Smith, Symmetric spectra, *J. Amer. Math. Soc.* 13 (2000) 149–208.
- [18] D.C. Isaksen, Etale realization on the \mathbb{A}^1 -homotopy theory of schemes, *Adv. Math.* 184 (2004) 37–63.

- [19] D.C. Isaksen, Generalized cohomology of pro-spectra, preprint, arXiv: math.AT/0408179 v1, 2004.
- [20] D.C. Isaksen, Completions of pro-spaces, *Math. Z.* 250 (2005) 113–143.
- [21] D.C. Isaksen, Flasque model structures for simplicial presheaves, *K-Theory* 36 (2006) 371–395.
- [22] J.F. Jardine, Motivic symmetric spectra, *Doc. Math.* 5 (2000) 445–553.
- [23] R. Joshua, Spanier–Whitehead duality in étale homotopy, *Trans. Amer. Math. Soc.* 296 (1) (1986) 151–166.
- [24] M. Levine, Inverting the motivic Bott element, *K-Theory* 19 (1) (2000) 1–28.
- [25] M. Levine, F. Morel, *Algebraic Cobordism*, Springer Monogr. Math., Springer-Verlag, 2007.
- [26] J.S. Milne, *Étale Cohomology*, Princeton Univ. Press, 1980.
- [27] F. Morel, Ensembles profinis simpliciaux et interprétation géométrique du foncteur T, *Bull. Soc. Math. France* 124 (1996) 347–373.
- [28] F. Morel, On the motivic π_0 of the sphere spectrum, in: J.P.C. Greenlees (Ed.), *Axiomatic, Enriched and Motivic Homotopy Theory*, Kluwer Acad. Publ., 2004, pp. 219–260.
- [29] F. Morel, V. Voevodsky, \mathbb{A}^1 -homotopy theory of schemes, *Inst. Hautes Études Sci. Publ. Math.* 90 (1999) 45–143.
- [30] I. Panin, Oriented cohomology theories of algebraic varieties, *K-Theory* 30 (2003) 265–314.
- [31] G. Quick, Profinite étale cobordism, preprint, arXiv: math.AG/0506332, 2005.
- [32] A. Schmidt, On the étale homotopy type of Voevodsky’s spaces, preprint, www.math.uiuc.edu/K-theory/0675, 2003.
- [33] V.P. Snaith, Algebraic cobordism and K-theory, *Mem. Amer. Math. Soc.* 21 (221) (1979).
- [34] R.W. Thomason, Algebraic K-theory and étale cohomology, *Ann. Sci. École Norm. Sup.* 13 (1980) 437–552.
- [35] V. Voevodsky, \mathbb{A}^1 -homotopy theory, in: *Proceedings of the Internat. Congress of Mathematicians, Berlin, 1998*, in: *Doc. Math.*, vol. I (extra), 1998, pp. 579–604.
- [36] V. Voevodsky, Open problems in the motivic stable homotopy theory, I, in: *Motives, Polylogarithms and Hodge Theory, Part I*, Irvine, CA, 1998, in: *Int. Press Lect. Ser.*, vol. 3 I, Int. Press, Somerville, MA, 2002, pp. 3–34.