A NOTE ON THE THOM MORPHISM FOR THE CLASSIFYING SPACE OF CERTAIN LIE GROUPS AND GAUGE GROUPS

EIOLF KASPERSEN AND GEREON QUICK

ABSTRACT. We give a complete description of which non-torsion generators are not in the image of the Thom morphism from complex cobordism to integral cohomology for the classifying space of exceptional Lie groups except for E_8 . We then show that the Thom morphism is not surjective for the classifying space of the gauge group of a principal E_7 -bundle over the four-dimensional sphere. We use the results to detect nontrivial elements in the kernel of the reduced Thom morphism for Lie groups and their classifying spaces.

1. INTRODUCTION

Let G be a compact, connected Lie group, and let ξ denote a principal G-bundle

$$G \longleftrightarrow P \xrightarrow{\pi} X$$

over a paracompact space X. Recall that the gauge group $\mathcal{G}(\xi)$ of ξ is defined to be the group of automorphisms of ξ , i.e.,

$$\mathcal{G}(\xi) = \{ \phi \in \operatorname{Aut}_G(P) \mid \pi \circ \phi = \pi \}.$$

Gauge groups play an important role in geometry, topology and mathematical physics. Moreover, the classifying space $B\mathcal{G}(\xi)$ is homotopy equivalent to the moduli space of connections on ξ . Integral singular cohomology $H^*(B\mathcal{G}(\xi);\mathbb{Z})$ is a fundamental invariant of $B\mathcal{G}(\xi)$, and it is an important question which elements in $H^*(B\mathcal{G}(\xi);\mathbb{Z})$ are the fundamental class of a smooth manifold $M \to B\mathcal{G}(\xi)$, i.e., for which classes [c] in $H^*(B\mathcal{G}(\xi);\mathbb{Z})$ is there an oriented, compact, smooth manifold M and a continuous map $f: M \to B\mathcal{G}(\xi)$ such that $f_*([M]) = [c]$ where [M] denotes the Poincaré dual of the fundamental class of M in $H_*(M;\mathbb{Z})$. This question is closely related to the Thom morphism

$$\tau \colon MU^*(B\mathcal{G}(\xi)) \longrightarrow H^*(B\mathcal{G}(\xi);\mathbb{Z})$$

from complex cobordism to singular cohomology since complex cobordism classes over $B\mathcal{G}(\xi)$ can be represented by complex-oriented proper maps $f: M \to B\mathcal{G}(\xi)$, where M is a smooth manifold, and τ sends the cobordism class [f] to the cohomology class $f_*[M]$. In fact, the Thom morphism plays a key role in algebraic and geometric topology.

The purpose of the present paper is to show that the Thom morphism for the classifying space of certain gauge groups is not surjective and thereby to show that there is a restriction for how non-torsion classes can be represented by fundamental

²⁰²⁰ Mathematics Subject Classification. 57T10, 57R77, 55R35, 55S05, 55S10.

Key words and phrases. Complex cobordism, Lie groups, gauge groups, classifying spaces.

The second-named author was partially supported by the RCN Project No. 313472.

classes.

The cohomology of $B\mathcal{G}(\xi)$ is closely related to that of BG, the classifying space of the Lie group G. The torsion in the cohomology of $B\mathcal{G}(\xi)$ has been studied in [9], [19] and [22], see also [12]. When the cohomology has torsion, the Thom morphism can be non-surjective. The question of when the Thom morphism is surjective for BG has previously been studied in for example [21] and [3] where the non-surjectivity of the Thom morphism for classifying spaces of certain Lie groups is used to detect new phenomena of the cycle map and Deligne cohomology in algebraic geometry.

In Section 2, we determine the image of the Thom morphism for the classifying spaces of SO(2n) as well as the exceptional Lie groups, with the exception of E_8 where we only provide a partial result. We note that partial results were already known for SO(4) and G_2 . Therefore, our results significantly extend the known cases by providing a complete list of which non-torsion generators are in the image for G_2 , F_4 , E_6 and E_7 . It turns out that the classifying space of E_7 behaves in a slightly different way compared to the other exceptional Lie groups. We do not know of a geometric explanation of this phenomenon. We note that some of our results could also have been deduced from the computations on *BP*-cohomology of Kono and Yagita in [14, Theorem 5.5]. In Section 3 we study the image of the Thom morphism for the classifying spaces of the gauge groups of principal SO(6)and E_7 -bundles over spheres. Our main result is the following:

Theorem 1.1. Let ξ be a principal E_7 -bundle over S^4 . Then there is a non-torsion generator in $H^4(B\mathcal{G}(\xi);\mathbb{Z})$ which is not in the image of the Thom morphism.

As explained in [21], the non-surjectivity of τ can be used to construct examples where the reduced Thom morphism

 $\bar{\tau}: MU^*(X \times B\mathbb{Z}/p) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^*(X \times B\mathbb{Z}/p;\mathbb{Z})$

is not injective. The non-injectivity of the reduced Thom morphism is crucial for the applications in algebraic geometry in [21]. In Section 4, we show that $\bar{\tau}$ is not injective when X is the classifying space of SO(n) with n even or an exceptional Lie group and p = 2. We note that the work of Kono and Yagita in [14] implies a stronger statement on the integral reduced Thom morphism for the classifying spaces of F_4 and E_6 . In Theorem 4.5, we provide a complete list of the simplest cases of non-injectivity of $\bar{\tau}$ for X a compact connected Lie group with simple Lie algebra.

Acknowledgement. We are grateful to Nobuaki Yagita and the anonymous referee for helpful comments and suggestions.

2. Classifying spaces of Lie groups

Let X denote a finite CW-complex. It is a well-known fact that all Steenrod operations of odd degree vanish on the image of $MU^*(X)$ in $H^*(X; \mathbb{Z}/p)$ for all prime numbers p (see for example [21, page 468], [3, Proposition 3.6], [4]). Hence, in order to show that an element $x \in H^*(X; \mathbb{Z})$ is not in the image of the Thom morphism, it suffices to find a Steenrod operation of odd degree which does not vanish on r(x) where $r: H^*(X; \mathbb{Z}) \to H^*(X; \mathbb{Z}/p)$ denotes the reduction map. We will now apply this observation to the cases where X is BSO(n) or the classifying space of an exceptional Lie group. We let $H^*_{\text{free}}(X;\mathbb{Z})$ denote the quotient of $H^*(X;\mathbb{Z})$ by the torsion subgroup. When there is no risk of confusion we often use the same notation for an element in $H^*(X;\mathbb{Z})$ and its image in $H^*_{\text{free}}(X;\mathbb{Z})$.

2.1. Special Orthogonal Groups. Recall from [18, Theorem III.5.16] that the free cohomology of BSO(n) is given by

$$H^*_{\text{free}}(BSO(n);\mathbb{Z}) \cong \begin{cases} \mathbb{Z}[e_4, e_8, \dots, e_{2n-2}], & n \text{ odd} \\ \mathbb{Z}[e_4, e_8, \dots, e_{2n-4}, \chi_n], & n \text{ even.} \end{cases}$$

Theorem 2.1. Let $n \ge 4$ be even and let $z \in H^n(BSO(n); \mathbb{Z})$ be a torsion class (or 0). Then $(\chi_n + z) \in H^n(BSO(n); \mathbb{Z})$ is not in the image of the Thom morphism.

Proof. From [18, Theorems III.3.19 and III.5.12] we know that the mod 2 cohomology of BSO(n) is given by

$$H^*(BSO(n); \mathbb{Z}/2) \cong \mathbb{Z}/2[y_2, y_3, \dots, y_n],$$

with

(1)
$$\operatorname{Sq}^{j}(y_{k}) = \sum_{i=0}^{j} \binom{k-i-1}{j-i} y_{k+j-i} y_{i}$$

Let r denote the mod 2-reduction map $H^*(BSO(n);\mathbb{Z}) \to H^*(BSO(n);\mathbb{Z}/2)$. Then

$$r(e_{4k}) = y_{2k}^2 + \text{(other terms)}$$
$$r(\chi_n) = y_n + \text{(other terms)},$$

since y_{2k}^2 and y_n are neither the image nor the source of a nontrivial Bockstein homomorphism. Furthermore, we have

$$Sq^{3}(y_{n}) = \sum_{i=0}^{3} {\binom{n-1-i}{3-i}} y_{n+3-i}y_{i} = y_{3}y_{n} \neq 0.$$

We will now show that no other term in $r(\chi_n + z)$ is mapped to y_3y_n by Sq³. It follows from equation (1) that the only other element of $H^{n-3}(BSO(n); \mathbb{Z}/2)$ which can map to y_3y_n under Sq³ is y_2y_{n-2} . For this element, the relevant part of the Bockstein diagram for BSO(n) is the following:





where there are no other nontrivial Bocksteins going into or out of any of these elements (see for example [7, Chapter 3E] and [11, Section 2.2] for the use of Bockstein cohomology). It follows that y_2y_{n-2} does not generate a nontrivial element of the Bockstein cohomology of BSO(n), and thus it is not one of the summands in $r(\chi_n + z)$. Thus, we have $\operatorname{Sq}^3(r(\chi_n + z)) \neq 0$. The statement then follows from the fact that all odd-degree elements of the Steenrod algebra vanish on the image of the Thom morphism.

2.2. Exceptional Groups. Recall from [18, Theorems VI.5.5 and VI.5.10] that the free cohomologies of the classifying spaces of the simply connected exceptional Lie groups are given by

$$H^*_{\text{free}}(BG_2; \mathbb{Z}) \cong \mathbb{Z}[e_4, e_{12}]$$

$$H^*_{\text{free}}(BF_4; \mathbb{Z}) \cong \mathbb{Z}[e_4, e_{12}, e_{16}, e_{24}]$$

$$H^*_{\text{free}}(BE_6; \mathbb{Z}) \cong \mathbb{Z}[e_4, e_{10}, e_{12}, e_{16}, e_{18}, e_{24}]$$

$$H^*_{\text{free}}(BE_7; \mathbb{Z}) \cong \mathbb{Z}[e_4, e_{12}, e_{16}, e_{20}, e_{24}, e_{28}, e_{36}]$$

$$H^*_{\text{free}}(BE_8; \mathbb{Z}) \cong \mathbb{Z}[e_4, e_{16}, e_{24}, e_{28}, e_{36}, e_{40}, e_{48}, e_{60}].$$

Theorem 2.2. Let $G = G_2$, F_4 or E_6 . Then the generator $e_4 \in H^*(BG; \mathbb{Z})$ is not in the image of the Thom morphism, while all other non-torsion generators are in the image.

Proof. The mod 2 cohomology of each of the classifying spaces is given by

$$H^*(BG_2; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_4, y_6, y_7, y_{10}]$$

$$H^*(BF_4; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_4, y_6, y_7, y_{16}, y_{24}]$$

$$H^*(BE_6; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_4, y_6, y_7, y_{10}, y_{18}, y_{32}, y_{34}, y_{48}]/I,$$

where I denotes the ideal given by

 $I = \langle y_7 y_{10}, y_7 y_{18}, y_7 y_{34}, y_{34}^2 + y_{18}^2 y_{32} + y_{10}^2 y_{48} + y_6 y_{10} y_{18} y_{34} + y_4 y_{10} y_{18}^3 + y_4 y_{10}^3 y_{34} \rangle$ see [18, Corollary VII.6.3 and Theorem VII.6.6] and [10, Theorem 1.1 and Proposition 5.1]. In each of the four cases, the reduction map $H^*(BG;\mathbb{Z}) \to H^*(BG;\mathbb{Z}/2)$ sends e_4 to y_4 , and in each case $\operatorname{Sq}^3(y_4) = y_7$. This proves the first claim since Sq^3 vanishes on the image of the Thom morphism as we pointed out at the beginning of Section 2. For the other generators, the claim follows from the fact that all differentials in the Atiyah–Hirzebruch spectral sequence vanish as explained, for example, in [11, Section 2.1] or [21, page 471]. To prove the latter assertion, we note first that all differentials in the Atiyah–Hirzebruch spectral sequence for the respective spaces are torsion, i.e., their images are contained in the subgroup of torsion elements. For BG_2 , the cohomology has only 2-torsion. Therefore, it suffices to show that all odd degree compositions of Steenrod squares vanish. This can be checked directly for e_{12} . For BF_4 and BE_6 , the respective cohomology groups contain both 2- and 3-torsion. The cohomology operations in $H^*(BF_4; \mathbb{Z}/3)$ and $H^*(BE_6; \mathbb{Z}/3)$ are all power operations for p = 3. The differentials in the Atiyah–Hirzebruch spectral sequence are compositions of Steenrod squares and power operations. We can then check directly that all such compositions which increase the cohomological degree by an odd number vanish on the respective generators. This proves the second claim and finishes the proof of the theorem.

As the following theorem shows, the situation is a bit different for the classifying space BE_7 . While in the previous cases only the generator in degree 4 is not hit by the Thom morphism, for E_7 there are several generators which are not in the image of τ .

Theorem 2.3. The generators $e_4, e_{16}, e_{24}, e_{28} \in H^*(BE_7; \mathbb{Z})$ are not in the image of the Thom morphism, and nor are the sums of any of these generators with a 2-torsion element in the same degree. The non-torsion generators e_{12} and e_{20} are in the image of the Thom morphism. For the generator $e_{36} \in H_{free}^{36}(BE_7; \mathbb{Z})$, there

exist lifts to $H^{36}(BE_7;\mathbb{Z})$ which are not in the image of the Thom morphism, while other lifts are in the image.

Proof. The mod 2 cohomology of BE_7 is given by

$$H^*(BE_7; \mathbb{Z}/2) \cong \mathbb{Z}/2[y_4, y_6, y_7, y_{10}, y_{11}, y_{18}, y_{19}, y_{34}, y_{35}, y_{64}, y_{66}, y_{67}, y_{96}, y_{112}]/J,$$

where J denotes the ideal given by

$$\begin{split} J = & \langle y_6 y_{11} + y_7 y_{10}, \ y_6 y_{19} + y_7 y_{18}, \ y_{10} y_{19} + y_{11} y_{18}, \ y_{11}^3 + y_7^2 y_{19}, \\ & y_6 y_{35} + y_7 y_{34}, \ y_{10} y_{35} + y_{11} y_{34}, \ y_{11} y_{19}^2, \ y_{18} y_{35} + y_{19} y_{34}, \\ & y_{19}^3, \ y_6 y_{67} + y_7 y_{66}, \ y_{10} y_{67} + y_{11} y_{66}, \ y_{18} y_{67} + y_{19} y_{66}, \\ & y_7^2 y_{67} + y_{11} y_{35}^2, \ y_{11}^2 y_{67} + y_{19} y_{35}^2, \ y_{34} y_{67} + y_{35} y_{66}, \ y_{19}^2 y_{67}, \\ & y_{66}^2 + y_{10}^2 y_{112} + y_{18}^2 y_{96}, \ y_{66} y_{67} + y_{10} y_{11} y_{112} + y_{18} y_{19} y_{96}, \\ & y_{34}^2 + y_{6}^4 y_{112} + y_{19}^2 y_{96}, \ y_{35}^2 y_{67} + y_7^2 y_{11} y_{112} + y_{11}^2 y_{19} y_{96}, \\ & y_{34}^3 y_{35} + y_6^3 y_7 y_{112} + y_{10}^3 y_{11} y_{96} + y_{18}^3 y_{19} y_{64}, \\ & y_{34}^2 y_{35}^2 + y_{6}^2 y_7^2 y_{112} + y_{10}^2 y_{11}^2 y_{96} + y_{18}^2 y_{19}^2 y_{64}, \\ & y_{34} y_{35}^3 + y_{6} y_7^2 y_{112} + y_7^2 y_{11} y_{18} y_{96}, \ y_{35}^4 + y_7^4 y_{112} + y_{11}^4 y_{96} \rangle, \end{split}$$

see [17, Theorem 4.4]. From [13, page 276 and Corollary 6.9] we then get that the Steenrod algebra acts on the following generators as

$$\begin{aligned} & \operatorname{Sq}^{3}(r(e_{4})) = \operatorname{Sq}^{3}(y_{4}) = y_{7} \\ & \operatorname{Sq}^{15}(r(e_{16})) = \operatorname{Sq}^{15}(y_{6}y_{10}) = y_{6}^{2}y_{19} + y_{10}^{2}y_{11} + y_{4}y_{7}y_{10}^{2} \\ & \operatorname{Sq}^{3}(r(e_{24})) = \operatorname{Sq}^{3}(y_{6}y_{18}) = y_{7}y_{10}^{2}. \end{aligned}$$

This implies, in particular, that these elements are not in the image of the Thom morphism, since all Steenrod operations of odd degree vanish on the image of the Thom morphism (see the beginning of Section 2 for references for this claim). By analysing the Bockstein homomorphisms, we see that an element of $H^{28}(BE_7;\mathbb{Z})$ which maps to the generator $e_{28} \in H^{28}_{\text{free}}(BE_7;\mathbb{Z})$ can be mapped to either $y_{10}y_{18}$ or $y_{10}y_{18} + y_7^4$ by r. Both of these elements are mapped to $y_{10}^2y_{11}$ by Sq³, so e_{28} plus torsion is not in the image of the Thom morphism either.

For the generators e_{12} and e_{20} , the claim follows again from the fact that all differentials in the Atiyah–Hirzebruch spectral sequence vanish as explained, for example, in [11, Section 2.1] or [21, page 471]. To prove the latter assertion, we use again that all differentials in the Atiyah–Hirzebruch spectral sequence for the respective spaces are torsion. The cohomology groups of BE_7 contain both 2- and 3-torsion. The cohomology operations in $H^*(BE_7; \mathbb{Z}/3)$ are all power operations for p = 3. The differentials in the Atiyah–Hirzebruch spectral sequence are thus compositions of Steenrod squares and power operations. We can then check directly that all such odd degree compositions vanish on the images of the generators e_{12} and e_{20} .

35	36	37
$y_4 y_7^3 y_{10}$ —	$\longrightarrow y_4 y_7^3 y_{11}$	$y_{18}y_{19}$
$y_7^2 y_{10} y_{11}$ —	$\longrightarrow y_7^2 y_{11}^2$	$y_4 y_7^2 y_{19}$
$y_4^2 y_6 y_7^3$	$\longrightarrow y_4^2 y_7^4$	$y_4^2 y_{11} y_{18}$
y_{35}	$y_7 y_{11} y_{18}$ —	$\longrightarrow y_7 y_{11} y_{19}$
$y_{4}^{4}y_{19}$	$y_4 y_{10} y_{11}^2$ —	$\longrightarrow y_4 y_{11}^3$
$y_7 y_{10} y_{18}$	$y_4 y_6^3 y_7^2$	$\longrightarrow y_4 y_6^2 y_7^3$
$y_4 y_6 y_7 y_{18}$	$y_4^4 y_6 y_7^2$ —	$\longrightarrow y_4^4 y_7^3$
$y_4 y_{10}^2 y_{11}$	y_{18}^2	$y_6^2 y_7 y_{18}$
$y_4^6 y_{11}$	$y_4^2 y_{10} y_{18}$	$y_4^3 y_7 y_{18}$
$y_4^2 y_7 y_{10}^2$	$y_6^3 y_{18}$	$y_4^2 y_7 y_{11}^2$
$y_6^3 y_7 y_{10}$	$y_4^3 y_6 y_{18}$	$y_4^4 y_{10} y_{11}$
$y_4^3 y_6 y_7 y_{10}$	$y_6 y_{10}^3$	$y_4^3 y_7^2 y_{11}$
$y_4 y_6^4 y_7$	$y_4^4 y_{10}^2$	$y_7 y_{10}^3$
$y_4^4 y_6^2 y_7$	$y_4 y_6^2 y_{10}^2$	$y_4 y_6 y_7 y_{10}^2$
$y_{4}^{7}y_{7}$	$y_4^2 y_6^3 y_{10}$	$y_6 y_7^3 y_{10}$
	$y_4^5 y_6 y_{10}$	$y_4^2 y_6^2 y_7 y_{10}$
	y_6^6	$y_4^5 y_7 y_{10}$
	$y_4^3 y_6^4$	$y_4^3 y_6^3 y_7$
	$y_{4}^{6}y_{6}^{2}$	$y_{6}^{5}y_{7}$
	y_4^9	$y_{4}^{6}y_{6}y_{7}$

Finally, we have the following Bockstein diagram in degree 36:

The diagram shows that an element of $H^{36}(BE_7; \mathbb{Z})$ corresponding to the generator $e_{36} \in H^{36}_{\text{free}}(BE_7; \mathbb{Z})$ is mapped to $y_{18}^2 + L$ by r, where L is some linear combination of the elements

$$y_4y_7^3y_{11}, y_7^2y_{11}^2, y_4^2y_7^4.$$

While all odd-degree elements of the Steenrod algebra act trivially on y_{18}^2 , $y_7^2 y_{11}^2$ and $y_4^2 y_7^4$, we have $\operatorname{Sq}^3(y_4 y_7^3 y_{11}) = y_7^4 y_{11}$. Thus, any lift of e_{36} which contains the term $y_4 y_7^2 y_{11}$ is not in the image of the Thom morphism, while any lift which does not contain that term is in the image. This proves the last claim and finishes the proof of the theorem.

Remark 2.4. As for the other exceptional Lie groups, the generator $e_4 \in H^4(BE_8; \mathbb{Z})$ is not in the image of the Thom morphism. This can for example be shown using the fact that BE_8 and the Eilenberg–MacLane space $K(\mathbb{Z}, 4)$ have homotopy equivalent 15-skeletons [8, page 185]. However, since the mod 2 cohomology of BE_8 is not known, we cannot give a complete answer to which other generators are in the image of the Thom morphism.

Remark 2.5. We note that some of our results could have been deduced from the computations of *BP*-cohomology of Kono and Yagita. Moreover, it follows from

[14, Theorem 5.5] that e_4 in $H^4(BG;\mathbb{Z})$ is not in the image of τ for G = SO(4) whereas $2e_4$ is in the image.

3. Gauge Groups

Let X be a paracompact space, and let ξ be a principal G-bundle over X. The bundle ξ is classified by a map $f: X \to BG$. Let $\operatorname{Map}_f(X, BG)$ denote the path component of $\operatorname{Map}(X, BG)$ which contains the map f. By [2, Proposition 2.4], $B\mathcal{G}(\xi)$ equals $\operatorname{Map}_f(X, BG)$ in homotopy theory, and we will consider $\operatorname{Map}_f(X, BG)$ as a model for $B\mathcal{G}(\xi)$. For $X = S^n$, there is a fibre sequence

(2)
$$\Omega_f^n(BG) \longrightarrow \operatorname{Map}_f(S^n, BG) \xrightarrow{\operatorname{ev}} BG,$$

where $\Omega_f^n(BG)$ denotes the path component of $\Omega^n(BG)$ which contains f and ev denotes the evaluation map at the basepoint of S^n . Note that maps in Map (S^n, BG) are not required to be pointed, while maps in $\Omega^n(BG)$ are. We further note that $\Omega BG \cong G$ and that all path-components of $\Omega^{n-1}(G)$ are homotopy equivalent. Therefore, sequence (2) simplifies to

(3)
$$\Omega_0^{n-1}(G) \longrightarrow \operatorname{Map}_f(S^n, BG) \xrightarrow{\operatorname{ev}} BG,$$

where $\Omega_0^{n-1}(G)$ denotes the path-component of $\Omega^{n-1}(G)$ which contains the constant map. Since there is a cross-section $s \colon BG \to \operatorname{Map}_f(S^n, BG)$ such that $\operatorname{ev} \circ s$ is the identity, the Thom morphism is easily seen to be non-surjective for $\operatorname{Map}_f(S^n, BG)$ if it is non-surjective for BG. When the Thom morphism is not surjective for BG, however, then the question of surjectivity of the Thom morphism for $\operatorname{Map}_f(S^n, BG)$ is more interesting. First we need the following lemma.

Lemma 3.1. The cohomology group $\widetilde{H}^k(\Omega_0^3(E_7); \mathbb{Z}/2)$ is trivial for $k \leq 7$.

Proof. Let Q_i denote the *i*th Dyer–Lashof operation [1, Definition 4.1]

$$Q_i \colon H_k(\Omega^n X; \mathbb{Z}/2) \longrightarrow H_{2k+i}(\Omega^n X; \mathbb{Z}/2)$$

for $0 \leq i \leq n-1$, and let $\beta: H_k(X, \mathbb{Z}/2) \to H_{k-1}(X; \mathbb{Z}/2)$ denote the Bockstein homomorphism. By [6, Theorem 3.15] there is an isomorphism

$$H_*(\Omega_0^3(E_7); \mathbb{Z}/2) \cong \mathbb{Z}/2[Q_1^a \beta(u_{30})] \otimes \mathbb{Z}/2[Q_1^a Q_2^b(u_{30})] \otimes \left(\bigotimes_{k \in J} H_*(\Omega^3(S^k); \mathbb{Z}/2)\right),$$

where $J = \{11, 15, 19, 23, 27, 35\}$ and where *a* and *b* range over all non-negative integers. Since the reduced homology of the triple loop space of a sphere of dimension at least 11 is concentrated in degrees ≥ 8 (see [5, page 74]), we get that $\widetilde{H}_k(\Omega_0^3(E_7); \mathbb{Z}/2)$ is trivial for $k \leq 7$. By the universal coefficient theorem the same holds for $\widetilde{H}^k(\Omega_0^3(E_7); \mathbb{Z}/2)$.

We now prove our main result.

Theorem 3.2. Let ξ be a principal E_7 -bundle over S^4 . Then the image of $ev^*(e_4)$ in $H^4(B\mathcal{G}(\xi);\mathbb{Z})$ is not in the image of the Thom morphism.

Proof. Due to the homotopy equivalence $B\mathcal{G}(\xi) \simeq \operatorname{Map}_f(S^4, BE_7)$, it suffices to show that $\operatorname{ev}^*(e_4) \in H^4(\operatorname{Map}_f(S^4, BE_7); \mathbb{Z})$ is not in the image of the Thom morphism. We consider the mod 2 cohomology Serre spectral sequence of the fibre sequence

$$\Omega_0^3(E_7) \longleftrightarrow \operatorname{Map}_f(S^4, BE_7) \xrightarrow{\operatorname{ev}} BE_7.$$

The E_2 -page is given by

$$E_2^{p,q} = H^p(BE_7; \mathbb{Z}/2) \otimes H^q(\Omega_0^3(E_7); \mathbb{Z}/2)$$

By Lemma 3.1, the element $y_7 \in H^7(BE_7; \mathbb{Z}/2) \cong E_2^{7,0}$ is not in the image of any nontrivial differentials for degree reasons. Thus, $ev^*(y_7) \in H^7(\operatorname{Map}_f(S^4, BE_7); \mathbb{Z}/2)$ is nonzero. Furthermore, the proof of Theorem 2.3 shows that $\operatorname{Sq}^3(r(e_4)) = y_7$. The commutative diagram

$$\begin{array}{c|c} H^4(BE_7;\mathbb{Z}) & \stackrel{\mathrm{ev}^*}{\longrightarrow} & H^4(\mathrm{Map}_f(S^4, BE_7);\mathbb{Z}) \\ & & & & \downarrow^r \\ H^4(BE_7;\mathbb{Z}/2) & \stackrel{\mathrm{ev}^*}{\longrightarrow} & H^4(\mathrm{Map}_f(S^4, BE_7);\mathbb{Z}/2) \\ & & & & \downarrow^{\mathrm{Sq}^3} \\ H^7(BE_7;\mathbb{Z}/2) & \stackrel{\mathrm{ev}^*}{\longrightarrow} & H^7(\mathrm{Map}_f(S^4, BE_7);\mathbb{Z}/2) \end{array}$$

then shows that $\operatorname{Sq}^3 \circ r$ acts non-trivially on $\operatorname{ev}^*(e_4)$, which completes the proof. \Box

Remark 3.3. We can similarly examine whether the Thom morphism is surjective for principal SO(n)-bundles. Let ξ denote a principal SO(n)-bundle over S^2 where the classifying map f is not homotopic to the constant map. It was shown in [19, Theorem 1.2] that $B\mathcal{G}(\xi)$ has torsion if and only if $n \geq 5$. Since we found in Theorem 2.1 that the Thom morphism is non-surjective for BSO(n) when n is even, a natural candidate for a gauge group whose classifying space has a non-surjective Thom morphism is ξ as above when n = 6. Since $\Omega_f^2(BSO(6)) \cong \Omega_0(SO(6)) \cong$ $\Omega(\text{Spin}(6))$, there is a fiber sequence

(4)
$$\Omega(\operatorname{Spin}(6)) \longrightarrow \operatorname{Map}_f(S^2, BSO(6)) \xrightarrow{\operatorname{ev}} BSO(6).$$

By [5, Lemma 2.2], we know that

$$H^*(\Omega(\operatorname{Spin}(6)); \mathbb{Z}/2) \cong \mathbb{Z}/2[a_2]/(a_2^4) \otimes \Gamma[c_6] \otimes \left(\bigotimes_{i=0}^{\infty} \mathbb{Z}/2[\gamma_i(b_4)]/(\gamma_i(b_4)^4)\right),$$

where $\Gamma[x]$ denotes the divided power algebra generated by $\{\gamma_0(x), \gamma_1(x), \ldots\}$. It is then possible that the differential $d_9: H^8(\Omega(\text{Spin}(6)); \mathbb{Z}/2) \longrightarrow H^9(BSO(6); \mathbb{Z}/2)$ in the Serre spectral sequence of sequence (4) maps $\gamma_1(b_4)$ to y_3y_6 , which would imply that the element y_3y_6 does not survive to the E_{∞} -page. However, we were unable to determine whether this differential is trivial or not and we therefore do not know whether the Thom morphism is surjective in this case.

4. Non-injectivity of the integral Thom Morphism

Let X be a CW-complex and let X_k denote the k-skeleton. Recall that $MU^*(X)$ is said to satisfy the Mittag-Leffler condition if for all $n \ge 0$ there exists some $m \ge n$ such that Im $(MU^*(X) \to MU^*(X_n)) = \text{Im}(MU^*(X_m) \to MU^*(X_n))$. It follows from the Milnor short exact sequence that if $MU^*(X)$ satisfies the Mittag-Leffler condition, then $MU^*(X) = \varprojlim MU^*(X_n)$. Letting MU^* act on \mathbb{Z} (or \mathbb{Z}/p) by having all generators in nonzero degree act trivially on \mathbb{Z} , we get a tensor product $MU^*(X) \otimes_{MU^*} \mathbb{Z}$. This is isomorphic to $MU^*(X)$ modulo an ideal contained in the kernel of the Thom morphism. Thus, the reduced Thom morphism

$$MU^*(X) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^*(X;\mathbb{Z})$$

is well-defined (see also [21, page 470]). The following lemma is a general version of [21, Corollary 5.3] which is formulated for classifying spaces of compact Lie groups only. The proof, however, is the same as in [21]. For completeness, we provide the reader with the full argument here since we will apply the assertion later in Theorem 4.5.

Lemma 4.1. Let X be a CW-complex of finite type such that $MU^*(X)$ satisfies the Mittag–Leffler condition. Let p be a prime, k an integer and $x \in H^k(X; \mathbb{Z})$. Suppose that the image of the Thom morphism $MU^k(X) \to H^k(X; \mathbb{Z})$ contains px but no element y such that py = px. Then the Thom morphism

$$MU^{k+2}(X \times B\mathbb{Z}/p) \otimes_{MU^*} \mathbb{Z} \to H^{k+2}(X \times B\mathbb{Z}/p;\mathbb{Z})$$

is not injective.

Proof. We first show that the map $MU^k(X) \otimes_{MU^*} \mathbb{Z}/p \to H^k(X; \mathbb{Z}/p)$ is not injective. Suppose that $\alpha \in MU^k(X)$ maps to px under the Thom morphism. If the element $(\alpha \otimes 1) \in MU^k(X) \otimes_{MU^*} \mathbb{Z}/p$ is zero, then $\alpha = p\beta$ for some $\beta \in MU^k(X)$. It follows that the Thom morphism maps β to some y with py = px, which contradicts the assumption. Therefore, $\alpha \otimes 1$ is nonzero. However, the Thom morphism maps $\alpha \otimes 1$ to $0 \in H^k(X; \mathbb{Z}/p)$. Thus $MU^k(X) \otimes_{MU^*} \mathbb{Z}/p \to H^k(X; \mathbb{Z}/p)$ is not injective.

For all finite CW-complexes Y, there is an isomorphism

$$MU^*(Y \times B\mathbb{Z}/p) \cong MU^*(Y) \otimes_{MU^*} MU^*(B\mathbb{Z}/p)$$

by [15, Theorem 2']. This implies

$$MU^*(X \times B\mathbb{Z}/p) \cong \varprojlim MU^*(X_n \times B\mathbb{Z}/p) \cong \varprojlim (MU^*(X_n) \otimes_{MU^*} MU^*(B\mathbb{Z}/p))$$
$$\cong MU^*(X) \otimes_{MU^*} MU^*(B\mathbb{Z}/p).$$

It then follows that

$$MU^{*}(X \times B\mathbb{Z}/p) \otimes_{MU^{*}} \mathbb{Z} \cong (MU^{*}(X) \otimes_{MU^{*}} MU^{*}(B\mathbb{Z}/p)) \otimes_{MU^{*}} \mathbb{Z}$$
$$\cong (MU^{*}(X) \otimes_{MU^{*}} \mathbb{Z}) \otimes_{MU^{*}} (MU^{*}(B\mathbb{Z}/p) \otimes_{\mathbb{Z}} \mathbb{Z})$$
$$\cong (MU^{*}(X) \otimes_{MU^{*}} \mathbb{Z}) \otimes_{\mathbb{Z}} (MU^{*}(B\mathbb{Z}/p) \otimes_{MU^{*}} \mathbb{Z})$$
$$\cong (MU^{*}(X) \otimes_{MU^{*}} \mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}[c]/(pc)),$$

where |c| = 2 and where the final isomorphism comes from the fact that the Thom morphism

$$MU^*(B\mathbb{Z}/p) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^*(B\mathbb{Z}/p;\mathbb{Z}) \cong \mathbb{Z}[c]/(pc)$$

is an isomorphism. Moreover, we get

$$(MU^*(X) \otimes_{MU^*} \mathbb{Z}) \otimes_{\mathbb{Z}} (\mathbb{Z}[c]/(pc)) \cong (MU^*(X) \otimes_{MU^*} \mathbb{Z})[c]/(pc)$$
$$\cong (MU^*(X) \otimes_{MU^*} \mathbb{Z}) \oplus \prod_{i=1}^{\infty} (MU^*(X) \otimes_{MU^*} \mathbb{Z}) c^i/(pc^i)$$
$$\cong (MU^*(X) \otimes_{MU^*} \mathbb{Z}) \oplus \prod_{i=1}^{\infty} (MU^*(X) \otimes_{MU^*} \mathbb{Z}/p) c^i.$$

Thus, the integral Thom morphism for the space $X\times B\mathbb{Z}/p$ is given by the composite map

Since $MU^k(X) \otimes_{MU^*} \mathbb{Z}/p \to H^k(X; \mathbb{Z}/p)$ is not injective, we get that the subgroup $(MU^k(X) \otimes_{MU^*} \mathbb{Z}/p)c$ maps noninjectively to $H^{k+2}(X \times B\mathbb{Z}/p)$, and the statement follows.

By combining the previous lemma with the results in Section 2, we get multiple examples of when the Thom morphism is not injective.

Theorem 4.2. Let $G = G_2$, F_4 , E_6 , E_7 or SO(n) with $n \ge 4$ even. Then the Thom morphism

$$MU^6(BG \times B\mathbb{Z}/2) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^6(BG \times B\mathbb{Z}/2;\mathbb{Z})$$

is not injective.

Proof. We first note that $H^4(BG; \mathbb{Z}) \cong \mathbb{Z}$, where e_4 denotes the canonical generator. By Theorems 2.1, 2.2 and 2.3 e_4 is not in the image of the Thom morphism. However, by [16, Theorem 1], $MU^*(BG)$ satisfies the Mittag–Leffler condition, from which it follows that some integer multiple ne_4 is in the image of the Thom morphism. Assume that n is minimal. Since Sq^3 acts non-trivially on e_4 , it follows that n is a multiple of 2. Setting p = 2, we then see that the conditions of Lemma 4.1 are satisfied, and the statement follows. In fact, assuming that $\alpha \in MU^4(BG)$ maps to ne_4 , we see that the element

$$(\alpha \otimes 1)c \in (MU^4(BG) \otimes_{MU^*} \mathbb{Z}/2)c \subseteq MU^6(BG \times B\mathbb{Z}/2)$$

is mapped to 0 in $H^k(BG \times B\mathbb{Z}/2;\mathbb{Z})$.

Remark 4.3. For E_7 , we note that also other generators can be used to construct non-trivial elements in the kernel by Theorem 2.3.

Remark 4.4. We note that, for some of the groups, the work of Kono and Yagita allows for a stronger statement. One can deduce from [14, pages 795-796] that the reduced Thom morphism $MU^*(BG) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^*(BG;\mathbb{Z})$ is not injective for $G = F_4$ and $G = E_6$.

If X is a Lie group, and not the classifying space, the Thom morphism is in many cases not injective. In the following theorem we give a complete list for the simplest cases where injectivity fails.

Theorem 4.5. Let G be a compact connected Lie group with simple Lie algebra. For an integer $n \ge 1$, let r denote the smallest natural number such that $2^r \mid n$. The Thom morphism

 $MU^k(G \times B\mathbb{Z}/p) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^k(G \times B\mathbb{Z}/p;\mathbb{Z})$

is not injective in the following cases:

Group	n	k	p
$\operatorname{Spin}(n)$	$n \ge 7$	5	2
SO(n)	$n \ge 5$	5	2
Ss(n)	$8 \mid n$	5	2
Ss(n)	$8 \nmid n$	9	2
PSO(n)	$8 \mid n$	5	2
PSO(n)	$8 \nmid n, n \ge 10$	9	2
PSp(n)	n even	$2^{r+1} + 1$	2
$SU(n)/\Gamma_l$	$4 \mid n, \ l \equiv 2 \pmod{4}$	$2^r + 1$	2
G_2		5	2
F_4		5	2 or 3
E_6		5	2 or 3
E_6/Γ_3		5	2
E_7		5,17	2
E_7/Γ_2		5	3
E_7/Γ_2		17	2
E_8		5, 17, 25, 29	2

Proof. This follows from [11, Theorem 1.1] and Lemma 4.1.

Remark 4.6. If $MU^*(B\mathcal{G}(\xi))$ satisfies the Mittag–Leffler condition for the bundle ξ of Theorem 3.2, then the Thom morphism

 $MU^6(B\mathcal{G}(\xi) \times B\mathbb{Z}/2) \otimes_{MU^*} \mathbb{Z} \longrightarrow H^6(B\mathcal{G}(\xi) \times B\mathbb{Z}/2;\mathbb{Z})$

is not injective. However, we were unable to determine whether this is the case.

References

- T. Kudo, S. Araki, Topology of H_n-spaces and H-squaring operations, Mem. Fac. Sci. Kyūsyū Univ. A10 (1956), 85–120.
- [2] M. F. Atiyah, R. Bott, The Yang-Mills equations over Riemann surfaces, Philos. Trans. Roy. Soc. London Ser. A308 (1983), no. 1505, 523–615.
- [3] O. Benoist, J.C. Ottem, Two coniveau filtrations, Duke Math. J. 170 (2021), 2719–2753.
- [4] H. Cartan, Séminaire Henri Cartan de l'Ecole Normale Supérieure, 1954/1955. Algèbres d'Eilenberg-MacLane et homotopie, Secrétariat Math., Paris, 1955.
- [5] Y. Choi, Homology of the double and triple loop space of SO(n), Math. Z. 222 (1996), no. 1, 59–80.

- [6] Y. Choi, S. Yoon, Homology of the double and the triple loop spaces of E₆, E₇, and E₈, Manuscripta Math. 103 (2000), no. 1, 101–116.
- [7] A. Hatcher, Algebraic topology, Cambridge University Press, Cambridge, 2002.
- [8] M. A. Hill, The String bordism of BE₈ and BE₈ × BE₈ through dimension 14, Illinois J. Math. 53 (2009), no. 1, 183–196.
- [9] M. Kameko, Torsion in classifying spaces of gauge groups, Proc. Math. Roy. Soc. Edinb. A, published online 2024:1–16.
- [10] M. Kameko, M. Nakagawa, T. Nishimoto, On the mod 2 cohomology of the classifying space of the exceptional Lie group E6, Proc. Japan Acad. Ser. A Math. Sci. 95 (2019), no. 9, 91–96.
- [11] E. Kaspersen, G. Quick, On the cokernel of the Thom morphism for compact Lie groups, Homology Homotopy Appl. 27 (2025), 323–353.
- [12] D.Kishimoto, S. Theriault, The mod-p homology of the classifying spaces of certain gauge groups, Proceedings of the Royal Society of Edinburgh, 153 (2023), 1805–1817.
- [13] A. Kono, M. Mimura, N. Shimada, On the cohomology mod 2 of the classifying space of the 1-connected exceptional Lie group E_7 , J. Pure Appl. Algebra, 8 (1976), no. 3, 267–283.
- [14] A. Kono, N. Yagita, Brown–Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups, Trans. AMS 339 (1993), 781–798.
- [15] P. S. Landweber, Coherence, flatness and cobordism of classifying spaces, Proc. Advanced Study Inst. on Algebraic Topology, Aarhus, 1970, Vol. II, pp. 256–269.
- [16] P. S. Landweber, Elements of infinite filtration in complex cobordism, Math. Scand. 30 (1972), 223–226.
- [17] M. Mimura, The characteristic classes for the exceptional Lie groups, Adams Memorial Symposium on Algebraic Topology, 1 (Manchester, 1990), 103–130. London Math. Soc. Lecture Note Ser., 175, Cambridge University Press, Cambridge, 1992.
- [18] M. Mimura, H. Toda, Topology of Lie groups. I and II, Translations of Mathematical Monographs, 91, American Mathematical Society, Providence, RI, 1991.
- [19] Y. Minowa, On the cohomology of the classifying spaces of SO(n)–gauge groups over $S^2,\,2023,\,\mathrm{arXiv}:2304.08702 \text{v}1.$
- [20] A. Ohsita, On the Stiefel–Whitney classes of the adjoint representation of E_8 , J. Math. Kyoto Univ. 44 (2004), no.3, 679–684.
- [21] B. Totaro, Torsion algebraic cycles and complex cobordism, J. Amer. Math. Soc. 10 (1997), no. 2, 467–493.
- [22] S. Tsukuda, On the cohomology of the classifying space of a certain gauge group, Proc. Roy. Soc. Edinburgh Sect. A127 (1997), no. 2, 407–409.

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU, NO-7491 TRONDHEIM, NORWAY *Email address*: eiolf.kaspersen@ntnu.no

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU, NO-7491 TRONDHEIM, NORWAY *Email address:* gereon.quick@ntnu.no